



15th International Symposium of Health Sciences

i-SIHAT 2026

in conjunction with the


**3RD AUAP INTERNATIONAL
HEALTH SCIENCE CONFERENCE**

**Advancing Health Frontiers:
Innovation and Global Partnerships**

ABSTRACT BOOK



23-24 JUNE 2026



**BERJAYA TIMES SQUARE HOTEL,
KUALA LUMPUR**



Invited Speaker

From Awareness To Empowerment: Reimagining Breast Health Education Through Human-Centric Artificial Intelligence

NORHAYATI MOHD ZAIN*

*Medical Imaging Department, School of Integrative Medicine & Life Sciences, KPJ Healthcare University, Nilai
71800, Negeri Sembilan, Malaysia*

**Correspondence: norhayati@kpju.edu.my*

ABSTRACT

Breast cancer remains one of the most prevalent health challenges affecting women worldwide, yet gaps in breast health literacy continue to hinder early detection and informed healthcare decision-making. In the era of Society 5.0, where technology is increasingly designed to serve human needs, artificial intelligence presents new opportunities to transform health education into a more accessible, engaging, and empowering experience. This plenary session introduces the Breast Imaging Education Bot (BieBot), a human-centric artificial intelligence (AI) initiative developed to enhance breast health literacy and promote proactive health behaviours among communities. Designed as a bilingual conversational chatbot, BieBot provides evidence-based information on breast cancer awareness, risk factors, screening options, and breast self-examination through an interactive and user-friendly digital platform. By integrating educational content, multimedia resources, instructional videos, and direct access to screening-related services, the initiative seeks to bridge knowledge gaps while improving access to reliable health information. The presentation will showcase the development journey of BieBot, highlighting how human-centred design principles, stakeholder engagement, and healthcare expertise were combined to create a meaningful digital health solution. Community evaluation findings reveal high levels of user acceptance, demonstrating the chatbot's effectiveness in supporting self-directed learning and increasing awareness of breast health practices. Furthermore, the initiative illustrates the potential of AI-powered health education to extend beyond awareness campaigns by fostering empowerment, informed decision-making, and greater participation in preventive healthcare. By sharing the BieBot experience, this session offers a practical example of how human-centric AI can be leveraged to create inclusive, scalable, and impactful healthcare innovations. The discussion will provide valuable insights for healthcare professionals, educators, researchers, and policymakers seeking to harness emerging technologies to improve health literacy and strengthen community wellbeing in the digital age.

Keywords: Breast Imaging Education Bot (BieBot); breast health literacy; conversational chatbot; human-centric AI

YI002

Caring for the Caregivers: The Evaluation of a Psychological Intervention Program for Caregivers of Childhood Cancer

AGNES CHONG SHU SZE^{1*}, MAHADIR AHMAD¹, HAMIDAH ALIAS²,
RIZUANA IQBAL HUSSAIN³, AFIFI LATEH⁴, CARYN MEI HSIEN CHAN¹

¹*Clinical Psychology and Behavioural Health Program/Center for Community Health Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Department of Paediatrics, Faculty of Medicine, Universiti Kebangsaan Malaysia, Faculty of Medicine, 56000 Cheras, Kuala Lumpur, Malaysia*

³*Department of Radiology, Faculty of Medicine Universiti Kebangsaan Malaysia, Faculty of Medicine, 56000 Cheras, Kuala Lumpur, Malaysia*

⁴*Department of Educational Evaluation and Research, Faculty of Education, Prince of Songkla University, Pattani Campus, Amphoe Mueang Pattani, Pattani 94000, Thailand*

*Correspondence: agnes@ukm.edu.my

ABSTRACT

Caregivers of children with acute lymphoblastic leukaemia (ALL) often report high levels of psychological distress impacting their psychological well-being and indirectly, their child's outcomes. Despite this, interventions targeting caregiver mental health remain limited. Therefore, the aim of this study was to evaluate a psychological intervention program designed to address traumatic stress, depression, and anxiety among caregivers. In this randomised controlled trial, 59 caregivers were assigned to either the intervention group (n = 29) or treatment-as-usual control group (n = 30) using the sequentially numbered, opaque, sealed envelopes method. The intervention consisted of a 2-week psychosocial self-help guidebook. Psychological outcomes were assessed at baseline, post-intervention, and 1-month follow-up using validated measures namely the Posttraumatic Stress Disorder Checklist, Beck Depression Inventory and Beck Anxiety Inventory. Results indicated a statistically significant reduction in post-traumatic stress symptoms in the intervention group at post-intervention ($F(1, 57) = 5.760, p = 0.020, \eta^2 = 0.093$). While there was no statistical difference in terms of depression and anxiety symptoms, a close to medium effect size was observed at 1-month follow-up. The psychosocial module developed and tested in this study shows potential to improve traumatic stress and depressive symptoms among caregivers of children with ALL, and suggests potential to benefit caregivers with higher baseline distress.

Keywords: Anxiety; cancer; caregiver; depression; traumatic stress

YI004

Clinician-Informed Usability and Design Guidelines for Home-Based Rehabilitation Mobile Applications

NGAN KUEN LAI^{1*}, GLORIA YING LE CHONG^{1,2}, TS. MUHAMMAD NUR AFFENDY NOR'A¹, JEFFERY JESELEE SIJORE¹, WEI YANG CHEN¹, TIMOTHY TSIN JIEN CHENG³

¹*School of Computing, Asia Pacific University of Technology & Innovation, 57000 Kuala Lumpur, Malaysia*

²*Johor Matriculation College, 84900 Tangkak, Johor, Malaysia*

³*Department of Orthopaedic, Hospital Duchess of Kent, 90000 Sandakan, Sabah, Malaysia*

**Correspondence: ngan.kuen@apu.edu.my*

ABSTRACT

Patients recovering from trauma injuries require continuous rehabilitation; however, conventional rehabilitation often relies on passive instructions, standardised protocols, and limited interactivity, making it less suitable for home-based settings. Although home-based rehabilitation mobile applications exist, many lack rigorous investigative foundations, clinician input, and affordability considerations. This study aims to derive clinician-validated usability and design guidelines for an interactive home-based rehabilitation mobile application. A qualitative approach was employed using face-to-face semi-structured interviews with 23 healthcare professionals from two Malaysian hospitals conducted over five months. A storyboard prototype was presented to facilitate discussion. Interviews were transcribed verbatim and analysed using NVivo to identify emerging themes based on thematic coding frequency, with data saturation achieved at 23 participants. Findings indicate that System Functionalities and Content (29%) were the most critical requirements, followed by Usefulness (21%), Positive User Experience (16%), Accessibility (14%), Affordability (12%) and Ease of Use (8%). Participants emphasised features such as posture guidance, strength monitoring, motivational feedback, and exercise tracking to support functional recovery and pain reduction. The study provides clinician-informed design guidelines for developing clinically reliable, user-friendly, affordable, and context-sensitive home rehabilitation applications tailored to Malaysia's linguistic, cultural, and infrastructural environment. The findings are limited to clinician perspectives within two hospitals and may not fully capture patient-specific usability challenges.

Keywords: Home-based rehabilitation; mobile health; patient-centred care; rehabilitation technology; usability design

Acknowledgement: This study is funded by the Asia Pacific University of Technology and Innovation Research Development Internal Grant (RDIG/07/2024).

YI005

Effectiveness of a Structured Physiotherapy Intervention on Motor Performance and Behavioural Outcomes in Children with Autism Spectrum Disorder: A Randomised Controlled Trial

NAZURAH ALWI¹, ASFARINA ZANUDIN^{1*}, MAHADIR AHMAD²

¹*Physiotherapy Programme, Centre for Rehabilitation and Special Needs Studies, Faculty of Health Science, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Clinical Psychology Programme, Centre for Community Health Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

**Correspondence: asfarina.zanudin@ukm.edu.my*

ABSTRACT

Motor impairments and behavioral difficulties are highly prevalent among children with autism spectrum disorder (ASD), significantly limiting functional performance and participation. Although exercise-based interventions have demonstrated benefits, many are implemented as recreational or activity-based programs rather than being systematically structured within a therapeutic framework. Consequently, these approaches remain heterogeneous and lack structured physiotherapy interventions grounded in clinical principles, limiting reproducibility and clinical translation. Therefore, this study aimed to evaluate the effectiveness of a structured physiotherapy intervention on motor performance and behavioral outcomes among children with mild-to-moderate ASD. A single-blinded, parallel randomised controlled trial included 64 children aged 6–10 years with mild-to-moderate ASD, classified using the Gilliam Autism Rating Scale, Third Edition (GARS-3). Participants were randomly allocated (1:1) to either the intervention or control group. Both groups received standardised usual care therapy. In addition, the intervention group underwent a structured physiotherapy program integrating coordination, balance, strength, and endurance training, delivered twice weekly in 60-minute sessions over 12 weeks, guided by progressive training principles. Outcomes were assessed at baseline and post-intervention using the Bruininks–Oseretsky Test of Motor Proficiency, Second Edition Short Form (BOT-2 SF) for motor performance and the Child Behavior Checklist (CBCL) for behavioral outcomes. Data were analysed using a mixed-design ANOVA. The intervention group demonstrated significantly greater improvements compared to the control group. Motor performance improved across key domains, including balance and coordination, with statistically significant group × time interaction effects ($p < 0.05$). Behavioral outcomes also showed significant reductions in behavioral problems ($p < 0.05$), indicating meaningful functional gains. In conclusion, a structured, progressive physiotherapy intervention significantly enhances both motor performance and behavioral outcomes in children with ASD. These findings provide evidence supporting the integration of structured physiotherapy interventions within multidisciplinary rehabilitation to improve functional and behavioral outcomes.

Keywords: Autism spectrum disorder; behavioral outcomes; motor performance; physiotherapy

YI006

Mapping Theory-Based Game Mechanics of “Wabak X” to Empower Outbreak Preparedness among the Orang Asli

AMEERAH SU’AD ABDUL SHAKOR^{1*}, MARIAM MOHAMAD¹, KHALID IBRAHIM¹, IZANDIS MOHAMAD SAYED²

¹*Department of Public Health Medicine, Faculty of Medicine, Universiti Teknologi MARA, 47000 Sungai Buloh, Selangor, Malaysia*

²*Hospital Orang Asli, Gombak, 53100 Kuala Lumpur, Selangor, Malaysia*

**Correspondence: ameerahsuad@moh.gov.my*

ABSTRACT

Orang Asli communities are highly vulnerable to outbreaks, often resulting in severe outcomes including death. Low health awareness, socioeconomic challenges, and reliance on deforested environments increase their exposure to emerging pathogens, including Disease X. Despite low levels of preparedness, they also show limited interest in current preparedness initiatives. This highlights the need for innovative, engaging, and culturally appropriate strategies to address this gap. This study presents “Wabak X”, a card-based serious game that applies game-based learning to deliver outbreak preparedness education through theory-aligned gameplay. The game is structured around the Health Belief Model (HBM) to achieve five learning objectives: helping players remember items needed in an outbreak preparedness kit; raising awareness that Disease X and outbreaks can occur at any time; informing players that animals can transmit Disease X to humans; highlighting that infectious diseases can spread through close contact; and demonstrating that preparedness can protect against outbreaks. Stakeholder input, including from the Department of Orang Asli Development and community members, informed culturally appropriate visual elements, while veterinary and public health experts guided the appropriate depiction of animals and disease transmission. To ensure that learning is embedded within player interactions, game mechanics are explicitly aligned with these objectives and mapped to key HBM constructs: perceived susceptibility, perceived severity, perceived benefits, and self-efficacy. The game design comprises four card types: item, preparedness, event, and action cards. Item and preparedness cards work together through a kit-building mechanic to strengthen self-efficacy and perceived benefits. Event cards simulate unpredictable zoonotic outbreaks and close-contact transmission, reinforcing perceived susceptibility and severity. Action cards regulate player interactions by triggering or mitigating events, illustrating disease dynamics while reinforcing the value of preventive behaviors. In conclusion, “Wabak X” translates complex outbreak concepts into accessible and practical knowledge, demonstrating how theory-driven game mechanics can support community-level preparedness.

Keywords: Game-based learning; health education; indigenous health; outbreak preparedness

YI008

Muscle Strength Following Critical Illness: The Relative Role of Protein Delivery and Organ Dysfunction Trajectory

NUR SHAZLEEN NISHA ROZALI, NURANIZA AZAHARI*

Department of Nutrition Sciences, Kuliyah of Allied Health Sciences, International Islamic University

Malaysia, 25200 Kuantan, Pahang, Malaysia

**Correspondence: nuraniza@iiu.edu.my*

ABSTRACT

Muscle weakness is a prevalent long-term consequence of critical illness and is associated with slow recovery and physical activity limitations. While protein provision remains foundational for muscle mass preservation, illness severity may exert an independent effect on functional muscle strength. This study aimed to assess the association between demographics and nutritional factors with handgrip strength in critically ill patients prior to intensive care unit discharge, and to determine whether illness severity provides additional explanatory value. A prospective observational cohort study was conducted among adult mechanically ventilated intensive care unit (ICU) patients. Average protein adequacy via enteral and/or parenteral nutrition was calculated as the percentage of protein delivered over the protein target. Daily Sequential Organ Failure Assessment (SOFA) score was utilised to represent the trajectory of illness severity. Patients performed handgrip strength assessment prior to ICU discharge, after weaning from mechanical ventilation. To assess the independent contribution of protein adequacy and admission SOFA score onto muscle strength, a hierarchical linear regression was conducted with adjustment for age and sex. In the final model, 65 patients with a mean age of 63 years (IQR 55.8 – 70.0) were included. Sex ($B = -0.38$, $p = 0.001$), average protein adequacy ($B = -0.28$, $p = 0.013$), and admission SOFA score ($B = -0.25$, $p = 0.040$) were associated with handgrip muscle strength and altogether explained 29.4% variance in the final model. The hierarchical linear regression analysis revealed that the average protein adequacy and admission SOFA score significantly explained an additional 7.2% and 5.1% of the variance in handgrip strength respectively. Men, lower average protein adequacy, and lower organ failure score were significantly associated with higher handgrip muscle strength. Admission SOFA score was independently associated with handgrip strength, even after controlling for demographic and nutritional factors. This factor should be considered alongside individualised nutrition strategies for the critically ill patients.

Keywords: Critical illness; hand strength; nutritional support; organ dysfunction scores

YI009

Knowledge, Attitude and Practice of Healthcare Providers Toward Multidomain Lifestyle Intervention for Dementia Prevention: A Preliminary Needs Assessment

SYAFIY A.AZMAN*, SUZANA SHAHAR, DEVINDER KAUR AJIT SINGH,
ARIMI FITRI MAT LUDIN, PONNUSAMY SUBRAMANIAM, NURUL
HIDAYAH MD FADZIL

*Center for Healthy Aging and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300
Kuala Lumpur, Malaysia*

**Correspondence: syafiy.azman03@gmail.com*

ABSTRACT

Multidomain lifestyle interventions have increasingly been recognised as an effective approach for dementia prevention by addressing modifiable risk factors across physical exercise, cognitive, nutrition, vascular and psychosocial domains. As part of an ongoing research project titled “Implementation and Integration of a Multidomain Lifestyle Intervention for Dementia Prevention Framework in Healthcare Settings”, Phase 1 focuses on adapting, harmonising and preparing the intervention module for implementation through structured training of healthcare professionals. These preliminary needs assessment aimed to explore the knowledge, attitude, and practice (KAP) of healthcare providers regarding the delivery of a multidomain lifestyle intervention for older adults. A pre- and post-workshop survey was conducted among seven healthcare providers from a healthcare institution selected for future program implementation. Participants were recruited using purposive sampling and completed a self-designed questionnaire before and after the training to assess sociodemographic characteristics and self-confidence related to delivering intervention components including exercise, cognitive stimulation, nutrition, vascular risk management and psychosocial engagement. Descriptive statistics were used to summarise participant characteristics and scores, while paired sample t-tests were used to examine changes following the training. All participants were female healthcare professionals from multidisciplinary backgrounds. The mean confidence score increased from 3.49 ± 0.66 at baseline to 4.53 ± 0.50 following the workshop. A statistically significant improvement was observed after the training ($t(6) = 7.92, p < 0.001$), indicating increased readiness among participants to deliver the multidomain intervention. Overall, the findings suggest that healthcare providers demonstrated acceptable knowledge and confidence following the training. These results support the feasibility of implementing the multidomain lifestyle intervention in healthcare settings and highlight the importance of capacity building among healthcare professionals in preparation for the subsequent phases of the ongoing implementation study.

Keywords: Dementia prevention; healthy ageing; healthcare providers; multidomain intervention; training of trainers

YI010

Combination Exercise Preserves Muscle Mass and Strength While Reducing Adiposity in a Neurodegenerative Rat Model: Preliminary Findings

NUR AISYAH ANUAR^{1,2*}, ARIMI FITRI MAT LUDIN¹, MUHAMMAD HAFIZ ZUHDI FAIROF^{1,2}, AMIRUL HAFIZ AHMAD ABDULLAH^{2,3}, THENG CHOON OOI⁴, FARAH WAHIDA IBRAHIM^{2,3}, NOR FADILAH RAJAB^{1,2}

¹Centre for Healthy Ageing and Wellness, Faculty of Health Sciences, 50300 Kuala Lumpur, Wilayah Persekutuan, Malaysia

²Biomedical Science Program, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Wilayah Persekutuan, Malaysia

³Center for Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

⁴Premier Integrated Labs Sdn. Bhd, Kuala Lumpur 55100, Malaysia

*Correspondence: p153408@siswa.ukm.edu.my

ABSTRACT

Neurodegeneration leads to progressive declines in muscle function, while increased visceral adiposity has been associated with impaired brain health. Aerobic and resistance exercise selectively improve metabolic and neuromuscular outcomes, respectively, reducing adiposity and enhancing muscle function. This suggests that combination exercise may be a promising strategy to simultaneously target multiple pathways affected in neurodegeneration. This study aimed to investigate the effects of an alternate-day combination exercise protocol on muscle and adiposity in a neurodegenerative rat model. Fifty male Wistar rats were assigned to Sham, Control, Aerobic+Al, Resistance+Al, or Combination+Al groups. Neurodegeneration was induced using 200 mg/kg body weight aluminium chloride (AlCl₃) for 8 weeks. Exercise began at week 4 until week 8, where aerobic (30 min/session, 5 days/week) and resistance exercise (14-16 reps/session, weekly load increment, 3 days/week) while alternate days for combined exercise (5 days/week). Grip strength was assessed at three time points and analysed using repeated-measures two-way ANOVA, whilst adiposity and muscle mass were measured post-intervention and analysed using two-way univariate ANOVA. Four weeks of aluminium exposure significantly reduced grip strength compared with baseline and controls (both $p < 0.001$). Exercise training improved grip strength across all modalities versus Control (all $p < 0.001$), with resistance training producing the greatest strength gains. However, modality-specific differences emerged in body composition. Resistance training induced divergent muscle adaptations, showing the lowest soleus but highest gastrocnemius mass ($p < 0.05$), and was associated with the highest adiposity among intervention groups ($p < 0.001$). In contrast, aerobic and combination training significantly reduced adiposity compared with Control, with aerobic showing the lowest fat accumulation. In a neurodegenerative rat model, the combination exercise protocol preserved neuromuscular function while maintaining lower adiposity, despite a total weekly dose

below standard exercise recommendations. Notably, the combination intervention outcomes consistently showed effects closer to aerobic exercise while not fully achieving resistance-specific benefits. These findings suggest that exercise distribution may be a key determinant of training efficacy and warrant further investigation of the exercise dose.

Keywords: Adiposity; combined exercise; neurodegeneration; resistance exercise; skeletal muscle

YI011

Breaking the Silence: From Nutrition Knowledge, Attitudes and Practices Assessment to the Development of a Sign Language Healthy Eating Video for Deaf Adults in Malaysia

YEE XING YOU^{1*}, FARISAH NAJWA ZAMAN², AIN NASUHA AZLAN²,
SYAR MEEZE MOHD RASHID³

¹*Centre of Healthy Ageing & Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Dietetics Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

³*Centre For Research Education and Community Wellness, Faculty of Education, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia*

*Correspondence: youyeexing@ukm.edu.my

ABSTRACT

In Malaysia, about 1.6 million adults (7.6%) have a hearing disability, according to the 2019 National Health and Morbidity Survey. Deaf adults often face challenges accessing nutrition education due to communication barriers and a lack of education materials in sign language. While studies abroad have found that people with hearing disabilities are more likely to be overweight or obese, however there is limited study in Malaysia. This study aimed to determine the nutrition-related knowledge, attitudes and practices (KAP) of healthy eating among deaf adults and subsequently develop and validate a sign language healthy eating video. This cross-sectional study involved 92 deaf adults from three community centres in Klang Valley, Negeri Sembilan and Malacca, chosen using purposive sampling. Sociodemographic characteristics and nutrition-related knowledge, attitudes and practices were collected using validated questionnaires. Height, weight and body mass index were measured. Based on the identified knowledge gaps, an educational video on healthy eating was developed using Malaysian Sign Language, visual graphics and captions to enhance comprehension among deaf audiences. Dietitians, nutritionists, and sign language interpreters reviewed the video to assess its content validity. The video's understandability and actionability were measured with the Patient Education Materials Assessment Tool Audiovisual (PEMAT-AV). Of those who took part, 29.3% were overweight and 25% were obese. Mean scores for nutrition-related KAP were 53.1 ± 17.5 , 4.2 ± 1.7 , and 7.4 ± 2.2 , respectively. The video achieved a content validity index of at least 0.83. PEMAT results showed high scores for understandability (92%) and actionability (88%), indicating that participants found the material clear and easy to act on. This project highlights the need for nutrition education that is accessible to deaf adults and introduces the first locally developed sign language healthy eating video in Malaysia, which could help make health promotion more inclusive.

Keywords: Audio-visual; deaf adults; healthy eating; nutrition education; sign language

YI013

Subclinical Cardiac Dysfunction in Young Adults with Hypertension: Insights from Speckle Tracking Echocardiography

LYANA SHAHIRAH MOHAMAD YAMIN^{1,2}, MARYAM ALSHARQI^{3,4},
MOHD ASYIQ AL-FARD MOHD RAFFALI⁵, AKMAL SABARUDIN¹,
NORLELA SUKOR⁵, IZA NURZAWANI CHE ISA¹, AFIFAH MOHAMED^{1*}

¹Centre for Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Centre for Medical Imaging Studies, Faculty of Health Sciences, Universiti Teknologi MARA, 42300 Puncak Alam, Selangor, Malaysia

³Institute for Medical Engineering and Science, Massachusetts Institute of Technology, 77 Massachusetts Ave., Cambridge, MA, 02139, USA

⁴Cardiac Technology Department, College of Applied Medical Sciences, Imam Abdulrahman Bin Faisal University, Dammam, Saudi Arabia

⁵Department of Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia

*Correspondence: afifahmohamed@ukm.edu.my

ABSTRACT

Young adults with hypertension represent a critical window for early cardiac intervention, yet subclinical myocardial dysfunction is frequently missed by conventional echocardiography. This cross-sectional study aimed to compare cardiac structure, systolic and diastolic function, and speckle tracking echocardiography (STE) strain parameters among 132 young adults aged 18–39. Participants were divided into three groups: normotensive controls (NT, n = 58), non-medicated hypertensives (HT-NonMeds, n = 37), and medicated hypertensives (HT-Meds, n = 37). Bonferroni correction was applied for post-hoc multiple comparisons. Both HT groups demonstrated concentric LV remodelling versus NT, with increased LV mass index (HT-NonMeds 86.27 ± 16.01 ; HT-Meds 92.76 ± 17.02 vs NT 71.53 ± 15.12 g/m²; both $p < 0.001$), relative wall thickness (HT-nM 0.48 ± 0.10 ; HT-M 0.51 ± 0.10 vs NT 0.41 ± 0.07 ; both $p < 0.001$), alongside diastolic dysfunction with progressively impaired septal e' velocity (NT 10.97 ± 1.97 , HT-NonMeds 9.28 ± 1.95 , HT-Meds 8.12 ± 1.97 cm/s; all $p < 0.001$) and elevated E/e' (both $p < 0.001$ vs NT). Despite preserved ejection fraction ($p = 0.067$), LVGLS was significantly impaired in HT-NonMeds (-16.74 [-18.72 to -14.14]) and HT-Meds (-16.70 [-18.67 to -14.90]) versus NT (-20.33 [-22.81 to -19.11]), both $p < 0.001$. LA conduit strain was reduced in both HT groups compared with NT (both $p < 0.003$), and LA reservoir strain was impaired only in HT-Meds ($p = 0.006$). HT-NonMeds and HT-Meds did not differ significantly for most parameters. Hierarchical regression identified hypertensive status ($\beta = 0.379$), BMI ($\beta = 0.313$), and female sex ($\beta = 0.253$) as independent predictors of LVGLS ($R^2 = 0.463$, $p < 0.001$). Adding septal e' velocity as a functional predictor further improved the model ($\Delta R^2 = 0.026$, $p = 0.013$), with the

final model explaining 48.9% of the variance in LVGLS. These findings demonstrate that STE reveals subclinical myocardial dysfunction in young hypertensive adults undetected by ejection fraction or diastolic grading, supporting its routine integration into echocardiographic surveillance for earlier risk stratification.

Keywords: Hypertension; global longitudinal strain; left atrial strain; speckle tracking echocardiography; young adults

YI014

Differential Topographical Mapping of the OFF Visual Pathway in Myopic and Emmetropic Peripheral Retinas

JIA YI CHONG¹, PUI JUAN WOI^{1*}, MOHD IZZUDDIN BIN HAIROL²

¹*Optometry and Vision Science Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Center for Community Health Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

**Correspondence: woipj@ukm.edu.my*

ABSTRACT

The ON and OFF visual pathways are specialised for processing light increments and decrements, respectively. While the OFF pathway has been linked to myopia progression, its functional distribution across the peripheral retina remains underexplored. This study compared the topographical performance of these pathways between myopic and emmetropic adults. A custom experimental paradigm was developed in PsychoPy to isolate and evaluate the reaction time and accuracy of the ON and OFF visual pathways. Participants included nine myopic and ten emmetropic (control group) adults aged 18 to 28 years. The experiment was conducted across seven horizontal retinal eccentricities (fovea to 30° nasal and temporal). A performance index was calculated as the product of accuracy and the inverse of reaction time. Both pathways showed a significant main effect of eccentricity ($p < 0.001$), with performance declining (lower accuracy and longer reaction time) toward the periphery. However, the OFF pathway exhibited greater robustness at higher eccentricities compared to the ON pathway. Crucially, a significant interaction was found between eccentricity and refractive status for the OFF pathway performance index ($p = 0.033$). This indicates that the functional efficiency of the OFF pathway across the peripheral retina follows a distinct profile in myopic individuals compared to emmetropic controls. The findings reveal that myopic eyes process peripheral OFF pathway information differently from emmetropic eyes. This topographical divergence suggests that the peripheral retina's sensitivity to dark-increment stimuli may be a factor in refractive development, providing a baseline for future targeted myopia control interventions.

Keywords: Myopia; ON-OFF visual pathways; psychophysics; retinal eccentricity

YI015

Associations between Donning and Doffing Difficulty, Body Mass Index and Wearing Comfort of Pressure Garments among Healthy Adults

JIA MIN LEE¹, SIAW CHUI CHAI^{1*}, KOK BENG GAN², SHIN YING CHU³,
KUICHENG LI⁴, AI LIAN LIM⁵

¹Center for Rehabilitation and Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Department of Electrical, Electronic & Systems Engineering, Faculty of Engineering and Built Environment, Universiti Kebangsaan Malaysia, 43600 Bangi, Malaysia

³Centre for Healthy Ageing and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

⁴School of Rehabilitation Medicine, Shandong Second Medical University, Weifang 261053, Shandong, China

⁵Grossman School of Medicine, New York University, New York, NY 10016, United States

*Correspondence: sc.chai@ukm.edu.my

ABSTRACT

Pressure garments are customised, tight-fitting garments commonly used in rehabilitation. Donning and doffing difficulty of pressure garments, body mass index (BMI) and wearing comfort of wearers may play important roles in ensuring therapeutic outcomes and wearers' adherence. This cross-sectional study established baseline utility of pressure garments by investigating their associations. Twenty healthy adult participants (mean age = 26.35 ± 1.26), fitted with a customised long-glove pressure garment with warp orientation, were asked to perform a functional task. A 0 - 10 numeric rating scale was used to measure donning and doffing difficulty, and wearing comfort. Difficulty scores between the donning and doffing phases were compared using a Wilcoxon Signed-Rank test. Associations were assessed using Spearman's rho (ρ) for donning, and Pearson's correlations (r) for doffing and BMI with bias-adjusted 95% confidence intervals (CI). Participants, with a mean BMI of $22.67 \pm 0.79 \text{ kg/m}^2$, reported a mean wearing comfort of 4.95 ± 0.37 . Donning (mean = 6.40 ± 0.55) was found to be significantly more difficult than doffing (mean = 3.90 ± 0.47) ($z = 2.92$; $p < 0.01$). Analysis revealed moderate negative associations between wearing comfort and BMI ($r = -0.38$; $p = 0.10$; 95%CI = $-0.70 - 0.09$) as well as donning difficulty ($\rho = -0.35$; $p = 0.13$; 95%CI = $-0.69 - 0.12$). Doffing difficulty exhibited a weaker negative association with wearing comfort ($r = -0.23$; $p = 0.34$; 95%CI = $-0.60 - 0.25$). Although the confidence intervals included zero, the consistent trends suggest that the initial donning phase and BMI were associated with poorer wearing comfort. Although the findings were non-significant, the results suggest that clinical focus should be placed on specialised donning techniques and garment design. Future large-scale studies are warranted to further confirm these variables as predictors of wearers' adherence in clinical populations.

Keywords: Body mass index; doffing difficulty; donning difficulty; pressure garment; wearing comfort

YI016

MINDCARE™: An Artificial Intelligence-Driven Digital Application Incorporating Malaysian-MIND (MY-MINDD®) Diet Scores for Predicting Mild Cognitive Impairment among Malaysian Older Adults

MUHAMAD MUSTAQIM M ZAPAWI¹, YEE XING YOU^{1*}, LAILATUL QADRI ZAKARIA², SHAHRUL AZMAN MOHD NOAH², PONNUSAMY SUBRAMANIAM³, MOHD RAZIF SHAHRIL⁴, SUZANA SHAHAR¹

¹Dietetic Program, Centre for Healthy Ageing and Wellness Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Centre for Artificial Intelligence Technology, Faculty of Information Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

³Clinical Psychology and Behavioral Health Program, Centre for Healthy Ageing and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

⁴Nutrition Program, Centre for Healthy Ageing and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

**Correspondence: youyeexing@ukm.edu.my*

ABSTRACT

Mild cognitive impairment (MCI) represents an intermediate stage between normal cognitive ageing and dementia, where early identification is crucial for timely preventive strategies. Advances in Artificial Intelligence (AI) provide opportunities to analyse complex interactions between multiple risk factors. Among these, adherence to the Malaysian adaptation of the Mediterranean-DASH Intervention for Neurodegenerative Delay (MIND) diet, known as Malaysian-MIND (MY-MINDD®) diet, has shown potential protective effects on cognitive health. This study aimed to develop an AI-driven digital application for MCI prediction by integrating MY-MINDD® scores among Malaysian older adults. Data from 1368 participants of the Long-Term Research Grant Scheme-Towards Useful Ageing (LRGS-TUA) cohort with five-year follow-up were analysed. Multiple machine learning algorithms were trained and evaluated using an 80:20 training-testing split, with model performance assessed using accuracy, area-under-curve (AUC), precision, sensitivity, specificity, and F1-score. Model interpretability was further examined using feature importance plots and Shapley Additive Explanations (SHAP). The Gradient Boosting Machine demonstrated the best predictive performance, achieving an accuracy of 91.97%, AUC of 0.9714, precision of 92.59%, sensitivity of 91.24%, specificity of 92.70%, and F1-score of 91.90%. MY-MINDD® diet scores emerged as the most influential predictor with an importance score of 0.4502. SHAP analysis indicated a transition threshold around scores of 5-6 where predicted MCI risk changes substantially. Following this model, an AI-driven digital application known as MINDCARE™ was developed and subsequently evaluated for usability among 105 older adults in both rural and urban settings using the Malay version of the mHealth Application Usability Questionnaire (M-MAUQ), yielding a high overall

usability score of 6.29 ± 0.80 out of a maximum score of seven. These findings demonstrated the integration of diet and AI for early MCI screening and highlight the potential of MINDCARE™ as a culturally relevant digital solution for clinical and community-based screening to support early intervention and preventive strategies among Malaysian older adults.

Keywords: Cognitive impairment; machine learning; Malaysian-MIND diet; older adult; prediction

YI017

Longitudinal Determinants of Intrinsic Capacity Trajectories Among Community-Dwelling Older Adults in Malaysia

NURUL HIDAYAH MD FADZIL, SUZANA SHAHAR, NURUL FATIN
MALEK RIVAN, DEVINDER KAUR AJIT SINGH*

Centre for Healthy Ageing and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia

**Correspondence: devinder@ukm.edu.my*

ABSTRACT

Intrinsic capacity (IC), defined as the composite of an individual's physical and mental abilities, plays a crucial role in sustaining independence in later life. This research aimed to identify factors associated with IC trajectories among community-dwelling Malaysian older adults. Data were gathered from the Long-term Research Grant Scheme Towards Useful Ageing (LRGS TUA) longitudinal cohort conducted between 2015 and 2020. As baseline, 2322 participants were recruited and 983 completed the five-year follow-up for trajectory analysis. Multinomial logistic regression was applied to determine associations between baseline determinants such as sociodemographic characteristics, biomarkers, body composition, physical fitness and cognitive outcome with IC trajectory categories (stable, improved or declined). Over 5-years period, almost half of the participants experienced a decline in IC (49.6%), 31.1% remained stable and 19.2% indicated improvement. The greatest deterioration was observed in the sensory domain, followed by cognitive, psychosocial, locomotor and vitality functions. No significant determinants were identified for IC improvement. However, IC decline was significantly associated with female gender [Odd ratio (OR) = 1.69, 95% Confidence Interval (CI): 1.28–2.24], older age (OR = 1.09, 95% CI: 1.06–1.12), older age (OR = 1.85, 95% CI: 1.33–2.57) and being single or divorced (OR = 1.85, 95% CI: 1.33–2.57). Lower educational attainment (OR = 0.941, 95% CI: 0.909–0.974) and poorer body composition indicators, including mid-upper arm circumference (MUAC) (OR = 0.942, 95% CI: 0.896–0.978), calf circumference (CC) (OR = 0.954, 95% CI: 0.918–0.992), and skeletal muscle mass (SMM) (OR = 0.940, 95% CI: 0.911–0.969) were also significant predictors. In addition, poorer cognitive performance (VR II) (OR = 0.961, 95% CI: 0.940–0.983), weaker handgrip strength (OR = 0.960, 95% CI: 0.942–0.979), and reduced functional status (OR = 0.904, 95% CI: 0.836–0.977) were linked to IC decline. These findings indicate that IC trajectories in Malaysian older adults are influenced by modifiable social, educational, and health factors. The results support implementing the World Health Organisation's Integrated Care for Older People (ICOPE) framework, with comprehensive IC assessment and targeted interventions to promote healthy ageing.

Keywords: Healthy ageing; ICOPE; intrinsic capacity; longitudinal cohort; older adults

YI018

From Regression to Machine Learning: Improving Prediction of Accelerated Ageing in Breast Cancer Survivors

NUR AFIFAH MURSYIDA ZAUJAN¹, MOHD RAZIF SHAHRIL^{1*}, SUZANA SHAHAR¹, EZANEE AZLINA MOHAMAD HANIF², NUR FA'IZAH AB MUJIN³, RAZINAH SHARIF¹, PONNUSAMY SUBRAMANIAM¹, AZIZI ABDULLAH⁴, MICHAEL FENECH⁵

¹Centre for Healthy Ageing and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300, Kuala Lumpur, Malaysia

²UKM Medical Molecular Biology Institute, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia

³Department of Radiotherapy and Oncology, Hospital Canselor Tuanku Muhriz, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia

⁴Centre for Artificial Intelligence Technology, Faculty of Information Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

⁵School of Agriculture, Food and Wine, College of Sciences, Adelaide University, Adelaide, SA, 5000 Australia

*Correspondence: raziifshahril@ukm.edu.my

ABSTRACT

Breast cancer survivors may experience accelerated ageing due to cancer pathology and treatment-related physiological stress. This study compared the predictive accuracy of a Multivariable Linear Regression (MLR) model and a machine learning approach to identify the optimal model for accelerated ageing prediction. A cross-sectional study was conducted among 262 breast cancer survivors and 262 cancer-free controls recruited through purposive sampling at Hospital Canselor Tuanku Muhriz, Universiti Kebangsaan Malaysia as part of Cancer Survivors' Trajectories of Ageing Research (C*STAR) project. Ageing phenotypes and behavioural determinants were assessed using validated instruments, including cognitive assessments, anthropometric measurements, physical fitness tests, the Pittsburgh Sleep Quality Index, the Instrumental Activities of Daily Living, the Food Frequency Questionnaire, the International Physical Activity Questionnaire, and the Perceived Stress Scale-10. Telomere length was used as the biomarker of accelerated ageing. For the MLR model, key assumptions were evaluated, including normality (histograms and P-P plots), multicollinearity (Tolerance and variance inflation factor), and linearity (scatterplots). For the machine learning approach, Extreme Gradient Boosting (XGBoost) models were developed using training, validation, and testing split (60:20:20). An XGBoost model integrated with the Synthetic Minority Over-sampling Technique for Regression with Gaussian Noise (SMO-GN) was further optimised using Particle Swarm Optimisation. Model performance was evaluated using Mean Absolute Error (MAE). The MLR model met the assumptions of normality and multicollinearity but violated the assumptions of linearity and homoscedasticity, indicating potential limitations in modelling complex non-linear relationships. The XGBoost-SMO-GN model demonstrated better predictive performance compared with both the MLR and standard XGBoost models in breast cancer survivors (MAE: 0.007 vs. 201.961).

vs. 0.1785) and controls (MAE: 0.0074 vs. 185.768 vs. 0.1592). In conclusion, the optimised XGBoost-SMOGN model showed better accuracy in predicting accelerated ageing, highlighting the utility of machine learning for identifying modifiable determinants to support precision healthy ageing strategies among breast cancer survivors.

Keywords: Age acceleration; ageing phenotypes; behavioural determinants of health; multivariable linear regression; machine learning

YI022

Artificial Intelligence and Mobile Health Technologies in the Assessment and Management of Voice Disorders: A Systematic Review

SAFWAN YUSOF^{1*}, ALIAA ARIFF²

¹*Center for Rehabilitation and Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan
Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Speech-Language Therapy Unit, Department of Otorhinolaryngology, Hospital Kuala Lumpur, 50586 Kuala
Lumpur, Malaysia*

**Correspondence: safwanyusof@ukm.edu.my*

ABSTRACT

Digital health technologies are transforming the assessment and rehabilitation of voice disorders by improving accessibility, remote monitoring, and personalised care. This systematic review evaluated the role of artificial intelligence (AI), mobile health (mHealth), and digital therapeutics in voice disorder management. This systematic review was conducted in accordance with PRISMA guidelines. A literature search identified 754 records from PubMed (n = 134), Scopus (n = 375), and Web of Science (n = 245). After duplicate removal and eligibility screening, 34 studies were included in the final analysis. AI and machine learning models, particularly Convolutional Neural Networks, Random Forests, and Support Vector Machines, demonstrated high accuracy in detecting, classifying, and grading vocal pathologies, often matching or surpassing clinical specialists. In contrast, generic Large Language Models showed limited reliability for auditory-perceptual voice assessment. mHealth applications enabled effective ambulatory monitoring and point-of-care screening through smartphone-based extraction of acoustic markers such as the Acoustic Voice Quality Index, Voice Wellness Index and Cepstral Peak Prominence, with strong agreement against gold-standard tools like Praat and studio microphones. However, performance declined in severely dysphonic voices. Mobile-based voice therapy improved adherence, self-efficacy, and vocal outcomes comparable to in-person rehabilitation. Serious games, covert public practice simulations, smartphone-adapted high-speed digital imaging, and computer vision tools further supported accessible therapy and objective vocal fold analysis. mHealth and AI technologies offer accessible, low-cost, and reliable solutions for voice screening, remote rehabilitation and longitudinal monitoring. These tools show strong clinical utility as adjuncts to, rather than replacements for, comprehensive clinical voice evaluation. Future work should improve accuracy for severe dysphonia, strengthen data privacy, and integrate multimodal clinical data for personalised voice care.

Keywords: Artificial intelligence; digital therapeutics; mobile health; voice disorders; voice rehabilitation

YI023

Monsoon Variability of Bacterial Communities in Airborne Particulate Matters of Kuala Lumpur, Malaysia

NURUL TASHA ZULKIFLE¹, HUI-MIN NEOH², MUHAMMAD IKRAM A WAHAB¹, SHARIFAH MAZRAH SAYED MOHAMED ZAIN³, ZURAHANIM FASHA ANUAL³, WEI-KANG LEE⁴, YEE JIA KEE⁴, SITI SHAHARA ZULFAKAR^{1*}

¹*Environmental Health and Industrial Safety Program, Centre for Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*UKM Medical Molecular Biology Institute, Universiti Kebangsaan Malaysia, Cheras, 56000 Kuala Lumpur, Malaysia*

³*Institute for Medical Research, National Institutes of Health, 40170 Shah Alam, Selangor, Malaysia*

⁴*Codon Genomics Sdn Bhd, Seri Kembangan, 43200 Selangor, Malaysia*

*Correspondence: sitishahara.zulfakar@ukm.edu.my

ABSTRACT

Airborne particulate matter (PM) carries diverse microbial communities that can influence atmospheric processes and public health. In Malaysia, monsoon-driven environmental changes may affect the distribution and composition of airborne bacteria, yet their relationships across monsoon cycles remain incompletely understood. This study investigated the bacterial community in PM across four monsoon periods, including Southwest Monsoon (SW), Intermonsoon II (IM II), Northeast Monsoon (NE) and Intermonsoon I (IM I). PM_{2.5} and PM₁₀ samples were collected using high- and low volume samplers followed by shotgun metagenomic sequencing to characterise taxonomic composition. The effects of monsoon-related environmental factors on bacterial community structure were also evaluated. Bacteria dominated the biological fraction of airborne PM. The bacterial community composition exhibited pronounced seasonal variation associated with monsoon periods, with *Bacillus* and *Staphylococcus* consistently dominating across all four monsoon phases and accounting for up to 90% of the relative abundance. Other genera, including *Pseudomonas*, *Ralstonia*, *Enterococcus* and *Clostridium* displayed seasonal variations, indicating environmental influences on airborne bacterial distribution. Strong correlations between environmental variables and bacterial composition were observed ($r > 0.80$, $p < 0.05$), indicating that environmental factors play a key role in shaping community structure. Overall, these findings demonstrate that monsoon-associated environmental factors significantly shape the composition and seasonal variation of airborne bacterial communities in Malaysia.

Keywords: Airborne particulate matter; bacterial community; environmental drivers; monsoon variability

YI024

Mosquito Magnet: An Innovative Solution to an Old Problem in Entomological Surveillance for Zoonotic Malaria

NANTHA KUMAR JEYAPRAKASAM^{1*}, INDRA VYTHILINGAM²,
SANDTHYA PRAMASIVAN², JONATHAN WEE KENT LIEW³, VAN LUN
LOW⁴, WAN-YUSOFF WAN-SULAIMAN²

¹Biomedical Science Programme, Center for Toxicology and Health Risk Studies, Faculty of Health Sciences,
Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Department of Parasitology, Faculty of Medicine, Universiti Malaya, 50603 Kuala Lumpur, Malaysia

³Environmental Health Institute, National Environment Agency, 138667 Singapore

⁴Tropical Infectious Diseases Research and Education Centre, University of Malaya, 50603 Kuala Lumpur,
Malaysia

*Correspondence: nanthakumar@ukm.edu.my

ABSTRACT

Vector surveillance is essential for determining the geographical distribution of mosquito vectors and for understanding the dynamics of malaria transmission. In Malaysia, while human-only malaria species have been largely eliminated, cases of zoonotic knowlesi malaria in humans continue to increase. Thus, it necessitates intensive vector studies using safer trapping methods that are both field-efficient and capable of attracting local vector populations. This study evaluated the ability of mosquito magnet to serve as an alternative tool for collecting *Anopheles* mosquitoes, vectors of zoonotic malaria. A randomised 4 x 4 Latin-square designed experiment was conducted to compare the efficiency of the mosquito magnet against human landing catch (HLC), Centers for Disease Control (CDC) light trap and human-baited trap (HBT). A Latin-square designed experiment showed that HLC caught the greatest number of *Anopheles* mosquitoes (n = 321) compared to HBT (n = 87), Mosquito Magnet (n = 58), and CDC light trap (n = 13). The Generalized Linear Mixed Model (GLMM) analysis showed that the HLC method caught significantly higher numbers of *Anopheles* mosquitoes compared to the mosquito magnet (p = 0.049). However, there was no significant difference in mean nightly catch of *Anopheles* mosquitoes between mosquito magnet and the other two trapping methods: HBT (p = 0.646) and CDC light traps (p = 0.197). On the other hand, the mean nightly catch for both *An. introlatus* (9.33 ± 4.341) and *An. cracens* (4.00 ± 2.273) caught using HLC was higher than those caught using mosquito magnet, though the differences were not statistically significant (p > 0.05). This is in contrast with the mean nightly catch of *An. sinensis* (15.75 ± 5.640) and *An. maculatus* (15.78 ± 3.479), where HLC showed a significantly higher number of mosquito catches compared to mosquito magnet (p < 0.05). Mosquito magnet has a promising ability to catch *An. introlatus* and *An. cracens*, the important vectors of knowlesi malaria in Peninsular Malaysia. Mosquito magnet's ability to catch some *Anopheles* species is comparable to HLC, making it an ethical and safer alternative.

Keywords: Malaria; mosquito magnet; surveillance; zoonotic

YI025

NMR-based Metabolomics Profiling and Bioactivity Evaluation of Selected Malaysian Herbs for Radical Scavenging and Senotherapeutic Properties

NOR MUHAMMAD HILMI HUSSIN¹, AHMED MEDIANI², MICHAEL FENECH³, RAZINAH SHARIF^{1*}

¹Centre for Healthy Aging and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Institute of Systems Biology, Universiti Kebangsaan Malaysia, Kuala Lumpur, 43600 Selangor, Malaysia

³Health and Biomedical Innovation, Clinical and Health Sciences, University of South Australia, 5000 Adelaide, Australia

*Correspondence: razinahsharif@ukm.edu.my

ABSTRACT

Inflammaging, a chronic, low-grade inflammation, is a shared mechanism of aging and age-related pathologies. Malaysian medicinal plants are traditionally known and used for their therapeutic effects to treat numerous diseases, including aging and related conditions. This study aimed to assess the biological activity of six selected Malaysian herbs with a focus on their antioxidant and anti-aging potential. Following this preliminary screening, we aimed to evaluate the metabolite variations of the three most active plants or herbs using proton nuclear magnetic resonance (1HNMR) and correlate them with radical scavenging activity and anti-aging properties. Ethanolic extracts (50%, 80% and 100%) of *Polygonum minus*, *Centella asiatica*, *Astragalus membranaceus*, *Cosmos caudatus*, *Momordica charantia* and *Morinda citrifolia* were screened for total phenolic content (TPC), DPPH radical scavenging, Ferric Reducing Antioxidant Power (FRAP) and 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulfonic acid (ABTS). The top three most active herbs were further analysed using 1H-NMR metabolomics coupled with multivariate data analysis (MVDA) to correlate specific metabolites with bioactivities. The 80% ethanolic extracts of *P. Minus*, *C. Caudatus* and *M. citrifolia* demonstrated the highest TPC, radical scavenging activity (DPPH and ABTS) and FRAP values, indicating robust antioxidant capacity. Metabolomics profiling using partial least squares (PLS) biplot revealed the presence of potent metabolites of these herbs and those responsible for antioxidant and anti-inflammatory or senomorphic effects such as quercetin, quercetin-3-O-rhamnoside (quercitrin), myricetin derivatives, catechin, isorhamnetin, astragaloside and apigenin. Eighty percent ethanolic extract of *P. minus*, *C. caudatus* and *M. citrifolia* can be considered promising sources of anti-aging ingredients and effective free radical eradicators. Therefore, these findings will offer a comprehensive overview of medicinal ingredients applicable in senotherapeutic food and product development.

Keywords: Antioxidant; inflammaging; Malaysian Medicinal Herbs Metabolomics; NMR metabolomics; radical scavenging

YI026

Chemical Profiling and Persistence of Gasoline Residues on Natural and Synthetic Porous Substrates under Tropical Weathering Conditions

NOOR HAZFALINDA HAMZAH^{1*}, KHAIRUL OSMAN¹, SUVITRAA PARAMASIVAM¹, ROGAYAH ABU HASSAN¹, GINA FRANCESCA GABRIEL², WAN NUR SYUHAILA MAT DESA³

¹*Forensic Science Program, School of Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia*

²*Netherlands Measurement Institute, Thijsseweg 11, 2629 JA Delft*

³*Forensic Science Programme, School of Health Sciences, Universiti Sains Malaysia (Health Campus), 16150 Kubang Kerian, Kelantan, Malaysia*

**Correspondence: raviera@yahoo.com*

ABSTRACT

Understanding the persistence of gasoline residues on porous substrates is fundamental to arson investigations, providing critical evidence of malicious intent through the identification of ignitable liquid residues. In forensic practice, delays between fire suppression and evidence collection frequently result in the recovery of weathered debris, where volatile organic compounds have been significantly depleted by environmental factors. In tropical regions such as Malaysia, high ambient temperatures and relative humidity are hypothesised to accelerate this evaporation process, potentially hindering residue recovery. This study evaluates the retention and chemical profiling of gasoline residues on natural (100% cotton) and synthetic (100% polyester) substrates subjected to outdoor tropical weathering. Following controlled ignition and suppression, samples were exposed to environmental conditions for durations ranging from 30 minutes to 960 minutes (16 hours). Residues were extracted via passive headspace adsorption and analysed using gas chromatography–mass spectrometry (GC–MS) in strict accordance with ASTM E1618-19 standards. Nine diagnostic target compounds, including critical alkylbenzenes and indane, were consistently identified using the National Institute of Standards and Technology library with a minimum match quality of 85%. The findings demonstrated that both natural and synthetic substrates successfully retained identifiable gasoline residues throughout the 960-minute exposure period, despite the aggressive tropical environment. This confirms a viable window for short-term ILR recovery in warm, humid climates and underscores the significance of substrate-environment interactions in environmental forensic investigations.

Keywords: ASTM E1618-19; exposure duration; fire investigation; substrate-liquid interaction; tropical weathering

RP002

Development of the Malay Version of the Multidimensional Fall Efficacy Scale: Translation, Cultural Adaptation and Preliminary Validation

LEE MIAO ZHUANG¹, DEVINDER KAUR AJIT SINGH¹, SHAWN ENG HSEIN SOH², NORMALA MESBAH^{1*}

¹*Physiotherapy Programme, Centre for Healthy Ageing & Wellness, Faculty of Health Sciences, University Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Health and Social Sciences Cluster, Singapore Institute of Technology, 828608 Singapore*

**Correspondence: normala.mesbah@ukm.edu.my*

ABSTRACT

Falls remain a major public health concern among older adults, with fear of falling often under-recognised despite its substantial impact on function and participation. Most existing instruments primarily assess negative emotional responses to falling, while commonly used fall-related self-efficacy scales are largely unidimensional and do not capture confidence across the full fall process. The Multidimensional Fall Efficacy Scale (MdFES) was developed to address this gap by evaluating confidence across four stages of the fall continuum. In this study, we aimed to translate, culturally adapt, and establish preliminary validity (content and face validity) of the Malay version of the MdFES for community-dwelling older adults in Malaysia. Phase 1 involved forward-backward translation and cultural adaptation with AI-assisted translation support and expert panel review. Phase 2 evaluated preliminary validity. Content validity was assessed by six experts ($n = 6$) using the Item-Level Content Validity Index (I-CVI) and Scale-Level Content Validity Index (S-CVI/Ave). Face validity will be examined among 30 community-dwelling older adults ($n = 30$) using the Item-Level Face Validity Index (I-FVI) and Scale-Level Face Validity Index (S-FVI/Ave). Preliminary results showed that the Malay-MdFES demonstrated acceptable content validity, with I-CVI values ranging from 0.83 to 1.00 and S-CVI/Ave of 0.98, indicating excellent content validity and expert agreement. The Malay-MdFES is expected to be relevant, clear and culturally appropriate for Malaysian older adults. Its availability may help healthcare providers accurately identify older adults with reduced confidence at specific stages, enabling effective and targeted interventions that promote healthy ageing.

Keywords: Content validity; cultural adaptation; Malay version; multidimensional fall efficacy scale; older adults

RP005

From Affection to Connection: The Pivotal Role of Maternal Openness and Interdependence in Predicting Emerging Adults' Psychological Well-being

NOVI QONITATIN*, ERIN RATNA KUSTANTI

Faculty of Psychology, Universitas Diponegoro, Kota Semarang, Jawa Tengah 50275, Indonesia

**Correspondence: qonitatin@gmail.com*

ABSTRACT

In collectivist cultures such as Indonesia, the transition to adulthood necessitates a complex renegotiation of the mother-child relationship, which remains pivotal to mental health. This study seeks to investigate the impact of maternal togetherness, warmth, openness, and interdependence on the psychological well-being (PWB) of emerging adults. Data were gathered from 312 university students ($N = 312$, ages 18–29) in Semarang, Indonesia, employing a predictive correlational methodology. Participants completed four validated mother-specific relational scales: Togetherness (13 items, $\alpha = 0.936$), Warmth (32 items, $\alpha = 0.969$), Openness (22 items, $\alpha = 0.949$), and Interdependence (18 items, $\alpha = 0.919$), along with the modified Psychological Well-being Scale (25 items, $\alpha = 0.936$). The findings from multiple linear regression analysis indicated that psychological well-being was significantly predicted by maternal relational dynamics ($F = 6.503$, $p < 0.001$), explaining 7.8% of the variance ($R^2 = 0.078$). Notably, Warmth and Togetherness, traditionally emotive variables, were not significant predictors in this model, as revealed by a partial analysis. Conversely, Maternal Openness ($\beta = 0.151$, $p < 0.05$) and Interdependence ($\beta = 0.150$, $p < 0.05$) emerged as the only significant positive predictors. These findings align with the autonomous-relatedness paradigm, suggesting a societal shift where egalitarian communication and functional support are more crucial for psychological well-being than mere physical presence or attachment. Consequently, family interventions for this demographic should prioritise fostering open communication and supportive interdependence, rather than solely emphasising closeness.

Keywords: Emerging adulthood; Indonesian culture; maternal interdependence; mother-child communication; openness; psychological well-being

RP008

Determinants of Return to Work After Stroke in Asia: A Scoping Review

ALISAFAR ALI ZOHOR ALI^{1*}, NURSUHAILI MOHD AMIN²

¹*School of Physiotherapy, Faculty of Allied Health Professions, AIMST University, 08100 Bedong, Kedah, Malaysia*

²*School of Occupational Safety and Health, Faculty of Allied Health Professions, AIMST University, 08100 Bedong, Kedah, Malaysia*

**Correspondence: alifsafarali@aimst.edu.my*

ABSTRACT

Return to work (RTW) after stroke represents a critical milestone in functional recovery, economic productivity and long-term social participation among working-age survivors. Although determinants of RTW have been widely studied, existing evidence is predominantly derived from Western populations, limiting contextual relevance to Asian countries undergoing rapid demographic ageing and labour market transitions. This scoping review aimed to synthesise evidence on facilitators and barriers to RTW among stroke survivors in Asia and to classify identified factors according to the International Classification of Functioning, Disability and Health framework. A systematic search of electronic databases identified quantitative studies examining factors associated with RTW after stroke. Extracted data included study characteristics, operational definitions of RTW and statistically significant associated variables. Six studies met the inclusion criteria. Consistent facilitators included younger age, lower stroke severity, better functional mobility and preserved cognitive performance. Frequently reported barriers comprised severe physical impairment and cognitive deficits, with psychological factors also emerging in several studies. Environmental and occupational determinants, including job demands and workplace accommodation, were comparatively underexplored. These findings underscore the multidimensional and context-sensitive nature of workforce reintegration following stroke in Asian settings. Greater emphasis on environmental and policy-level determinants is required to support sustainable vocational rehabilitation strategies and inform culturally responsive workforce reintegration frameworks in Malaysia and the wider Asia-Pacific region.

Keywords: Asia; disability and health; international classification of functioning; return to work; stroke; working age

RP010

DNMT3A/DNMT3B-Mediated Epigenetic Regulation of TRPV4 Drives Extracellular Matrix Stiffening in Lung Squamous Cell Carcinoma

KHAIRUNNISA MOHD KAMAL¹, AHMAD ROHI GHAZALI¹, GAYATHRI THEVI SELVARAJAH², NURUL SYAKIMA AB MUTALIB³, NADIAH ABU⁴, ENG WEE CHUA⁵, SITI FATHIAH MASRE^{1*}

¹*Biomedical Science Programme, Centre for Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Department of Veterinary Clinical Studies, Faculty of Veterinary Medicine, Universiti Putra Malaysia, 43400 Serdang, Malaysia*

³*Department of Biological Sciences and Biotechnology, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia*

⁴*UKM Medical Molecular Biology Institute, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia*

⁵*Centre for Drug and Herbal Development, Faculty of Pharmacy, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

*Correspondence: sitifathiah@ukm.edu.my

ABSTRACT

Extracellular matrix (ECM) stiffening is a key biomechanical hallmark of lung squamous cell carcinoma (SCC) that actively contributes to the tumour microenvironment (TME). However, the epigenetic mechanisms orchestrating mechanotransduction during multistage tumour progression remain incompletely understood. This study provides the first multistage *in vivo* mechanistic mapping of a DNMT3A/DNMT3B-TRPV4 epigenetic-mechanobiological axis in an N-nitroso-trichloroethylurea (NCTU)-induced lung SCC model, uncovering evidence of early pre-malignant epigenetic priming. Female Balb/C mice were exposed to NTCU for 15 weeks (pre-malignant stage) or 30 weeks (malignant stage), and mice treated with 70% acetone comprised the control group. The expression levels of two DNA-methylating proteins, DNMT3A and DNMT3B, and TRPV4 were analysed by immunohistochemistry, ECM remodelling by Sirius Red staining, and TRPV4 promoter methylation by methylation-specific PCR. We found that DNMT3A was significantly upregulated at both stages ($p < 0.05$), with higher expression at the pre-malignant stage, indicating early *de novo* methylation activity preceding overt malignancy. DNMT3B exhibited moderate induction. Correspondingly, MSP detected TRPV4 amplification predominantly in the unmethylated reaction; however, reduced unmethylated band intensity in NTCU-treated tissues indicates a relative shift in promoter methylation. These stage-dependent methylation changes were associated with altered TRPV4 expression, supporting epigenetic modulation rather than complete silencing. The significantly elevated TRPV4 expression at the pre-malignant stage ($p < 0.05$), followed by a marked reduction in malignant lesions ($p < 0.05$). These molecular alterations coincided with significant

collagen accumulation in NTCU-treated lungs at both stages ($p < 0.05$), with malignant lesions demonstrating greater deposition than pre-malignant tissues, reflecting progressive ECM stiffening and suggesting dynamic recalibration of mechanosensory signalling across tumour evolution. Collectively, these findings identify early DNMT3A/DNMT3B-mediated epigenetic priming as a potential upstream driver of mechanotransduction rewiring and progressive ECM stiffening, providing a novel mechanistic framework for lung SCC evolution and early intervention strategies.

Keywords: Extracellular matrix; epigenetic regulation; lung squamous cell carcinoma; mechanotransduction; transient receptor potential vanilloid

RP011

Efficacy of Beta-Glucan 1,3/1,6 on Health-Related Outcomes among Adults: A Systematic Review

MUSHAVIA ASHRAF¹, NUR NADIA MOHAMAD HABIBULLAH¹,
MUNIRAH ISMAIL¹, SUMAIYAH MAT¹, NORHAYATI IBRAHIM², SHIRLEY
GEE HOON TANG³, MOHD FAISAL ABDUL HAMID⁴, KALAVATHY
RAMASAMY⁵, SUZANA SHAHAR^{1*}

¹Dietetic Programme, Centre for Health Aging and Wellness, Universiti Kebangsaan Malaysia, Kuala Lumpur
50300, Malaysia

²Clinical Psychology Programme Centre for Healthy Ageing and Wellness, Universiti Kebangsaan Malaysia,
Kuala Lumpur 50300, Malaysia

³Biomedical Science Programme, Centre for Toxicology and Health Risk Studies, Universiti Kebangsaan
Malaysia, Kuala Lumpur 50300, Malaysia

⁴Respiratory Unit, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur,
Malaysia

⁵Faculty of Pharmacy, Universiti Teknologi MARA, Cawangan Selangor, Kampus Puncak Alam, 42300 Bandar
Puncak Alam, Selangor Malaysia

*Correspondence: suzana.shahar@ukm.edu.my

ABSTRACT

Beta-glucan, a polysaccharide derived from the cell walls of yeast species and fruiting bodies of mushrooms, has gained significant attention in various clinical and experimental settings in recent years due to its potential immunomodulatory, anti-inflammatory, and antioxidant therapeutic effects on human health. This systematic review and meta-analysis aim to determine the therapeutic effects of consuming beta-(1,3/1,6)-D-glucan on health outcomes. A comprehensive literature search was performed in three electronic databases to retrieve studies from 2013 to 2025 that applied open-source randomised controlled trials to investigate the impact of exclusive oral administration of fungal beta-glucan in any form and at any dosage to adult subjects or patients. A total of 31 RCTs of the 1431 records retrieved and met the eligibility criteria, thus included in the present review. The dosage of the supplementation ranged from 1mg to 8000 mg daily for up to 252 days. The physiological outcome of the majority of the interventions was immunomodulation at both low dose (<250 mg/d) and medium dose (250-750 mg/d), but medium dose had a more consistent effect, which resulted in (i) enhanced the immune system with SMD = -0.26 (95% CI: -0.41 to -0.12), $I^2 = 24\%$, leading to a decrease in the occurrence and severity of colds, flu, and other respiratory infections and alleviated allergic symptoms; (ii) contributed to an overall improvement in Quality-of-Life at medium dose (250-750 mg/d) with SMD = -0.74 (95% CI: -1.14 to -0.34); (iii) Improvement in the clinical outcomes such as URTI episodes, lipid profile, and weight reduction at high dose (>750 mg/d) with SMD = -0.32 (95% CI: -0.63 to -0.01); however, heterogeneity ($I^2 = 93-84\%$) was high. Participants who received beta-glucans reported elevated mood states and a general sense of well-being. At the same time, it might also be useful as a complementary agent to patients undergoing cancer therapies

and decrease comorbid conditions associated with obesity and osteoarthritis. Adverse effects (mild GI discomfort) were minimal, supporting the safety profile of beta-glucans. Supplementation with fungal beta-glucan is safe, well-tolerated, and shows promising immunomodulatory effects, such as enhanced immune defence, improved QoL and clinical outcomes. However, significant heterogeneity and variability in sources of beta-glucan and study populations highlight the need for more high-quality research.

Keywords: Clinical trial; fungal beta-glucans; health outcomes; systematic review

RP012

Pilot Study: Adaptation of Resilience-Building Intervention among Youth with Adverse Childhood Experience in Malaysia

**JUN KIAT NG¹, MAHADIR AHMAD^{1*}, CARYN MEI HSIEN CHAN¹,
SATIRAH ZAINALABIDIN², PONNUSAMY SUBRAMANIAM¹,
MICHAEL UNGAR³**

¹Department of Clinical and Health Psychology, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Kuala Lumpur 50300, Malaysia

²Biomedical Science Program, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

³Resilience Research Centre, Dalhousie University, Halifax, NS B3H 4R2, Canada

**Correspondence: mahadir@ukm.edu.my*

ABSTRACT

As child abuse cases are reported throughout the world, their impact on a person's life is pervasive. Adverse childhood experiences are associated with obesity, cancer, difficulties with self-regulation, and emotion regulation. Hence, this study adapted and modified a resilience-fostering intervention among individuals with adverse childhood experiences in Malaysia. This intervention study was designed based on Ungar's model and modified accordingly to the context of Malaysia by using a biopsychosocial-ecological model. For this study, it is a pilot study evaluating the feasibility of the adapted intervention study as part of the Assessment-Decision-Administration-Production-Topical experts-Training-Testing (ADPTT-ITT) framework. Ten participants were recruited and randomised into the control group and the intervention group. The intervention was conducted in both Malay and English languages. Non parametric tests were conducted to calculate resilience and surrogate resilience measures. For between-group comparison analysis between the control and intervention groups, adult resilience, patient health and well-being were found to be significantly different. The results provide insight for the clinical study.

Keywords: Adverse childhood experience; pilot study; resilience; resilience-building intervention; youth

RP016

miR-155 Modulation Induces Apoptosis and Regulates PTEN/MARCH7 Signalling in Breast Cancer-Derived Cell Lines

XIAOHUA CHEN^{1,2}, MING THONG ONG^{1*}¹*Institute for Research in Molecular Medicine, Universiti Sains Malaysia, 11800 USM, Pulau Pinang, Malaysia*²*Medical College of Shaoguan University, 512005 Shaoguan, China**Correspondence: omt@usm.my

ABSTRACT

Breast cancer (BC) progression is governed by intricate microRNA regulatory networks that modulate cellular processes, including survival, cell cycle progression, and metastasis. This study focuses on the functional role of microRNA-155 (miR-155) in BC-derived cell lines (MCF-10A, MCF-7, and MDA-MB-231). Cells were transfected with miR-155 mimics and inhibitors to evaluate the effects of over-expression and inhibition, respectively. Cell viability was assessed using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assays at 24, 48, and 72-hour intervals. Apoptosis and cell-cycle distribution were analysed by flow cytometry. In addition, the expression of selected genes and proteins, including PTEN, MARCH7, Akt and Caspase-3, was evaluated using RT-qPCR and Western blotting to provide preliminary insights into potential regulatory pathways. The results revealed that miR-155 plays a context-dependent role in BC cells. Gain-of-function assays demonstrated that miR-155 overexpression via mimics significantly induces apoptosis and suppresses BC cell survival in a time-dependent manner, with the most pronounced inhibitory effects observed at 48 and 72 hours. Cell cycle analysis revealed no significant G1 or G2/M arrest, indicating that miR-155 does not primarily inhibit cell growth through classical cell-cycle checkpoint regulation. Conversely, treatment with miR-155 inhibitors exhibited modest anti-apoptotic effects, partially preserving cell viability. Mechanistically, inhibition of miR-155 resulted in upregulation of MARCH7 and downregulation of PTEN, supporting their roles as downstream targets of miR-155. Furthermore, alteration of this signaling axis was associated with changes in cellular migratory and invasive behavior, suggesting that miR-155 may contribute to the metastatic processes in BC cells. However, a primary limitation of this study is the reliance on in vitro 2D cell culture models, which may not fully replicate the complexity of the tumor microenvironment. These findings highlight a context-dependent and non-canonical role of miR-155 in BC cells, contrasting with its widely reported oncogenic function. This supports its potential as both a biomarker and a therapeutic target for strategies aimed at modulating tumour growth and metastatic progression.

Keywords: Apoptosis; breast cancer; MARCH7; miR-155; PTEN

RP017

Factors Influencing Water Scarcity Preparedness in Urban Slums of Padang City: A Community Health Perspective

AZYYATI RIDHA ALFIAN^{1,2*}, AULIA ERID ANGELICA¹, AIDA SORAYA²,
LEE KHAI ERN², AZIEMAH ZULKIFLI²

¹*Faculty of Public Health Andalas University, Padang, Indonesia*

²*Institute for Environment and Development, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia*

**Correspondence: azyyatiridhaalfian@ph.unand.ac.id*

ABSTRACT

Indonesia faces a growing water crisis driven by rapid population growth, uneven resource distribution, and the escalating impacts of climate change. Despite government initiatives, access to safe drinking water remains limited, with only 11.9% of households having water that meets quality standards. In urban slums, particularly in Padang City, reliance on unsafe sources such as shallow wells increase the risk of contamination and waterborne diseases, posing significant public health challenges. Infrastructure gaps, limited community awareness, and inadequate supervision further exacerbate household vulnerability to water scarcity. Aligned with the theme Advancing Community Health through Research, Innovation and Partnership, this study investigates the determinants of water scarcity preparedness among households in urban slum areas of Padang City. The research employs a cross-sectional design involving 102 households selected through proportional random sampling from three risk categories across nine urban villages. Data were collected between December 2022 and July 2023 using structured questionnaires and secondary data from local government reports. The findings reveal that education level ($p = 0.011$), availability of facilities and infrastructure ($p = 0.023$), and supervision ($p = 0.034$) significantly influence community preparedness. These results underscore the critical role of community-based health education, infrastructure innovation, and strengthened monitoring systems in building resilience against water scarcity. This study highlights how evidence-based research can inform innovative public health interventions, such as community-led water safety planning, low-cost water treatment technologies, and integrated surveillance systems. Furthermore, it emphasises the importance of multi-sectoral partnerships among local government, public health institutions, community organisations, and academic stakeholders to enhance sustainable water management. By integrating research, innovation, and collaborative action, this work contributes to advancing community health resilience and improving equitable access to safe water in vulnerable urban populations.

Keywords: Community health; Padang city; preparedness; urban slums; water scarcity

RP018

Predictors of Self-Reported Practice in Ventilator-Associated Pneumonia Prevention among Critical Care Nurses in Sarawak Public Hospitals

TAN JIA HUI, CHE CHONG CHIN*, TANG LI YOONG, CHONG MEI CHAN

Department of Nursing Science, Faculty of Medicine, Universiti Malaya, Kuala Lumpur, Malaysia

**Correspondence: chechongchin@um.edu.my*

ABSTRACT

Ventilator-associated pneumonia (VAP) remains one of the most difficult infections to control in intensive care settings and continues to pose a serious threat to patient safety. As healthcare systems move toward greater innovation and international collaboration, it is increasingly important to understand how nurses apply evidence-based VAP prevention strategies in daily practice. This study assesses the level of knowledge, self-reported practices, barriers, and predictors of VAP prevention among critical care nurses in Sarawak, Malaysia. A cross-sectional survey involving 298 nurses from four public hospitals was conducted using a structured questionnaire that captured demographics, VAP-related knowledge, self-reported practices, and identify barriers related to VAP prevention. Hierarchical multiple regression was used to determine the key predictors of practice. Although most nurses reported a high level of compliance with VAP prevention measures (76.5%), two-thirds demonstrated poor knowledge. Knowledge and practice were only weakly correlated ($p < 0.001$). Practice was significantly influenced by the type of unit, bed capacity, nurse-to-patient ratio, and knowledge level, while the most frequently reported barriers included staffing shortages, forgetting certain steps, and the lack of written protocols. These findings highlight a clear gap between what nurses know and what is practiced at the bedside. Strengthening VAP prevention efforts will require not only education, but also better staffing support, clearer protocols, and stronger institutional commitment. Such improvements are essential for advancing patient safety and are consistent with the broader i-SIHAT 2026 emphasis on health innovation and global partnerships.

Keywords: Critical care nursing; evidence-based practice; infection prevention; Malaysia ventilator-associated pneumonia

RP019

Development, Content Validation and Usability Testing of the Mobile Application for Vestibular Rehabilitation Therapy in Adults with Vestibular Problem

NUR DINI IZZATI KHAIROL ANUR¹, NOR HANIZA ABDUL WAHAT^{1*},
FADILLA' ATYKA NOR RASHID²

¹Centre for Rehabilitation and Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300, Kuala Lumpur, Malaysia

²Centre of Artificial Intelligence Technology, Faculty of Information Science & Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

*Correspondence: nor_haniza@ukm.edu.my

ABSTRACT

Vestibular disorders commonly cause dizziness, imbalance, and increased risk of falls, significantly affecting daily functioning and quality of life. Although vestibular rehabilitation therapy (VRT) is an evidence-based intervention, access to structured rehabilitation programs remains limited due to reliance on supervised clinical sessions and resource constraints. Digital health technologies offer an opportunity to improve accessibility by enabling home-based rehabilitation supported by remote clinical monitoring. This study aimed to develop and evaluate the Malaysian Adult Vestibular Rehabilitation Therapy (MAVeRT) application as a digital platform and demonstrate the feasibility of translating conventional vestibular rehabilitation into a technology-enabled system. The MAVErT application was developed from scratch through a structured multi-phase research process. Initially, a VRT exercise module was adapted and produced as guided instructional videos tailored for home-based practice. The validity of the MAVErT scripts and videos was evaluated by an expert panel comprising audiologists and physiotherapists, with a mean age of 39.17 years (SD = 4.13) and an average professional experience of 13.33 years (SD = 7.94). Content validation demonstrated strong agreement among experts, with Item Content Validity Index (I-CVI) values ranging from 0.83 to 1.00 for both script and video components, while the Scale Content Validity Index average (S-CVI/Ave) was 0.96 for the scripts and 1.00 for the videos, exceeding the recommended validity thresholds. Subsequently, a MAVErT prototype application integrating the validated exercise videos and a digital diary was developed and evaluated for usability among pilot participants. The usability assessment yielded a mean "Skala Kebolehgunaan Aplikasi Mudah Alih" (SKAMA) score of 71.25 (SD = 19.63), exceeding the recommended benchmark of 68, indicating good usability for the MAVErT prototype. Following the usability evaluation, the prototype was further enhanced into a fully functional digital platform integrated with a clinician dashboard to enable remote monitoring of patient engagement and rehabilitation progress. Preliminary piloting among patients with vestibular-related symptoms and healthcare professionals suggests that the MAVErT platform is feasible and practical for delivering digital vestibular rehabilitation. Overall, the MAVErT platform demonstrates the potential to support the digitalisation of vestibular rehabilitation and to facilitate accessible, patient-centered, clinician-monitored rehabilitation services.

Keywords: Digital health; mobile health application; tele-rehabilitation; vestibular rehabilitation therapy

RP020

The Impact of Orthokeratology on Ocular Dimensions and Peripheral Eye Length in Myopic Children: A 3D Magnetic Resonance Imaging 12 months Study

YU CHEN LOW^{1*}, BARIAH MOHD-ALI¹, MIZHANIM MOHAMAD SHAHIMIN¹, NORHANI MOHIDIN¹, HAMZAINI ABDUL-HAMID², SITI SALASIAH MOKRI³

¹*Optometry and Vision Science Program and Research Centre for Community Health, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Department of Radiology, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia*

³*Department of Electrical, Electronics and Systems, Faculty of Engineering and Built, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia*

**Correspondence: yuchenlow@ukm.edu.my*

ABSTRACT

This study investigated 12-month changes in peripheral eye length (PEL), axial length (AL), and three-dimensional ocular dimensions (longitudinal axial length (LAL), horizontal width (HW), and vertical height (VH)) in myopic children undergoing Orthokeratology (Ortho-K) treatment compared to single-vision spectacle (SVS) wearers using 3-Tesla Magnetic Resonance Imaging (MRI). A total of 70 myopic Chinese schoolchildren aged 8–9 years old (Ortho-K, $n = 45$ and SVS, $n = 25$) were recruited. While the SVS group exhibited significant axial elongation and a shift towards peripheral hyperopic defocus, the Ortho-K group demonstrated significantly less elongation in axial length (AL) and PEL across most eccentricities ($p < 0.05$), and a shift towards myopic defocus from 10 -30 in nasal and temporal. LAL was significantly shorter in the Ortho-K group compared to SVS ($p < 0.05$). While individual HW, VW did not reach significance within eyeball shape of each group, repeated measures ANOVA confirmed significant between-group differences in global eyeball shape over 12 months ($p < 0.05$). Notably, Ortho-K induced a comprehensive reduction in global ocular geometry across LAL, HW and VH dimensions. Correlation analysis revealed that SE significantly correlated with LAL in the SVS group ($p < 0.05$); however, the Ortho-K group demonstrated a “decoupling” effect, where SE showed no significant correlation with ocular dimensions ($p > 0.05$), while FK correlated significantly with all dimensional changes ($p < 0.05$). These findings highlight that while myopic progression in the SVS group is primarily driven by axial elongation, refractive changes in the Ortho-K group are attributed to corneal reshaping. Over 12 months, Ortho-K wear modifies ocular dimensions and inhibits AL elongation more significantly than SVS. Hence, providing a structural basis for its efficacy in controlling myopic progression.

Keywords: Axial length; MRI; myopia; ocular dimension; orthokeratology

RP021

The Use of Augmentative and Alternative Communication by Children with Autism in Special Education Classrooms in Malaysia: Perspective of Special Education Teachers and Parents

HAZEL BEE LENG CHEAH^{1*}, SUSHEEL JOGINDER SINGH¹, IVONNE JINN YEE LING¹, YAZMIN AHMAD RUSLI¹, NATALIE R. ANDZIK²

¹*Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Department of Special and Early Education, Northern Illinois University, DeKalb, IL, USA*

**Correspondence: hazelbleng@gmail.com*

ABSTRACT

Many children with autism (CwA) remain non-verbal for extended periods and were introduced to Augmentative and Alternative Communication (AAC) to facilitate their communication. In Malaysia, many CwA who attend the Integrated Special Education Program (PPKI) struggle with the use of AAC in the classroom. This paper explores the perspectives of parents and special education teachers on AAC implementation in the classroom settings. A total of four focus group discussions were conducted with 12 parents and 15 special education teachers. Data were transcribed and analysed using the qualitative content analysis. Several challenges were identified. These included limited knowledge and skills, misconceptions about AAC, student-related challenges such as low motivation, and AAC system issues like limited vocabulary or device safety and maintenance. Teachers also highlighted workplace challenges such as a large number of students in a class, limited funding, and restrictive school policies. Despite these challenges, both parents and special education teachers acknowledged the benefits of AAC, including increased vocabulary, improved two-way communication, and reduced challenging behaviors. Their shared perspectives emphasised the importance of collaboration between teachers, families, and professionals for the effective implementation of AAC in school settings.

Keywords: Augmentative and alternative communication; Malaysia; special education teachers, autism spectrum disorder; speech-language pathologist

RP024

From Food Waste to Food Acceptance: The Teddy Breakfast Project for Hospitalised Children

NURUL HUDA RAZALLI^{1*}, MUHAMMAD ANWAR LOTPI¹, HUDA RAFIKA HIDZIR¹, NURUL FATIHAH AZHAR¹, SYAFIY A. AZMAN¹, NURUL ATILIA ABDULLAH²

¹*Dietetic Programme, Centre for Healthy Aging and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Dietetic and Nutrition Unit, Universiti Kebangsaan Malaysia Children's Specialist Hospital, 56000 Cheras, Kuala Lumpur, Malaysia*

**Correspondence: nurulhuda.razalli@ukm.edu.my*

ABSTRACT

Plate waste is frequently observed among hospitalised children and may limit dietary intake while contributing to avoidable food disposal. Reducing food waste is consistent with Sustainable Development Goal (SDG) 12, which promotes responsible consumption. To address low acceptance of the existing bread-based breakfast (French toast served with baked beans), Teddy Breakfast was introduced at the Specialist Children's Hospital, Universiti Kebangsaan Malaysia, as an alternative child-friendly option aimed at improving meal acceptance and reducing plate waste. The meal consists of white bread with peanut butter and chocolate spread, decorated with raisins and banana slices to resemble a teddy bear. A two-phase study was conducted to examine sensory acceptance and changes in plate waste. Phase 1 (February–July 2024) involved baseline plate waste measurement using the Comstock visual estimation method, followed by product development, nutrient analysis, and sensory evaluation. Phase 2 (February–July 2025) involved serving the Teddy Breakfast in place of the existing menu and reassessing plate waste using the same method. The intervention achieved high overall acceptance (acceptance index 82.6%; mean score 4.1). Average plate waste decreased significantly from 77.34% at baseline to 55.5% after the introduction of the new menu ($p < 0.01$), with similar reductions across both age groups. The findings suggest that simple, child-oriented meal presentation strategies can improve food acceptance and reduce waste in paediatric hospital settings, with potential application in other child-focused food service environments such as preschools and primary schools.

Keywords: Paediatric nutrition; plate waste; sensory acceptance

RP027

Mechanism of FODMAP Restriction in Functional Gastrointestinal Disorder Patients: Preliminary Data from a Randomised Controlled Trial

JAYSRIINA MAHALINGA MOORTHY¹, YOU YEE XING², DEBORAH CHEW CHIA HSIN³, VANITHA MARIAPPAN⁴, SYARATUL DALINA YUSOFF⁵, HUI-MIN NEOH⁶, LEE YEONG YEH⁷, SHANTHI KRISHNASAMY^{1*}

¹*Centre of Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Centre of Healthy Ageing & Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

³*Gastroenterology and Hepatology Unit, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia*

⁴*Center for Toxicology & Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

⁵*Centre for Drug and Herbal Development, Faculty of Pharmacy, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

⁶*UKM Medical Molecular Biology Institute, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia*

⁷*School of Medical Sciences, GI Function and Motility Unit, Universiti Sains Malaysia, 15200 Kota Bharu, Kelantan, Malaysia*

**Correspondence: shanthikrishnasamy@ukm.edu.my*

ABSTRACT

Functional Gastrointestinal Disorders (FGIDs), including irritable bowel syndrome (IBS) and functional dyspepsia (FD) are frequently managed with restriction of fermentable carbohydrates (FODMAPs). FGID patients often present with diminished quality of life (QoL), increased food avoidance behaviours and have been linked with orthorexia nervosa, a pathological preoccupation with “healthy eating”. A parallel-group randomised controlled trial was conducted to determine the effectiveness of diets with varying FODMAPs and their impact on FGID patients. Eligible subjects are screened at baseline and randomly assigned to one of three diet intervention arms: Low FODMAP diet, Gentle FODMAP diet, and Traditional Dietary Advice, which they were required to follow for two weeks. Baseline and post-intervention assessments included the IBS-Severity Scoring System (IBS-SSS), Gastrointestinal Symptom Rating Scale (GSRS), European QoL 5 Dimensions 5 Level (EQ-5D-5L), 4-day food diaries (nutrient analysis), Hospital Anxiety and Depression Scale (HADS), Orthorexia-Nervosa questionnaire (ORTHO-15), and the Avoidant/Restrictive Food-intake disorder scales (NIAS). The preliminary data was analysed to determine the effect of FODMAP restriction on these outcomes. Continuous variables were analysed using mixed ANOVA, while categorical variables were compared using chi-square tests. Based on the pre-post data obtained from 24

participants, symptom severity improved from baseline to post-intervention regardless of type of diet, indicating all three dietary approaches were beneficial ($p < 0.05$). The mobility ($p = 0.016$) and usual activities ($p = 0.031$) parameters from EQ-5D-5L were presented with significant arm-to-arm interactions. Energy intake fell markedly ($p = 0.007$) together with fat intake ($p < 0.001$) and iron intake ($p = 0.039$). ORTHO-15 scores improved significantly ($p = 0.004$) with the low FODMAP diet showing the largest estimated marginal-mean rise, indicating the biggest improvement in reducing orthorexia-related eating behaviour score ($p = 0.016$) which may be attributed to symptom improvement associated with the intervention. Descriptive analyses demonstrated shifts towards normal level in anxiety and depression categories, with heightened picky eating and fear post-intervention as expected. Preliminary findings suggest that dietary interventions, particularly the low FODMAP diet may improve gastrointestinal symptoms, QoL and eating-related behaviours in FGID patients.

Keywords: FODMAP restriction; food avoidance; functional dyspepsia; functional gastrointestinal disorder; irritable bowel syndrome

RP028

Psychologically Informed Practice in Tinnitus Management: Audiologist Perceived Barriers, Facilitators and Preparedness in Malaysian Government Hospitals

MOHAMAD AZMEER SADALI^{1,2}, WAN SYAFIRA ISHAK^{1,3*}, MAHADIR AHMAD^{4,5}, EMMA C. LAIRD⁶, CAROL MACDONALD^{7,8}, DEREK J. HOARE^{7,9}

¹Centre for Healthy Ageing & Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Audiology Unit, Department of Otorhinolaryngology, Hospital Sungai Buloh, 47000 Sungai Buloh, Selangor, Malaysia

³Audiology Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

⁴Centre for Community Health Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

⁵Clinical Psychology Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

⁶Department of Community and Clinical Health, La Trobe University, Bundoora VIC 3086, Australia

⁷National Institute for Health and Care Research, Nottingham Biomedical Research Centre, Hearing Sciences, Mental Health and Clinical Neurosciences, University of Nottingham, Nottingham NG1 5DU, United Kingdom

⁸Department of Psychology, University of Stirling, Stirling FK9 4LA, Scotland, United Kingdom

⁹Department of Speech and Hearing Sciences, University College Cork, Cork, T12 AK54, Ireland

*Correspondence: wsyafira@ukm.edu.my

ABSTRACT

Psychologically informed practice sits between traditional biomedical care and specialist mental health services, where healthcare practitioners address the psychological aspects of chronic physical conditions. First developed in physiotherapy, this approach has increasingly been adopted by audiologists for managing tinnitus disorder, especially in the United Kingdom the United States. With limited evidence from Southeast Asia, this study aimed to identify gaps in implementing psychologically informed tinnitus management by examining audiologists' perceived barriers, facilitators, and preparedness. A cross-sectional survey was conducted among audiologists in Malaysian government hospitals across multiple regions (n = 104) who reported working with adults presenting with tinnitus as part of their clinical practice. The survey collected information on current practices, experiences in managing tinnitus, and perspectives on psychologically informed approaches. Most participants reported at least one barrier to discussing mental health when caring for patients with tinnitus (98.1%), with the most common being insufficient knowledge or training in psychological approaches (83.7%). Interpreted using the COM-B model of behaviour change, barriers were primarily related to capability and opportunity factors, including limited

confidence in identifying distress and insufficient consultation time or workplace support for these discussions. In contrast, motivational factors appeared to act as facilitators, with most audiologists recognising the importance of mental health in tinnitus management and agreeing that counselling falls within their scope of practice. Many participants also reported preparedness to incorporate psychologically informed interventions, particularly structured referral protocols for psychological services (76.9%). Overall, audiologists in Malaysian government hospitals demonstrated strong motivation to incorporate psychologically informed approaches into tinnitus management; however, capability-related limitations and organisational constraints may hinder their implementation. These findings highlight key targets for future efforts in training, service development, and policy support to facilitate the integration of psychologically informed tinnitus interventions within Malaysian audiology services.

Keywords: Audiologists; COM-B model; implementation research; psychologically informed practice; tinnitus management

RP033

Association of Short-Term and Working Memory Capacity with Brain Activation among Adolescents with Internet Gaming Addiction Symptom: A Preliminary Finding

ZATI AMALINA MOHD ABDUL GHAFAR¹, AHMAD NAZLIM YUSOFF^{1*},
CHING SIN SIAU², ELZA AZRI OTHMAN³, NI ADILA SULAIMAN¹,
NOOR FADZLIYANA MOHD FADIL^{1,4}, FAIZAH MOHD ZAKI⁵,
MAZLYFARINA MOHAMAD¹, NOOR ALAUDIN ABDUL WAHAB⁶

¹Centre for Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Centre for Community Health Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

³School of Medical Imaging, Faculty of Health Sciences, Universiti Sultan Zainal Abidin, 21300 Kuala Nerus Terengganu, Malaysia

⁴Medical Imaging Department, School of Health Sciences, KPI Healthcare University, Kota Seriemas, 71800 Nilai, Negeri Sembilan, Malaysia

⁵Department of Radiology, Faculty of Medicine, Universiti Kebangsaan Kuala Lumpur, 56000 Cheras, Kuala Lumpur, Malaysia

⁶Center for Rehabilitation and Special Needs, Faculty of Health Science, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

*Correspondence: nazlimtrw@ukm.edu.my

ABSTRACT

Internet Gaming Addiction (IGA) has become a growing public health concern. Studies showed that moderate gaming can enhance cognition but excessive gaming may detrimentally impact memories. This study investigated brain activation patterns during auditory short-term (STM) and working memory (AWM) tasks among adolescents with IGA symptoms. Nine participants with IGA symptoms and twelve controls participated in this study. All participants took a STM test using the Malay Version of Auditory Learning Test (MVAULT) while AWM capacity was assessed using the digit span test (DST) and backward word repeat test (BWRT) during an functional magnetic resonance imaging (fMRI) scan. MVAULT and DST data were analysed using SPSS while the fMRI data were analysed using SPM12 for neural activation. Spatial activation from eight regions of interest (ROI) i.e. superior temporal gyrus (STG), precuneus (PRE), posterior cingulate cortex (PCC), anterior cingulate cortex (ACC), inferior frontal gyrus (IFG), dorsolateral prefrontal cortex (dlPFC), primary auditory cortex (PAC) and orbitofrontal cortex (OFC) were recorded. The Man-Whitney U test showed that there is no significant difference ($p > 0.05$) in the cognitive performance and brain activation between addicted and control groups. The highest average brain activations were found at STG and PRE. In the control group, Spearman's correlations revealed that A6 MVAULT scores was negatively correlated with IFG activation ($\rho = -0.446$, $p = 0.043$) while DST scores were negatively correlated with PRE activation ($\rho = -0.459$, $p = 0.036$). In the IGA group, MVAULT and DST

scores were strongly negatively correlated with STG, PCC, dl-PFC and PAC activation. Simple linear regression also showed that MVAULT, BWRT and Digit Span scores were poor predictors of both cognitive performance and brain activation across all eight regions ($r^2 < 0.10$). Consequently, the relationship between IGA and STM/AWM remains inconclusive. Future studies should utilise larger sample sizes to assess the impact of IGA on neurocognitive development.

Keywords: Activation; addiction; adolescent; brain; gaming

RP037

Toward Integrated Microstructural and Hemodynamic Brain Imaging with Intravoxel Incoherent Motion Imaging at 3T MRI

KINGKARN APHIWATTHANASUMET*

Department of Radiological Technology, Faculty of Allied Health Sciences, Naresuan University, Phitsanulok, Thailand

**Correspondence: kingkarna@nu.ac.th*

ABSTRACT

Conventional diffusion-weighted imaging (DWI) provides valuable insights into brain tissue microstructure but is limited in separating true diffusion from perfusion-related signal components. Intravoxel incoherent motion (IVIM) imaging enables simultaneous, non-contrast assessment of both diffusion and microvascular perfusion, offering potential advantages in characterising brain tumors. This study aimed to evaluate the utility of IVIM imaging at 3T magnetic resonance imaging (MRI) for differentiating normal brain tissue and gliomas of varying World Health Organization (WHO) grades II–IV, with a focus on microstructural and hemodynamic characterisation. IVIM MRI was performed on a 3T clinical system using a multi-b-value DWI sequence. The dataset used in this study was obtained from the OpenNeuro database, including 3D T1 TFE, 3D FLAIR and 2D multi-echo multi-shell HARDI diffusion-weighted imaging with TE = 85 ms, 1.5 mm isotropic voxel size, FOV = 160 × 160 × 96 mm³, and 40 b-values (including 0, 400, 800, 1600 and 3200 s/mm²). Quantitative parameters, including the true diffusion coefficient (D), pseudo-diffusion coefficient (D*), and perfusion fraction (f), were estimated using a bi-exponential model. Regions of interest were defined in healthy brain tissue and in histopathologically confirmed gliomas. Comparative analysis of IVIM parameters across groups was conducted to assess their diagnostic performance. IVIM-derived parameters demonstrated distinct patterns between healthy tissue and gliomas of different grades (II–IV). The diffusion coefficient (D) showed a decreasing trend with increasing tumor grade, reflecting higher cellularity in high-grade gliomas. In contrast, perfusion-related parameters (D* and f) tended to increase in higher-grade tumors, consistent with enhanced microvascular proliferation and angiogenesis. These findings suggest that IVIM imaging provides complementary information on both tissue microstructure and tumor vascularity. In conclusion, IVIM imaging at 3T MRI offers a promising, non-invasive approach for simultaneous assessment of brain microstructure and hemodynamics. It enables differentiation between healthy tissue and gliomas of different grades (II–IV), potentially supporting tumor grading and improving diagnostic accuracy without the need for contrast agents.

Keywords: Brain microstructure; diffusion MRI; glioma; IVIM imaging

RP040

Low Calorie Sweeteners Consumption: Associations with Caloric Compensation and Appetite-Regulating Hormones

WAN FATHIN FARIZA*

Department of Nutrition Sciences, Kulliyyah of Allied Health Sciences, International Islamic University

Malaysia, 25200 Kuantan, Pahang, Malaysia

**Correspondence: wfathin@iiu.edu.my*

ABSTRACT

The efficacy of low-calorie sweeteners (LCS) in achieving caloric reduction is frequently debated due to concerns over potential compensatory energy intake and disruption of appetite regulation. This cross-over, single-blind clinical study investigated the acute effects of sugar, stevia, and aspartame preloads on subsequent food intake and the concentrations of appetite-regulating hormones in 21 healthy adult subjects. Subjects completed three non-consecutive test days, consuming composite preloads sweetened with sugar, stevia, or aspartame, followed by an ad libitum test meal. Total energy intake (preload + lunch) was significantly lower in meals preceded by LCS compared to sugar meals (e.g., aspartame meals were 8.4% lower than sugar, $p = 0.035$). Subjects demonstrated incomplete energy compensation for the reduced calories in LCS preloads (stevia: 45.6%; aspartame: 54.1%), confirming no caloric overcompensation. In terms of appetite-regulating hormones, the sugar preload resulted in a significantly higher total area under the curve (AUC) for leptin ($p < 0.001$) and a significantly lower total AUC for ghrelin ($p = 0.029$ for stevia vs. sugar) compared to both LCS preloads. Insulin AUC was not statistically significant between groups, but leptin levels correlated positively with body fat across all sweeteners ($p < 0.001$). These findings indicate that replacing sugar with LCS (stevia or aspartame) successfully reduces total energy intake without inducing compensatory overeating or negatively impacting subjective appetite responses. This supports the use of LCS as a beneficial tool in a calorie-deficient food intake.

Keywords: Energy compensation; ghrelin; leptin; low-calorie sweeteners; stevia

RP041

Capturing the Patient's Experience in Virtual Care: Development and Validation of the MPEQ-VC for Malaysian Primary Health Clinics

NASIM KUTHOOSE, KHALID IBRAHIM, MOHAMAD RODI ISA*

*Public Health Medicine Department, Faculty of Medicine, University Teknologi MARA, 47000 Sungai Buloh,
Selangor, Malaysia*

**Correspondence: rodi@uitm.edu.my*

ABSTRACT

Virtual consultation services have expanded rapidly across Malaysian health clinics, with over 230 facilities now offering this modality of care to community-based primary care patients. However, no validated, culturally appropriate Malay-language instrument has been available to assess patient experience with these services, leaving the voices of Malaysian primary care patients systematically underrepresented in the evaluation of digital health innovations. This study aimed to develop and validate the Malay Version of the Patient Experience Questionnaire for Virtual Consultation Services (MPEQ-VC) for use in the government health clinics of Malaysia. A two-phase sequential design was adopted, guided by instrument development framework and the COSMIN reporting guidelines. Phase 1 comprised item generation in Malay, content validation by a nine-member multidisciplinary expert panel, and face validation by ten experienced patients. Phase 2 involved exploratory factor analysis on 211 respondents, internal consistency and test-retest reliability assessment, and confirmatory factor analysis on an independent sample of 113 respondents recruited from eight government health clinics across four districts in Selangor stratified by urbanicity. The final 25-item MPEQ-VC demonstrated a refined and validated three-domain structure comprising Ease of Use, Patient-Centered Care, and Involvement of Family Members. Confirmatory factor analysis confirmed acceptable model fit (CFI = 0.912; TLI = 0.900; RMSEA = 0.093). Composite Reliability was 0.890, Average Variance Extracted was 0.731, and Cronbach's alpha exceeded 0.93 across all domains. Discriminant validity was clearly established across all domain pairs. The MPEQ-VC is the first psychometrically validated Malay-language patient experience instrument for virtual consultation services in Malaysian primary care, offering a culturally grounded tool to amplify patient voices and inform community health quality improvement. Additionally, it serves as a guide for healthcare workers to gain a deeper understanding of the dimensions of patient experience within a screen-mediated platform, which is crucial for their service delivery.

Keywords: Development; patient experience; primary healthcare; questionnaire; validation

RP043

Development and Content Validation of the STAR Parenting Module to Improve Socio-Emotional Development for School Readiness of Children with Cochlear Implants

CHA LONG GEH, CILA UMAT*

Rehabilitation & Special Needs Studies, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

**Correspondence: cila@ukm.edu.my*

ABSTRACT

Parents of children with cochlear implants (CIs) face unique challenges in fostering their children's socio-emotional development, a critical component of school readiness. There is a recognised lack of culturally adapted, evidence-based interventions to support these parents effectively. This study aimed to develop and validate a psycho-educational module, the "Socio-emotional Teaching and Raising (STAR) Parenting" module, designed to enhance the socio-emotional school readiness skills of preschool children with CIs in Malaysia. The module was constructed using Module Development Model, comprising two phases: (i) module draft construction and (ii) module evaluation. The content was based on an earlier study on needs analysis and empirical literature, structured into six sessions covering parenting knowledge, facilitator roles, and practical workshop strategies. Content validity was assessed by a panel of five experts (audiologists, a speech-language pathologist, a clinical psychologist, and a special education teacher) using a 92-item Content Validity Questionnaire rated on a 5-point Likert scale. The Item-Content Validity Index (I-CVI) and Scale-Content Validity Index (S-CVI) were calculated. The STAR Parenting module demonstrated high content validity. The overall content validity percentage was 86.1%. The I-CVI scores ranged from 0.80 to 1.00. The S-CVI/Ave was 0.88, exceeding the acceptable threshold of 0.80. The inter-rater reliability (ICC(2,k)) was 0.65, indicating moderate consistency among experts. Feedback was incorporated to refine the module's language, clarity, and cultural relevance. The STAR Parenting module is a valid and reliable psycho-educational tool. It is ready for pilot implementation and holds promise for empowering parents in Malaysia to better support the socio-emotional development and school readiness of their children with CIs.

Keywords: Children with cochlear implants; parenting module; socio-emotional development

RP044

Musculoskeletal Pain and Quality of Life among Secondary School Teachers in Kinta Utara District

NUR ASMA LAILEE NISAE ABDULLAH¹, FAIRUS FARIZA ZAINUDIN^{1*},
NOR FADILA KASIM², MON REDEE SUT TXI²

¹*Department of Health Science, Faculty of Sports Science and Coaching, Universiti Pendidikan Sultan Idris,
35900 Tanjong Malim, Perak, Malaysia*

²*Department of Coaching Science, Faculty of Sports Science and Coaching, Universiti Pendidikan Sultan Idris,
35900 Tanjong Malim, Perak, Malaysia*

**Correspondence: fairus.fz@fsskj.upsi.edu.my*

ABSTRACT

Musculoskeletal pain shows a significant occupational health concern among teaching professionals, with prevalence rates ranging from 39% to 95%. This condition markedly impacts job performance, increases absenteeism, and diminishes quality of life (QOL). Despite global evidence suggesting higher vulnerability among female teachers, research addressing this issue within the Malaysian educational context remains limited. This study studied the patterns of musculoskeletal pain and QOL among secondary school teachers in the Kinta Utara district, with specific focus on gender differences. A cross-sectional study was conducted involving 200 secondary school teachers (35% male, 65% female) in the Kinta Utara district through simple random sampling. Ethical approval obtained with written informed consent prior to data collection. Validated instruments including the Standardised Nordic Musculoskeletal Questionnaire (NMQ) and the Health-Related Quality of Life Short Form-12 (HRQOL-SF12). Statistical analyses used descriptive statistics, comparative analysis (t-test, chi-square), and Pearson correlation analysis (r). Shoulder pain emerged as the most prevalence complaint (45%), followed by neck pain (38.5%), back pain (34%), and knee pain (32.5%), with feet pain being the least reported (10.5%). Gender-based analysis revealed significantly higher musculoskeletal pain prevalence among female teachers (49%) compared to their male counterparts (25%). However, no significant gender differences were observed in Physical Component Summary (PCS) and Mental Component Summary (MCS) scores. Correlation analysis demonstrated significant negative associations between the number of pain sites and QOL, particularly for PCS ($r = -0.493$) and MCS ($r = -0.252$), indicating that increased pain burden corresponds with diminished physical and mental well-being. Musculoskeletal pain negatively impacts teachers' quality of life, with implications for both physical and mental health domains. Targeted pain management and musculoskeletal prevention strategies should be developed and implemented to improve teachers' quality of life and musculoskeletal health outcomes

Keywords: Mental health; musculoskeletal pain; occupational health; physical health; quality of life; teachers

RP045

A Comprehensive Predictive Model for Cardiovascular Disease: Integrating Logistic Regression and Classification and Regression Tree for Early Risk Evaluation

AZAH SUMAIYAH ABDUL AZIZ, ZALINA MOHD ALI*

Department of Mathematical Sciences, Faculty of Science and Technology, Universiti Kebangsaan Malaysia

43600 Bangi, Selangor, Malaysia

**Correspondence: zalina99@ukm.edu.my*

ABSTRACT

Cardiovascular disease (CVD) remains a leading cause of global mortality, placing a continuous burden on healthcare systems and requiring more efficient predictive models for early detection. This study aims to identify key health and lifestyle factors associated with CVD and to develop an effective risk classification model using a thorough statistical approach. The analysis employed secondary data from Kaggle, encompassing over 60,000 individuals aged 40 and older in Russia. Russia was chosen because of its extensive, high-quality dataset and significant prevalence of cardiovascular disease, rendering it suitable for multivariate modeling and risk assessment. Descriptive statistics were initially utilised to analyse the distribution of cardiovascular disease among demographic groups. Binary logistic regression was employed to evaluate the influence of health and lifestyle risk factors on the likelihood of CVD. A decision tree approach using the Classification and Regression Tree (CART) algorithm was implemented to enhance predictive accuracy and interpretability by classifying individuals into distinct risk categories. The findings indicate that systolic and diastolic blood pressure, cholesterol levels, body mass index, age and alcohol consumption are significant predictors of CVD, while physical activity offers a protective advantage. Glucose level was recognised as a confounding variable, indicating that its influence on cardiovascular disease is mediated by other primary predictors. The CART model designated systolic blood pressure as the primary splitting variable, followed by cholesterol level, enabling the efficient categorisation of individuals into high-risk groups. This study demonstrates that integrating logistic regression with CART provides a robust and practical framework for predicting cardiovascular risk. The proposed hybrid approach improves classification accuracy and promotes interpretability, providing useful insights for early intervention methods and evidence-based public health policy development.

Keywords: Cardiovascular disease; classification and regression tree; logistic regression; risk factors; risk prediction

RP046

Factors and Associated Health Outcomes of Food Insecurity among Patients with Cancer

SAMIHAH ASLAMIAH AHMAD NAJDI¹, SUHANA YUSAK², ROS SUZANNA AHMAD BUSTAMAM³, CARYN CHAN MEI HSIEN^{1*}

¹Center for Community Health Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Institut Kanser Negara, 62250 Putrajaya, Malaysia

³Department of Radiotherapy and Oncology, Hospital Kuala Lumpur, 50586 Kuala Lumpur, Malaysia

*Correspondence: caryn@ukm.edu.my

ABSTRACT

Food insecurity is a critical yet often overlooked challenge for vulnerable populations, including patients with cancer, impacting their health and well-being. This study aimed to identify the sociodemographic and clinical factors associated with food insecurity among patients with cancer. A cross-sectional, multi-centre study using convenience sampling employed non-parametric tests, primarily the Kruskal–Wallis H test and Chi-square test, to analyse food insecurity [measured using the 8-item Food Insecurity Experience Scale (FIES)] among patients with cancer. Significant findings revealed that household income was a key factor, with patients in the B40 population (earning below RM4,849) demonstrating a statistically significant difference ($H(2) = 9.07$, $p = 0.011$). Findings also suggested higher food insecurity among rural participants ($p = 0.063$), those with lower education levels, individuals not working or retired, and patients with head and neck or skin cancers, though these were not statistically significant. Importantly, a moderate, statistically significant correlation between food insecurity and financial toxicity ($\rho = -0.460$, $p < 0.001$) was observed, with a highly significant Chi-square association ($p < 0.001$) confirming that greater financial toxicity is associated with greater food insecurity. Furthermore, food insecurity was significantly associated with higher levels of anxiety ($\rho = 0.195$, $p = 0.008$; χ^2 $p = 0.040$) and depression ($\rho = 0.331$, $p < 0.001$; χ^2 $p < 0.001$). These findings underscore that financial vulnerability is a significant driver of food insecurity among patients with cancer, which in turn exacerbates financial toxicity and negatively impacts psychological well-being. Therefore, comprehensive screening and targeted support programs are crucial to mitigate food insecurity and its debilitating consequences in this vulnerable patient population.

Keywords: Cancer; food insecurity; patients

RP047

Assessing the Effectiveness of the DemensiaKITA Mobile Health Application on Caregiver Knowledge, Attitude, Practice and Burden in Selangor, Malaysia

NURULIZZAH MAHFAR^{1*}, NIK NAIRAN ABDULLAH¹, DALILA ROSLAN¹,
KHALID IBRAHIM¹, UNGKU AHMAD AMEEN UNGKU MOHD ZAM²,
NORALIZA NOORDIN MERICAN³, NURUL SYAIREEN AB RASHID¹,
CHEN XIN WEE¹, MARIAM MOHAMAD¹

¹Department of Public Health Medicine, Faculty of Medicine, Universiti Teknologi MARA, 47000 Sungai Buloh, Selangor, Malaysia

²Tengku Ampuan Rahimah Hospital, 41200 Klang, Selangor, Malaysia

³Ministry of Health, 62590 Putrajaya, Malaysia

*Correspondence: izzahmahfar@gmail.com

ABSTRACT

Dementia is an increasing public health concern globally and in Malaysia, driven by rapid population ageing. In Malaysia, care is largely provided by informal caregivers who often experience high burden and gaps in knowledge, attitudes and practices (KAP). Mobile health (mHealth) applications offer a promising approach for caregiver support; however, evidence on their effectiveness remains limited. This study aimed to evaluate the effectiveness of the DemensiaKITA mHealth application in improving caregivers' KAP and reducing caregiver burden among dementia caregivers in Selangor, Malaysia. A non-randomised controlled trial was conducted in four public hospitals in Selangor, Malaysia. A total of 100 informal dementia caregivers were allocated into intervention (n = 50) and control (n = 50) groups and assessed at baseline, 1 month, and 3 months. Data were analysed using descriptive statistics and the Generalised Estimating Equations model. The mean age was comparable between groups (intervention: 50.84 ± 14.01 ; control: 48.44 ± 11.72), with no significant difference ($t(98) = 0.929$, $p = 0.355$). Most participants were female, Malay, married, and children of people living with dementia. The intervention significantly improved knowledge, with group time interactions at 1 month ($B = 1.640$, $p = 0.006$) and 3 months ($B = 2.340$, $p = 0.027$). Practice (caregiving difficulty) also improved significantly, with interaction effects at 1 month ($B = -2.440$, $p = 0.003$) and 3 months ($B = -2.320$, $p = 0.005$). Caregiver burden decreased significantly, with interaction effects at 1 month ($B = -2.580$, $p = 0.025$) and 3 months ($B = -3.860$, $p < 0.001$). No significant changes were observed in attitude ($p > 0.05$). The DemensiaKITA mHealth application is a promising and effective intervention for improving knowledge, enhancing caregiving practice, and reducing caregiver burden, although it did not significantly influence attitudes in the 3-month study period. It has a strong potential to support dementia caregivers in Malaysia.

Keywords: Attitudes; caregivers; caregiver burden; dementia; health knowledge; mobile applications; practice

RP048

Policy and Institutional Perspectives on the Use of Schools as Temporary Evacuation Centres during Floods in Southeast Asia

NOR AINSYAFIKAH MADIRAN, AZIEMAH ZULKIFLI*, AIDA SORAYA
SHAMSUDDIN, GOH CHOO TA, NURFASHAREENA MUHAMAD,
MUHAMMAD AMIRULAMRI MOHD AKHAIRI

Institute for Environment and Development, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

**Correspondence: aziemah_zul@ukm.edu.my*

ABSTRACT

Southeast Asia's high flood vulnerability frequently necessitates repurposing schools as temporary evacuation centres (TECs). However, balancing emergency response with educational continuity remains a challenge. This research aims to examine existing policies on schools as TECs, identify institutional roles and challenges, and propose recommendations for improved coordination. Using a qualitative comparative case study of Malaysia, Thailand, and Indonesia, this study reviewed government decrees, disaster guidelines, and academic literature. Key documents included Malaysia's NADMA Directive No. 1 and MKN No. 20, Thailand's OBEC disaster management and school safety guidelines, and Indonesia's Law No. 24 of 2007, BNPB Regulation No. 4 (2012), Ministerial Circular Letter No. 70a/MPN/SE/2010 and SNI 7937:2013. Findings indicate that school activation is governed by layered administrative frameworks rather than clear statutory instruments, leading to regulatory ambiguity. While Malaysia and Thailand rely on institutional directives, Indonesia has the most integrated framework, incorporating disaster risk reduction into educational governance. Despite these structures, all three countries face common challenges, including a lack of explicit school designation, limited teacher capacity, financial constraints, and inadequate planning for educational continuity. Significant regulatory gaps remain in managing the dual-use function of schools. Strengthening statutory designations, inter-agency coordination, and continuity planning is essential to support more resilient school-based disaster management.

Keywords: Disaster management; floods; schools; temporary evacuation centres

RP049

Effects of Blue and Ultraviolet Filtering Eye Drops on Contact Lens Light Transmittance

SADEQ NAEEM AJEENA*, HALIZA ABDUL MUTALIB, BASHIRAH ISHAK

*Programme of Optometry and Visual Science, Faculty of Health Sciences, Universiti Kebangsaan Malaysia,
50300 Kuala Lumpur, Malaysia*

**Correspondence: p146415@siswa.ukm.edu.my*

ABSTRACT

Blue and ultraviolet (UV) light can cause oxidative stress and visual fatigue in the eyes. Contact lenses (CLs) with blue and UV light filters can mitigate the effects of high-energy radiation. Filtered eye drops may provide additional protection. The filtered eye drop Miren® can be used concurrently with the CLs. The effect on blue and UV light transmittance by this combination has not been evaluated. This study investigated the effectiveness of the combination of filtered eye drops with different types of CL. An in vitro experiment evaluated the impact of blue and UV light transmittance of a combination of filtered eye drops (Miren®) with various CLs. Six commercially available soft CLs were tested: two clear contact lenses (CCLs), two blue light filtered contact lenses (BLCLs), and two UV filtered contact lenses (UVCLs). Transmitted light intensity was measured using the MK350S Premium spectrometer. Blue light was emitted using a 6.7-inch mobile device, while UV light was generated using a Smile Shark torch light. Baseline intensity was recorded prior to which the intensity was measured separately through each CL. The process was repeated after applying the filtered eye drops. Descriptive statistics, paired-sample t-test, and one-way ANOVA were used to analyse the data. The application of filtered eye drops significantly reduced blue and UV light transmittance across all tested CLs ($p < 0.001$). Under BL, no significant differences were observed between lens types after drop application ($p = 0.17$), whereas a significant difference was detected under UV light ($p = 0.028$). The magnitude of attenuation varied by lens type and light source. CCLs combined with drops showed the greatest additional reduction in transmittance (9.2–10.3% under BL; 7.0–8.2% under UV light). BLCLs combined with drops showed the greatest enhancement in overall filtering efficacy (26.2% total reduction), with the smallest incremental attenuation (4.9–6.8% under BL; 6.6–7.0% under UV light). UVCLs exhibited moderate attenuation, ranging from (7.4–8.4% under BL), and (6.4–7.8% under UV light). Filtered eye drops significantly enhance the light-filtering capabilities of CLs, with the magnitude of improvement depending on the lens type and light source, with greatest effect observed with BLCLs. This approach is particularly effective with clear lenses, suggesting a promising strategy for improving ocular protection against blue and UV light.

Keywords: Blue light; contact lens; filtering eye drop; light transmittance; UV light

RP053

Association of Sociodemographic and Environmental Factors of Tuberculosis Cases from 2014 to 2023 in Malaysia

NUR ADIBAH MOHIDEM^{1*}, SHARIFAH NORKHADIJAH SYED ISMAIL²

¹Public Health Unit, Department of Primary Health Care, Faculty of Medicine and Health Sciences, Universiti Sains Islam Malaysia, Bandar Baru Nilai, 71800 Nilai, Negeri Sembilan, Malaysia

²Department of Environmental and Occupational Health, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

*Correspondence: nuradibahmohidem@usim.edu.my

ABSTRACT

Tuberculosis (TB) is a major public health issue globally, and it remains one of the deadliest infectious diseases in Malaysia. This cross-sectional study aimed to determine the spatiotemporal distribution of TB cases with the association of sociodemographic and environmental factors in Malaysia. The sociodemographic data of 301,822 TB cases such as age, gender, nationality, country of origin, race, educational level, healthcare worker status, smoking status, and drug consumption from 1st January 2014 to 31st December 2023 in Malaysia district were collected from the Tuberculosis Information System (TBIS) database. Environmental data such as sulphur dioxide (SO₂), particulate matter 10 (PM₁₀), ozone (O₃), nitrogen dioxide (NO₂), carbon monoxide (CO), wind speed, relative humidity, temperature, rainfall, and atmospheric pressure were obtained from the Department of Environment Malaysia and Malaysia Meteorological Department. Ordinary least squares (OLS) and geographically weighted regression (GWR) models were employed to determine the spatial association of sociodemographic and environmental factors with the TB cases across districts in the country. Sociodemographic factors such as being aged less than 14 years old, country of origin from Nepal, country of origin from Thailand, Chinese racial, Indian racial, primary educational level, and tertiary educational level, as well as the environmental factor which is PM₁₀, were associated with TB cases ($p < 0.05$). The GWR model based solely on the sociodemographic factor i.e. GWR1 was the best model to determine the spatial distribution of TB cases based on the highest R² value i.e. 0.99. The maps of estimated local coefficients in GWR models reflected the spatial variability of sociodemographic and environmental factors on TB cases. This study concluded that the spatial analysis is crucial for identifying regions with a high TB burden based on characteristics related to the disease.

Keywords: Environment; geographically weighted regression; sociodemographic; spatial; tuberculosis

OR003

Watch, Learn, Improve: Video-Based Slit-Lamp Training Enhances Immediate Test Performance and Reveals High Instructional Motivation

KHADIJA DAYAH ABDULLAHI, MIZHANIM MOHAMAD SHAHIMIN*,
BARIAH MOHD ALI

*Optometry and Vision Science Program, Centre for Community Health Studies, Faculty of Health Sciences,
Universiti Kebangsaan Malaysia, Kuala Lumpur 50300, Malaysia*

**Correspondence: mizhanim@ukm.edu.my*

ABSTRACT

Digital video-based learning (VBL) has become a core innovation in rehabilitation and health sciences education, enabling scalable and standardised delivery of clinical skills training. However, empirical evidence supporting its effectiveness in optometry education and the role of learner motivation in digital environments remains limited. This study evaluated the effectiveness of slit-lamp VBL in enhancing immediate knowledge acquisition and explored the role of instructional motivation, measured using the Reduced Instructional Materials Motivation Survey (RIMMS), among undergraduate optometry students. Forty-four Year 3 Optometry students completed a written pre-test before engaging with digital slit-lamp instructional videos, followed by an immediate post-test and the RIMMS. Descriptive statistics, Wilcoxon signed-rank test, Hake normalised learning gains, Spearman's correlation, simple linear regression, and hierarchical multiple regression were performed to evaluate learning outcomes and the contribution of instructional motivation. Post-test scores (80.30 ± 14.55) were significantly higher than pre-test scores (63.13 ± 12.63 ; $Z = 5.051$, $p < 0.001$), demonstrating substantial immediate learning gains with a large effect size ($r = 0.76$) and standardised mean difference ($d = 1.27$). Under the Hake framework, 63.6% of students achieved medium-to-high normalised gains. Baseline pre-test scores did not significantly correlate with normalised gains ($\rho = 0.033$, $p = 0.833$; regression $p = 0.784$), indicating VBL was similarly effective across varying student baseline competency levels. Homogeneous, elevated instructional motivation scores were reported (mean = 4.4/5.0). However, RIMMS scores did not significantly correlate with normalized gains ($\rho = -0.09$, $p = 0.558$), or predict post-test performance when controlling for baseline knowledge ($\Delta R^2 = 0.001$, $p = 0.890$). Digital video-based slit-lamp instruction significantly enhances immediate clinical knowledge acquisition in optometry training, and appears to be an equitable preclinical learning strategy with high reported learning motivation. These findings support the integration of digital motivation-sensitive instructional design to optimise clinical skills development in rehabilitation and health sciences education.

Keywords: Clinical skills training; learning motivation; optometry education; RIMMS; video-based learning

OR004

Fear of Falling and Self-Efficacy for Physical Activity among Stroke Survivors with Home-Based Physiotherapy Experience

NITYA LETCHUMANAN^{1*}, ROSLIZAWATI NAWAWI¹, ISYAN FARAHIIN ISMAIL¹, NOR AZURA AZMI²

¹Physiotherapy Programme, School of Integrative Medicine & Life Sciences, KPJ Healthcare University, 71800 Nilai, Negeri Sembilan, Malaysia

²Physiotherapy Programme, Centre for Rehabilitation and Special Needs Studies, Faculty of Health, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

*Correspondence: aytin17@gmail.com

ABSTRACT

Despite physical recovery after stroke, fear of falling (FOF) continues to shape survivors' perceived capability to engage in physical activity. The perceived capability is reflected in self-efficacy for physical activity (SEPA), which influences independence. However, limited evidence has examined the prevalence of FOF and its association with SEPA among community-dwelling stroke survivors who underwent home-based physiotherapy. Therefore, this study aimed to determine the prevalence of FOF and SEPA and examine their association among stroke survivors. A cross-sectional study was conducted among 102 community-dwelling stroke survivors in the Klang Valley with prior experience of home-based physiotherapy, aged 41-79 years (mean 57.33 ± 8.68), with 69.6% males and 30.4% females. FOF and SEPA were assessed using self-administered Falls Efficacy Scale-International (FES-I) (scores: 16-19 low, 20-27 moderate, 28-64 high) and the Stroke Self-Efficacy Questionnaire (SSEQ), respectively, during the period from June 2025 to January 2026. FOF prevalence and SEPA scores were summarised using descriptive statistics, and the association between FOF and SEPA was analysed using Spearman's rank correlation analysis, with statistical significance determined at $p < 0.05$. The study showed that home-based physiotherapy stroke survivors were predominantly ischaemic (66.7%) and 54.9% reported at least one fall. High FOF was most prevalent (72.5%; FES-I 44.54 ± 9.82), followed by moderate FOF (22.5%; FES-I 23.61 ± 2.52) and low FOF (4.9%; FES-I 17.40 ± 1.52). SEPA showed marked variability (83.17 ± 30.99), suggesting differences in confidence to be physically active. FOF and SEPA were statistically associated, with a strong inverse relationship ($r_s = 0.666$, $p < 0.001$), indicating that greater concern about falling is linked to lower physical activity confidence. These findings highlight the need to address FOF alongside physical training in home-based and community stroke rehabilitation practice to support functional independence among stroke survivors.

Keywords: Fear of falling; home-based physiotherapy; physical activity; self-efficacy; stroke

OR006

Sleeping in on Weekends: Prevalence and Associations with Anthropometric Status and Academic Performance among Primary School Children in Selangor

DHANASRIY SATHIABALAN¹, FATIN HANANI MAZRI^{1*}, ZAHARA ABDUL MANAF¹, ARIMI FITRI MAT LUDIN², HASLINA ABDUL HAMID¹, NUR'AIN MARDHIYAH HARUN¹

¹Dietetics Program, Centre for Healthy Ageing and Wellness, Faculty of Health Science, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Biomedical Science Program, Centre for Healthy Ageing and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Kuala Lumpur 50300, Malaysia

**Correspondence: fatinhananimazri@ukm.edu.my*

ABSTRACT

Sleep deprivation is increasingly common in the modern world, leading to sleep extension on weekends to compensate for accumulated weekday sleep debt. This cross-sectional study aims to determine the prevalence of weekend catch-up sleep (WCUS) and its associations with anthropometric status and academic performance among primary school children in Selangor. This study involved 559 children recruited from urban and rural schools between June and August 2025. Data on sociodemographic characteristics, sleep parameters, screen time, physical activity, school attendance, cognitive performance, and classroom-based assessment (PBD) scores were collected. Anthropometric status was assessed using body mass index-for-age (BMI-for-age) and height-for-age indicators. Logistic and multiple linear regression analyses were performed to examine the associations between WCUS duration, anthropometric status, and academic performance. Overall, 75.7% (n = 423) of children extended their sleep on non-school days. Among WCUS practitioners, 26.7% reported >0–1 hour, 20.1% >1–2 hours, 18.9% >2–3 hours, and 34.3% ≥3 hours. Children reporting >2–3 hours of WCUS had significantly higher odds of having a normal BMI-for-age compared with those reporting 0 hours of WCUS (OR = 2.33; 95% CI: 1.02–5.31; p = 0.044). However, no significant association was observed between WCUS and height-for-age. Regarding academic performance, WCUS of <1 hour was significantly associated with higher Malay language PBD scores (β = 0.619; 95% CI: 0.244–0.993; p = 0.001), whereas WCUS of 1–2 hours was linked to higher Mathematics PBD scores (β = 0.401; 95% CI: 0.124–0.678; p = 0.005), compared with WCUS ≥3 hours. Furthermore, shorter WCUS durations (<2 hours) were significantly associated with higher school attendance compared with WCUS ≥3 hours. In conclusion, WCUS is highly prevalent among primary school children in Selangor. The associations vary by duration, emphasising the role of sleep patterns in supporting child health and academic performance.

Keywords: Academic performance; body weight; children; sleep extension; weekend catch-up sleep

OR007

Dietary Management of Disorders of Gut Brain Interactions: Knowledge, Attitudes and Practices among Gastroenterologists and Dietitians in Asia

SHANTHI KRISHNASAMY^{1*}, WEDSTER SCHUMARKER¹, FAZRYN SHAHIRA¹, KEWIN TIEN HO SIAH², SANJIV MAHADEVA³, KEE HUAT CHUAH³, MASUDUR RAHMAN⁴, HANG DAO VIET⁵, BRIDGETTE WILSON⁶

¹*Dietetics Department, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Department of Medicine, Yoo Soo Lin School of Medicine, National University of Singapore, 10 Medical Dr Singapore 117597 Singapore*

³*Department of Gastroenterology and Hepatology, Faculty of Medicine, Universiti Malaya, 50603 Kuala Lumpur, Malaysia*

⁴*Department of Gastroenterology, Sheikh Rusell National Gastroenterology Institute and Hospital, 1212 Dhaka, Bangladesh*

⁵*Department of Gastroenterology, Hanoi Medical University, Kim Lien Ward, V Dong Da District, Hanoi 100000, Vietnam*

⁶*Nutrition and Dietetics Department, Singapore Institute of Technology, 828608 Singapore*

**Correspondence: shanthikrishnasamy@ukm.edu.my*

ABSTRACT

Dietary management of disorders of gut-brain interaction (DGBI), particularly irritable bowel syndrome and functional dyspepsia, is complex. Since the implementation of evidence-based dietary therapies for DGBI has not been extensively studied in Asia, this study aimed to evaluate the knowledge, attitudes, practices, and perceived barriers among Asian gastroenterologists and dietitians regarding the dietary management of patients with DGBI. A cross-sectional face-to-face and online survey was conducted among gastroenterologists and dietitians across 8 Asian countries through websites, conferences, social media platforms and dietetic networks using convenience sampling. Data were analysed using SPSS version 27, with results presented as frequencies and means, and Spearman correlation analysis was used to determine the correlation between knowledge and attitudes. A total of 110 dietitians and 54 gastroenterologists with an average age of 36.00 ± 6.134 and 39.18 ± 6.886 years, completed the survey. Moderate knowledge was observed in 53.3% of dietitians, while 46.2% of gastroenterologists had unsatisfactory knowledge. Positive attitudes towards dietary interventions for DGBI were noted in 75.0% of dietitians and 60.4% of gastroenterologists. Although a large number of participants were aware of evidence-based dietary management for DGBI, this was not reflected in their practice. The primary barrier identified was a lack of resources and guidelines. No significant relationship was found between years of clinical experience and knowledge or attitudes ($p > 0.05$). Despite limited knowledge in managing DGBI, most dietitians and gastroenterologists have a positive attitude towards the importance of dietary

intervention. However, the implementation of dietary management was not according to evidence-based practice. Addressing identified barriers is essential to improving dietary management. Further research is needed on Asian DGBI treatment strategies. Moving forward, there is a strong need for culturally tailored guidelines, enhanced professional training, and greater international collaboration to improve dietary management of DGBI in Asian populations.

Keywords: Dietary management; functional dyspepsia; irritable bowel syndrome; low FODMAP diet

OR008

Sleep Patterns, Social Jetlag, Screen Time and Their Associations with School Attendance and Academic Performance among Primary School Children in Selangor, Malaysia

NUR'AIN MARDHIYAH HARUN¹, FATIN HANANI MAZRI^{1*}, ARIMI FITRI MAT LUDIN², ZAHARA ABDUL MANAF¹, HASLINA ABDUL HAMID³, DHANASRIY SATHIABALAN¹

¹*Dietetics Programme and Centre for Healthy Ageing and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Biomedical Science Programme and Centre for Healthy Ageing and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

³*Dietetics Programme and Centre for Community Health Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

*Correspondence: fatinhananimazri@ukm.edu.my

ABSTRACT

Sleep deficiency has become a growing global health concern among school-aged children, estimated 25% to 50% of children fail to achieve the recommended 9-11 hours of sleep per night globally. Beyond sleep duration, sleep-wake timing and digital screen exposure influence circadian alignment-the synchronisation of behavioural patterns with the body's internal biological clock. This study aimed to examine sleep-wake patterns during school and non-school days, daytime sleepiness and screen time, and their associations with academic performance and attendance among primary school children in Selangor, Malaysia. A cross-sectional study was conducted among 559 children aged 10-12 years old (54.7% boys; 81.8% Malay; 52.1% from urban schools). Data on sociodemographic characteristics, sleep patterns, daytime sleepiness, screen time, school attendance, and classroom-based assessment (PBD) scores were collected using structured questionnaires and school records. The average weekly sleep duration was 8.10 ± 0.57 hours, below recommended levels. Children reported significantly later sleep onset (23:19 vs 22:20, $p < 0.001$), sleep offset (08:34 vs 06:04, $p < 0.001$) and midpoint of sleep (03:18 vs 02:12, $p < 0.001$) during non-school days compared with school days, alongside longer sleep duration on non-school days (9.15 vs 7.44 hours, $p < 0.001$). Approximately 40.3% of children experienced more than 2 hours of social jetlag, while 46.9% reported high daytime sleepiness. Screen time was also significantly longer on non-school days compared with school days (6.15 vs 3.46 hours, $p < 0.001$). Greater social jetlag was associated with lower school attendance ($r = -0.10$, $p = 0.015$) and lower Malay language PBD scores ($r = -0.12$, $p = 0.004$). Higher daytime sleepiness score and longer screen time were associated with lower Math PBD scores ($r = -0.10$, $p = 0.018$; $r = -0.133$, $p = 0.002$). These findings suggest that sleep patterns and digital behaviours are important lifestyle factors linked to school engagement and academic performance. In modern societies, promoting healthy sleep habits and responsible screen use are important strategies to support children's learning and well-being.

Keywords: Academic performance; school children; screen time; sleep; social jetlag

OR009

Developmental Eye Movement Test Scores among Chinese Children in Klang Valley

SUMITHIRA NARAYANASAMY*, LOW LAY KHIM, MOHD IZZUDDIN HAIROL

Optometry & Vision Science Programme, Centre for Community Health Studies, Faculty of Health Science, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

**Correspondence: sumithira@ukm.edu.my*

ABSTRACT

Developmental eye movement (DEM) test is one of the most common tests that can be used to evaluate the oculomotor skills quantitatively and screen for reading problems. Previous studies have shown that DEM test scores are influenced by differences in languages and culture. Currently, there is lack of research on the impact of bilingual or multilingual on the DEM test scores. This study was aimed to explore the DEM performance among Chinese children in Klang Valley and investigate the differences in DEM test performance among Chinese children in three languages, which are Mandarin, Malay and English. Besides, the scoring of the DEM test of Malaysian Chinese children were compared with the normative data of other populations. The relationship between DEM test score and reading was also determined in this study. A total of 102 Chinese children in Klang Valley aged from 7-11 years old were recruited. The scores for vertical, adjusted horizontal and ratio components were computed. The results showed there were significant differences in the DEM scores in three languages and populations from other countries. The results showed most similar with Italian and followed by American English, China and Hong Kong. These showed that DEM scores may be affected by differences in languages, culture and educational systems among various populations. Adjusted horizontal scores and reading speed were positively correlated. This shows that DEM test can be used to screen children with reading problems. Further investigation of norms of DEM scores for Chinese children in Malaysia is needed.

Keywords: Chinese children; developmental eye movement test; oculomotor; reading performance

OR010

Translational Potential and Explainability of Artificial Intelligence Decision Support for Adults in Intensive Care: A Scoping Review

YODHA PRANATA^{1,2*}, YUSUF ACHMAD BAHTIAR³, HESTI PURWANTI⁴

¹Faculty of Nursing, Universitas Padjadjaran, Sumedang, West Java 45363, Indonesia

²Data Science Program, Faculty of Science and Technology, Universitas Terbuka, Tangerang Selatan, Banten 15418, Indonesia

³Department of Anesthesiology and Intensive Care, Ngudi Waluyo Regional General Hospital, Blitar, East Java Timur 66184 Indonesia

⁴Department of Internal Medicine, Ngudi Waluyo Regional General Hospital, Blitar, East Java 66184, Indonesia

*Correspondence: yodha24001@mail.unpad.ac.id

ABSTRACT

The intensive care unit (ICU) demands rapid, high-stakes decision-making. Although artificial intelligence (AI) offers superior predictive accuracy over traditional scoring, its “black-box” nature remains a barrier to clinical adoption. This scoping review systematically maps the translational potential and the characteristics of explainable artificial intelligence (XAI) strategies in AI/ML-based clinical decision support tools for adult ICU settings. The review followed the PRISMA-ScR guidelines. We conducted systematic searches of Web of Science, PubMed, Scopus and EBSCOhost for studies published up to January 2026. Data were synthesised using a narrative, descriptive approach with thematic mapping to characterise model architectures, XAI strategies, translational potential, and proposed clinical utility. A total of 808 records were identified, and 30 studies met the inclusion criteria. Most studies (83.3%) used tree-based ensemble models, with Shapley Additive exPlanations (SHAP) as the predominant XAI strategy (86.7%). A geographic imbalance was observed: 73.3% of studies originated from Asia (predominantly China), yet 80% relied heavily on North American public datasets (MIMIC/eICU). Translational outputs most commonly took the form of web-based prototypes (66.7%), whereas integration into electronic health record workflows was reported in only 16.7% of the studies. AI in adult ICU decision support is shifting from “black-box” prediction toward more transparent, clinician-centered systems. Web-based prototypes facilitate early access and usability testing, but full integration into routine clinical workflows remains a key challenge. Future research should prioritise automated data pipelines and prospective validation to enable reliable, real-world implementation of AI-driven predictions in everyday critical care practice.

Keywords: Artificial intelligence; clinical decision support; explainable AI; intensive care unit; translational research

OR011

Cross-Linguistic Acoustic Markers of Hearing Impairment: A Review of Vowel Production and Formant Centralisation

NUR 'AZZAH ZAKARIA^{1,4}, AKMALIZA ALI², CHONG FOONG YEN²,
JULIANA AMINAH MARHABAN^{1,4}, SITI MUNIRAH HARIS³, WAN
ASLYNN WAN AHMAD^{1,4*}

¹Department of Audiology and Speech-Language Pathology, Kuliyah of Allied Health Sciences, IIUM Centre for Natural & Health Sciences, Kuantan Campus, 25200 Bandar Indera Mahkota, Kuantan, Pahang, Malaysia

²Audiology Program, Centre for Rehabilitation and Special Needs Studies, Faculty of Health Sciences, National University of Malaysia, Kuala Lumpur Campus, 50300 Kuala Lumpur, Malaysia

³Speech Sciences Program, Centre for Rehabilitation and Special Needs Studies, Faculty of Health Sciences, National University of Malaysia, Kuala Lumpur Campus, 50300 Kuala Lumpur, Malaysia

⁴Speech & Language Science Collaboratives, Kuliyah of Allied Health Sciences, IIUM Centre for Natural & Health Sciences, Kuantan Campus, 25200 Bandar Indera Mahkota, Kuantan, Pahang, Malaysia

*Correspondence: wanaslynn@iium.edu.my

ABSTRACT

Vowel production serves as a critical acoustic indicator of speech development and auditory feedback efficiency. International literature consistently reports that individuals with hearing impairment exhibit reduced vowel space area, centralised formant patterns, and weakened articulatory contrast compared to typical hearing peers. These acoustic measures, specifically the first (F1) and second (F2) formants are increasingly utilised as objective markers of speech clarity and rehabilitation outcomes. This review synthesises global findings on vowel acoustics in hearing-impaired populations, focusing on the universal phonetic shifts observed across various linguistic systems. A synthesis of global comparative studies reveals that hearing impairment leads to systematic formant centralisation in Indo-European and Sinitic languages, this is characterised by a significant compression of the F2 range, representing restricted horizontal tongue movement, and a reduced F1 range, reflecting limited vertical articulatory displacement. While normal-hearing populations exhibit distinct F1/F2 clusters that vary by gender due to vocal tract anatomy, hearing-impaired populations show a universal trend toward a “neutralised” acoustic space. Although F1 and F2 acoustic markers are well-documented in Western and East Asian phonetic systems, a critical gap remains in the standardisation of these measures for the Malay language. This research void is particularly significant for clinical practice in Malaysia, where the lack of normative data prevents a precise comparison between typical-hearing individuals and the hearing-impaired population. Given the compact nature of the Malay six-vowel system, existing international benchmarks may not accurately detect the specific patterns of formant overlap and vowel centralisation unique to Malay-speaking patients. Therefore, establishing Malay-specific acoustic baselines is a vital prerequisite for conducting meaningful comparative studies, ensuring diagnostic accuracy, and optimising objective speech rehabilitation outcomes for the hearing-impaired community.

Keywords: Clinical rehabilitation; hearing impairment; phonetic gap; speech science; vowel acoustics

OR012

Karanjin Exerts Anti-Neuroinflammatory Activity in Lipopolysaccharide-Induced BV-2 Microglial Cells by Reducing Nitric Oxide Production and Targeting Key Inflammatory Mediators

AZIATUL MUNIRAH SHAFIE¹, YUAN SENG WU^{1,2*}, SANG LOON TAN³,
AMAR DAUD ISKANDAR ABDULLAH¹, SAIDUR RAHMAN⁴

¹Department of Biomedical Sciences, Sir Jeffrey Cheah Sunway Medical School, Faculty of Medical and Life Sciences, Sunway University, 47500 Subang Jaya, Selangor, Malaysia

²Sunway Microbiome Centre, Faculty of Medical and Life Sciences, Sunway University, 47500 Subang Jaya, Selangor, Malaysia

³Sunway Biofunctional Molecules Discovery Centre, Faculty of Medical and Life Sciences, Sunway University, 47500 Subang Jaya, Selangor, Malaysia

⁴Research Centre for Nano-Materials and Energy Technology, Faculty of Engineering and Technology, Sunway University, 47500 Subang Jaya, Selangor, Malaysia

*Correspondence: yuansengw@sunway.edu.my

ABSTRACT

Alzheimer's disease (AD) is a neurodegenerative disorder marked by amyloid-beta plaque and neurofibrillary tangle accumulation, leading to neuronal damage via neuroinflammation. Neuroinflammation involves elevated pro-inflammatory cytokines such as interleukin-6 (IL-6) and tumour necrosis factor alpha (TNF- α), along with mediators including nucleotide-binding domain leucine-rich repeat pyrin domain-containing protein 3 (NLRP3), inducible nitric oxide synthase (iNOS), and beta-site APP cleaving enzyme 1 (BACE1). Current AD drugs show limited efficacy and high toxicity, highlighting the need for effective and safer alternatives. Natural products are promising due to their abundance and lower toxicity. Karanjin, a furan flavonoid from *Pongamia pinnata* seeds, has been reported to exhibit anti-inflammatory properties, but its neuroprotective potential remains underexplored. This study aimed to evaluate the effects of karanjin on lipopolysaccharide (LPS)-induced neuroinflammation in BV-2 microglial cells using in vitro and in silico approaches. Cell viability assessment via 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay revealed that karanjin was non-cytotoxic to BV-2 cells at concentrations up to 12.5 μ M, with a slight reduction in viability at higher doses. Nitrite assay showed that LPS significantly increased nitrite levels (7.8563 μ M) compared to untreated controls (3.8625 μ M, $p < 0.05$). Interestingly, pretreatment with karanjin reduced nitrite production in a dose-dependent manner, with the highest inhibition at 50 μ M (4.0188 μ M; 48.85% inhibition), indicating suppression of nitric oxide synthesis. Molecular docking studies further supported these findings, revealing strong binding affinities of karanjin to key inflammatory mediators, including iNOS (PDB: 4CX7), NLRP3 (PDB: 7ALV), and BACE1 (PDB: 4DJU), with docking scores of -10.64, -9.13 and -8.64 kcal/mol, respectively. These interactions

suggest that karanjin may directly modulate inflammatory pathways relevant to neurodegeneration. Collectively, karanjin demonstrated significant anti-neuroinflammatory effects by inhibiting nitric oxide production and interacting with molecular targets implicated in neuroinflammation, thereby supporting its potential as a natural therapeutic agent for AD that warrants further investigation.

Keywords: Alzheimer's disease; karanjin; neuroinflammation

OR013

Cardiac Microenvironment Communication Under Hypoxic Stress Potentially Mediated by Cardiomyocyte-derived Exosome Delivery of Angiogenic Regulators

JIA ZHENG¹, ARIMI FITRI MAT LUDIN², NOR FADILAH RAJAB², NURUL FARHANA JUFRI^{1*}

¹Centre for Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Centre for Healthy Ageing and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

*Correspondence: nurulfarhana@ukm.edu.my

ABSTRACT

Myocardial ischemia triggers complex intercellular communication that coordinates angiogenic responses within the cardiac microenvironment. However, the way by which cardiomyocytes (CMs) regulate endothelial cell behaviour under hypoxic conditions, particularly through exosome-mediated signalling, remains insufficiently understood. This study investigated whether cardiomyocyte-derived exosomes (CM-exos) modulate angiogenic activity in cardiac microvascular endothelial cells (CMECs) and whether hypoxia enhances this communication through the delivery of angiogenesis-related genetic cargo. A hypoxic CMs model was established in H9C2 cells using cobalt chloride (600 μ M, 24 h). Exos were isolated and characterised, and their effects on CMECs were examined through two experimental systems: (i) CMs/CMECs co-culture and (ii) CMECs direct treatment with isolated CM-exos. Five experimental groups were established in each level, including normoxic or hypoxic CMs/exos with or without GW4869 (the exos secretion inhibitor). Exos uptake was evaluated by immunofluorescence (CD81), while RT-qPCR quantified angiogenesis-related regulators (MALAT1, Klotho, FGF23) and downstream markers (ET-1, NO, vWF, Ang-II). Functional effects on CMECs were assessed using CCK-8 proliferation, flow cytometry, scratch assay, transwell migration assays, and tube formation assays. Both normoxic and hypoxic CM-exos were efficiently endocytosed by CMECs, whereas GW4869 markedly reduced exos uptake. Exposure to CMs or CM-exos significantly increased the expression of MALAT1, FGF23, ET-1, NO, vWF, and Ang-II while suppressing Klotho expression ($p < 0.05$). Functionally, CM-exos enhanced CMECs' viability, migration, and angiogenic tube formation, with hypoxia-derived exos exhibiting stronger pro-angiogenic effects. These responses were consistently attenuated following inhibitor treatment, indicating that exosome-mediated signalling is a key driver of the observed endothelial responses. Collectively, these findings reveal that CMs communicate with CMECs through exos-dependent delivery of angiogenic regulatory molecules, thereby promoting endothelial migration and vascular network formation under hypoxic conditions. This exos-mediated signalling pathway may represent a critical mechanism for adaptive angiogenesis during myocardial ischemic stress and could provide a potential therapeutic target for coronary artery disease.

Keywords: Angiogenesis; coronary artery disease; exosome hypoxia; intercellular communication

OR014

Acceptable Quality Dose in Pediatric Abdominal Computed Tomography: A Multicenter Study Benchmarking against Diagnostic Reference Levels in Malaysia

MOHD SHAHRIL MOHD SHAMSUL^{1*}, AKMAL SABARUDIN¹,
FAIZAH MOHD ZAKI², HAMZAINI ABD HAMID², ROHAZLY ISMAIL³,
VIVIEN SURYANI AMRU⁴

¹Centre for Diagnostic, Therapeutic and Investigative Studies, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur

²Department of Radiology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Cheras, 56000 Kuala Lumpur, Malaysia

³Department of Radiology, Hospital Tunku Azizah Lot 25, 50300 Kuala Lumpur, Malaysia

⁴Department of Radiology, Sabah Women and Children Hospital, Karung Berkunci No 187, 88996 Kota Kinabalu, Sabah, Malaysia

*Correspondence: akmal.sabarudin@ukm.edu.my

ABSTRACT

Radiation dose optimisation in paediatric computed tomography (CT) is crucial to maintain diagnostic image quality while minimising radiation exposure. The Acceptable Quality Dose (AQD) has been proposed as an approach to balance image quality and radiation dose by linking radiation metrics with clinically acceptable image quality. This multicentre retrospective study analysed paediatric patients who underwent abdominal CT examinations. CT dose index (CTDI) and dose length product (DLP) were collected together with image quality assessments. Patients were divided into: neonates (0–30 days), infants (1 month–4 years), toddlers (4–10 years), childhood (10–14 years), and adolescents (14–18 years). Image quality was evaluated using the Image Quality Scoring Criteria (IQSC). The AQD was determined based on examinations that achieved diagnostically acceptable image quality, defined as an IQSC score of 3. A total of 120 paediatric abdominal CTs were evaluated using IQSC and the results showed score 3 (75.8%, n = 91), followed by score 4 (21.7%, n = 26), score 2 (2.5%, n = 3), and score 1 (0). Among the IQSC score 3 (n = 91), the distribution across age groups was as follows: neonates 4 (4.4%), infants 9 (9.9%), toddlers 23 (25.3%), children 30 (33.0%), and adolescents 25 (27.5%). Comparison of CTDI (mGy) and DLP (mGy cm) between AQD (IQSC score 3) and DRLs (75th percentile) demonstrated lower dose values across all age groups: neonates (2.2 and 24 vs 2.2 and 24), infants (1.9 and 103.7 vs 2.3 and 108.3), toddlers (2.9 and 168.3 vs 3.7 and 241), childhood (6.7 and 443.2 vs 7.7 and 444.2), and adolescents (10.3 and 491 vs 12 and 528.2), respectively. The findings demonstrate that diagnostically acceptable image quality in paediatric abdominal CT can be achieved at radiation dose levels below the DRL method, supporting the utility of the AQD approach for optimising paediatric CT imaging protocols.

Keywords: Acceptable quality dose; diagnostic reference levels; multicenter study; pediatric abdominal CT; radiation dose optimisation

OR016

Verification of GATE Modeling and Simulation for Clinac IX Linear Accelerator at 6 MV Photon

MICRAAJ ABDISALAM ADEN, SITI HAJAR ZUBER*, GUNALAN RAMACHANDRAN, MUHAMMAD SAFWAN AHMAD FADZIL

Centre for Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

**Correspondence: hajarzuber@ukm.edu.my*

ABSTRACT

Radiotherapy treatment planning relies heavily on accurate treatment planning to ensure precise delivery of radiation doses to cancerous tissues while minimising harm to healthy organs. This research aims to investigate the efficacy of the Geant4 Application for Tomographic Emission (GATE) in modelling and simulating the Clinac iX linear accelerator for radiotherapy treatment planning, particularly focusing on percentage depth dose (PDD) measurements. This research explored the potential of Monte Carlo GATE as a validation tool in the treatment planning of radiotherapy, establish the geometry of the Clinac iX linear accelerator in GATE, measure the PDD of a 6 MV photon beam, and compare the simulated and measured PDD at 6 MV photon. It was determined that an electron beam energy of 6.4 MeV provides the optimal modelling for the 6 MV photon beam of the Clinac iX linear accelerator. The PDD curve obtained from GATE simulations accurately reproduced the experimental dose distribution, demonstrating excellent agreement with measured data. GATE simulation achieved a 98% passing rate under the 3%/3 mm gamma index criterion, demonstrating its accuracy in dose verification. This research lays the groundwork for investigating the potential of GATE modelling and simulation in radiotherapy treatment planning, with the goal of improving treatment accuracy and patient outcomes.

Keywords: Clinac iX; GATE; Monte Carlo simulation; percentage depth dose; radiotherapy

OR017

Social Determinants of Cardiovascular Health: A Systematic Review and Meta-Analysis of Psychosocial Support on Treatment Adherence

RENI AFRIANA*, MUHAMMAD AFIIF AZIZ

Faculty of Nursing, Universitas Padjadjaran, Bandung, Jawa Barat 40161, Indonesia

**Correspondence: reniafriana001@gmail.com*

ABSTRACT

Hypertension remains a critical global health challenge, with treatment failure frequently driven by poor medication adherence. Addressing social determinants of health, specifically through psychosocial and community support, is recognised as a vital strategy for advancing community health. This study aims to systematically evaluate the impact of psychosocial support on medication adherence among adult hypertensive patients. Guided by the PRISMA framework, a systematic search was conducted across PubMed, Scopus and EBSCOhost for primary studies published over the last five years. Studies involving adults (≥ 18 years) with primary hypertension that evaluated psychosocial interventions, such as family, peer, or community support, were included. Quality assessment was conducted using the Risk of Bias tool. Statistical synthesis was performed using Review Manager 5.4, utilising the generic inverse variance method and a random-effects model. A total of 20 studies encompassing over 25,000 participants across diverse geographical settings met the inclusion criteria. The meta-analysis demonstrated that patients receiving strong psychosocial support had significantly higher odds of medication adherence compared to those with low or no support (Pooled Odds Ratio = 2.16; 95% CI: 1.34-3.48; $p = 0.002$). Substantial heterogeneity ($I^2 = 92\%$, $p < 0.00001$) was observed, reflecting diverse cultural contexts and intervention types. Psychosocial support acts as a crucial social determinant that significantly enhances medication adherence, doubling the odds of compliance. Integrating family and community-based partnerships into standard cardiovascular care is essential for optimising long-term health outcomes.

Keywords: Hypertension; medication adherence; meta-analysis; psychosocial support; social determinants of health

OR018

Cochlear Implant – Parental Support Scale: A Pilot Study

TANG ZHI LIM¹, CILA UMAT^{1,2*}, PEI-HUA CHEN³, CHUN HONG GAN⁴,
BEE SEE GOH^{5,6}

¹Center for Rehabilitation & Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan
Malaysia, 50300 Kuala Lumpur, Malaysia

²Center for Ear, Hearing & Speech, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala
Lumpur, Malaysia

³Speech and Hearing Science Research Institute, Children's Hearing Foundation, Taipei, Taiwan

⁴Clinical Psychology and Behavioural Health Programme, Faculty of Health Sciences, Universiti Kebangsaan
Malaysia, 50300 Kuala Lumpur, Malaysia

⁵Department of Otorhinolaryngology and Head and Neck Surgery, Faculty of Medicine, Universiti Kebangsaan
Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia

⁶Hospital Canselor Tuanku Muhriz, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia

*Correspondence: cila@ukm.edu.my

ABSTRACT

There is a lack of condition-specific instruments to assess contextual factors among parents of children who are cochlear implant (CI) users or candidates (CIc). Existing measures primarily focus on child outcomes, while parental assessments are typically generic and may not capture the unique challenges of the CI rehabilitation journey. This study aimed to validate a newly developed condition-specific instrument, the Cochlear Implants – Parental Support (CIPS) scale, designed to assist CI team professionals in evaluating parents' psychological well-being [Parental Psychological Status (PPS)] and ecological support systems (ESS). Using purposive sampling, parents of children with CI or CIc completed an online questionnaire comprising the CIPS (PPS and ESS subscales), the Parenting Stress Index–Short Form (PSI-SF), the Family Support Scale (FSS), and the Parenting Stress–Cochlear Implant (PSCI). Internal consistency was examined using Cronbach's alpha, while Pearson's correlations and inferential analyses assessed convergent, divergent, and known-group validity. The CIPS demonstrated excellent internal consistency (PPS: $\alpha = 0.97$; ESS: $\alpha = 0.91$). ESS was insignificantly correlated with PPS ($r = -0.03$, $p > 0.05$), supporting divergent validity. PPS was significantly correlated with PSI-SF ($r = 0.70$, $p < 0.001$) and PSCI ($r = 0.63$, $p < .001$), while ESS was significantly correlated with FSS ($r = 0.73$, $p < 0.001$), supporting convergent validity. For known-group validity, parents with high parenting stress on PSI-SF reported significantly higher PPS scores than those within the normal range (t -value = -2.68 , $p < 0.05$). No significant differences in PPS or ESS were found across demographic or child-related variables ($p > 0.05$). Despite the limited sample size, findings provide preliminary evidence supporting the reliability and validity of the CIPS. Future research should focus on item reduction to shorten the CIPS scale, reduce clinical administrative burden, and enhance its integration into routine CI services.

Keywords: Cochlear implants; parental support; psychological well-being; social support

OR019

Preliminary Histological Evaluation of Oral Pterostilbene in UVB-Induced Skin Photoaging Model in BALB/c Mice

MUHAMMAD HAIKAL BIN AHMAD¹, SITI FATHIAH MASRE¹, DAYANG FREDALINA BASRI², AHMAD ROHI GHAZALI^{1*}

¹Center for Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Center for Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

*Correspondence: rohi@ukm.edu.my

ABSTRACT

Ultraviolet B (UVB) radiation is a major environmental factor contributing to skin photoaging, leading to structural and pigmentary alterations in the skin. Chronic UVB exposure can induce epidermal thickening, collagen degradation, and increased melanin deposition, which are commonly used histological indicators of photoaging. Natural polyphenolic compounds such as pterostilbene have attracted increasing attention due to their potential antioxidant and photoprotective properties. However, the histological effects of orally administered pterostilbene on UVB-induced skin damage remain insufficiently characterised. This study aimed to perform a preliminary histological evaluation of skin changes in UVB-induced BALB/c mice following oral administration of pterostilbene. BALB/c mice were divided into four experimental groups: a normal control group, a UVB-exposed group, and two UVB-exposed groups treated with low-dose and high-dose pterostilbene, respectively. Following treatment, skin samples were collected and prepared for histological examination. Hematoxylin and Eosin (H&E) staining was used to evaluate general skin morphology, including epidermal thickness and structural alterations, while Fontana–Masson staining was performed to assess melanin deposition within the epidermal layer. Preliminary H&E observations revealed marked histological alterations in the UVB-exposed group, including increased epidermal thickness and disruption of normal skin architecture compared with the control group. In contrast, mice receiving oral pterostilbene showed observable differences in epidermal morphology relative to the UVB-exposed group, suggesting a potential protective effect against UVB-induced skin damage. Fontana–Masson staining also indicated variations in melanin distribution among the experimental groups. These preliminary findings suggest that orally administered pterostilbene may modulate histological features associated with UVB-induced skin photoaging. Further studies involving molecular and biochemical analyses are ongoing to elucidate the mechanisms underlying the potential photoprotective effects of pterostilbene.

Keywords: BALB/c mice; histological analysis; photoaging; pterostilbene; UVB

OR020

Metabolite Association and Pathway Analysis of COVID-19 Vaccine via LC-MS-Based Metabolomics

HAMZA AHMED PANTAMI^{1*}, KHADIJA HAMISU²

¹*Department of Pharmaceutical and Medicinal Chemistry, Faculty of Pharmaceutical Sciences, Gombe State University, PMB 127 Gombe Nigeria*

²*Department of Chemical Sciences, Faculty of Science, Gombe State University, PMB 127 Gombe Nigeria*

**Correspondence: hamza3983@gsu.edu.ng*

ABSTRACT

The COVID-19 pandemic marked a turning point in global health, driving unprecedented advances in vaccine development. This study examines the immune responses at molecular level triggered by COVID-19 vaccine using rat models. Eighteen rats were randomly assigned to two groups, with three replicates each: a control group (C) and a vaccinated group (VC). Both groups were maintained under identical conditions for four weeks, except for the vaccination received by VC group, after which blood samples were collected for analysis. Serum biochemical parameters, including globulin (GLB), total protein (TP), albumin (ALB), and aspartate aminotransferase (AST) were evaluated. To further characterise the vaccine, ¹H NMR and LC-MS analysis were conducted to analyse the metabolites present in the vaccine. LC-MS-based metabolomics was employed for the variation between control and vaccinated rats. Multivariate statistical models, including PCA and PLS-DA, effectively discriminated the metabolic profiles of the control and vaccinated groups. Results revealed no significant differences in TP, GLB and ALB levels between the control and vaccinated groups, suggesting no adverse systemic effects post vaccination. However, AST levels were Metabolite Association and significantly elevated in the vaccinated group, indicating localised metabolic activity without systemic toxicity. Variable importance in projection (VIP) scores identified key metabolites driving group separation. Pathway analysis and heatmap visualisation revealed that arachidonic acid, adenosine, stearic acid, oleic acid, and linoleic acid metabolites were upregulated in vaccinated rats, with arachidonic acid, adenosine, and stearic acid showing the most significant increase. These findings suggest these metabolites play critical roles in the vaccine-induced immune response. Conversely, metabolites such as lactate, palmitoleic acid, oleoyl glycine, sphingomyelin, oleamide, and prostaglandin D2 were higher in the control group, indicating downregulation post vaccination. These shifts point to potential metabolic pathways perturbed by the vaccination contributing to a comprehensive understanding of its immune responses.

Keywords: Biochemical test; COVID-19; immune response; metabolomics; VIP values

OR021

Bio-Based *Rhizophora* spp. Particleboard for Tissue-Equivalent Radiotherapy Phantom Applications

SAADIATUL NUR AQILAH MOHD ZAKARIA¹, SITI HAJAR ZUBER^{1*},
MOHD HAFIDZ JAAFAR², MUHAMMAD SAFWAN AHMAD FADZIL¹,
WAN NORDIANA W ABD RAHMAN³, MOHD ZAHRI ABDUL AZIZ⁴,
ABD AZIZ TAJUDDIN⁵

¹Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²School of Industrial Technology, Universiti Sains Malaysia, 11800 Penang, Malaysia

³Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Selangor, Malaysia

⁴Pusat Perubatan USM, Universiti Sains Malaysia, 13200 Pulau Pinang, Malaysia

⁵Albukhary International University, 05200 Kedah, Malaysia

*Correspondence: hajarzuber@ukm.edu.my

ABSTRACT

Three different sizes of *Rhizophora* spp. powder (0 to 104 μm , 105 to 210 μm , and 211 to 500 μm) were used to create particleboards with three different percentages of lignin and sodium alginate (0%, 3% and 6%) as adhesives. The goal of this study is to investigate the physical characteristics, morphology, chemical composition, and effective atomic number, Z_{eff} of *Rhizophora* spp. particleboards that have been incorporated with lignin and sodium alginate. Thickness swelling, water absorption, and the gravimetric method were used to study the physical properties of particleboards. The results showed that the produced particleboards have suitable radiological and physical properties for tissue-equivalent phantoms applications. In contrast, samples with a higher adhesive content and smaller particle sizes displayed less water absorption and thickness swelling. There was good agreement between the density readings and the required requirements. Increased adhesive content reduced the development of voids in the particleboard structure. Higher sodium alginate consumption was correlated with an increase in sodium content, and the calculated effective number further validated the suitability of the material. With a particle range of 0 to 104 μm and 6% adhesive, sample A6 demonstrated significant potential for use in the fabrication of tissue-equivalent phantoms.

Keywords: Alginate-lignin adhesive; *Rhizophora* spp.; particleboard; physical properties; radiotherapy

OR022

Nutrient Density of Commonly Consumed Foods: A Malaysian Perspective

RUHAYA SALLEH^{1,2*}, ROSLEE RAJIKAN², NOOR ANI AHMAD¹, MOHD
RAZIF SHAHRIL², SITI MASITAH ELIAS³, SUZANA SHAHAR²

¹*Institute for Public Health, National Institutes of Health, Ministry of Health, Setia Alam, 40170 Shah Alam, Selangor, Malaysia*

²*Centre for Healthy Ageing and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

³*Faculty of Science and Technology, Universiti Sains Islam Malaysia, Bandar Baru Nilai, 71800 Nilai, Negeri Sembilan, Malaysia*

*Correspondence: ruhaya.s@moh.gov.my and p106546@siswa.ukm.edu.my

ABSTRACT

Food-based dietary guidelines promote the consumption of a variety of nutritious foods to achieve optimal health and prevent chronic diseases. This study aimed to determine the nutrient density of the commonly consumed foods by adults in Malaysia and identify foods that provide the best nutritional value within each food group. A total of 130 food items frequently consumed were included in the analysis. The nutrient-rich food (NRF) score was used to assess the nutrient density. This model considers foods based on ten nutrients to encourage (protein, calcium, iron, potassium, vitamin A, vitamin C, thiamine, riboflavin, niacin, fibre) and three nutrients to limit (total fat, total sugar, and sodium). Nutrient composition data were obtained and standardised per 100 kcal. The NRF score was calculated by summing the percentage of daily recommended values for the encouraging nutrients and subtracting the three nutrients to limit. Foods were then ranked within each food group. Overall, vegetables were the most nutrient-dense group, with ulam raja (60.39), sweet potato shoots (51.55), and bitter melon (49.72) showing the highest scores. Among the fruits, papaya (26.95), star fruit (25.97), and guava (22.16) were the most nutrient-dense. In the legumes group, dhal showed high nutrient density. For milk and milk products, chocolate milk (15.9) had the highest score. Meat and poultry such as chicken liver (38.34) and chicken eggs (14.87) were highly nutrient-dense. Indian mackerel (14.81) and hairtail scud fish (13.9) ranked highest for the fish and seafood group. Wholemeal bread (14.93) and brown rice (11.51) were the most nutrient-dense grains. Fast foods like chicken burger (-7.46) and anchovies (-2.15) were among the least nutrient-dense. Confectionery, processed, and high-sodium foods scored the lowest in the NRF (0.27, -18.87) analysis. This study provides valuable information that can support nutrition policy, dietary guidelines, and public health interventions to promote healthier food choices among Malaysians.

Keywords: Malaysia; nutrient density; nutrient-rich food; nutritional assessment

OR023

Circadian Misalignment and Its Associated Factors in Malaysian Adults: Results from the Nationwide MYLIFERHYTHM Online Survey

**ZAHARA ABDUL MANAF^{1*}, FATIN HANANI MAZRI¹, ARIMI FITRI MAT
LUDIN², MAHADIR AHMAD³, MOHD RIZAL ABDUL MANAF⁴, SAFIAH
TAJUDDIN¹, MARIE-PIERRE ST-ONGE⁵**

¹Centre for Healthy Aging and Wellness and Dietetic Program, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Centre for Healthy Aging and Wellness and Biomedical Science Program Kuala Lumpur, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

³Centre for Community Health Studies and Clinical Psychology Program, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

⁴Department of Community Health, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia

⁵Centre of Excellence for Sleep & Circadian Research and Division of General Medicine, Columbia University Irving Medical Center, New York, NY 10032, United States

**Correspondence: zaharamanaf@ukm.edu.my*

ABSTRACT

Circadian rhythms regulate essential physiological processes over a 24-hour cycle. Disruptions such as circadian misalignment, also known as social jetlag represent a mismatch between biological timing and social obligations, and are associated with increased risks of cardiometabolic disorders and impaired mental health. Despite growing global concern, nationwide data regarding the prevalence of circadian misalignment and its specific associated factors in Malaysia remains limited. This nationwide cross-sectional study of 1,118 non-shift working Malaysian adults (mean age 33.9 ± 8.9 years) aimed to establish population-specific chronotypes, quantify circadian misalignment, and identify their associated factors. Methods: Participants were recruited via a web-based survey aimed at reflecting national population density. Sleep patterns and chronotypes were assessed using a culturally validated Malay version of the Munich Chronotype Questionnaire (MCTQ). Findings revealed an average sleep-debt corrected midpoint of sleep (MSFsc) of 03:24 ± 0:53. Most participants were intermediate chronotypes (68.3%), followed by evening (16.3%) and morning types (15.4%). While a significant proportion (33.9%) experienced circadian misalignment of one hour or more. Furthermore, the mean daily sleep duration was 6.54 ± 1.09 hours, falling below the internationally recommended minimum of 7 hours per night. Multivariate logistic regression identified obesity as the strongest associated factors of circadian misalignment (AOR: 1.85; 95% CI: 1.036–3.289; p = 0.038). Other significant factors included younger age (p = 0.049), being female (p = 0.002), and non-Malay ethnicity (p = 0.005). This study identifies a high prevalence of circadian misalignment

and insufficient sleep among Malaysian adults, with obesity emerging as the strongest associated factor alongside younger age, female gender, and non-Malay ethnicity. These findings underscore an urgent need for targeted public health interventions and flexible social policies to mitigate the associated cardiometabolic and mental health risks associated with disrupted circadian rhythms.

Keywords: Chronotype; circadian misalignment; Malaysian adult; social jetlag

OR024

Measuring How Students Perceive Their Physical Health: Evidence of Validity and Reliability

NIKEN FATIMAH NURHAYATI, DINIE RATRI DESININGRUM*, ANNISA
MAULIDYA

*Faculty of Psychology, Universitas Diponegoro, Jl. Prof. Mr. Sunario, Tembalang, Semarang, Jawa Tengah,
50275, Indonesia*

**Correspondence: diniedesiningrum@live.undip.ac.id*

ABSTRACT

Physical health perception refers to individuals' subjective evaluations of their bodily condition and functional capacity, which play a critical role in health behaviors, psychological well-being, and adjustment during the transition to university life. Despite its importance, validated instruments assessing physical health perception among Indonesian university students remain limited. **Objective:** This study aimed to examine the validity and reliability of the Physical Health Perception Scale (PHP) among first-year university students. A cross-sectional survey was conducted with 1,250 first-year students at a public university in Indonesia. The PHP scale was developed based on Kulsum and Jauhar's framework, comprising three dimensions: evaluation, potential, and activity. Confirmatory factor analysis (CFA) using LISREL 8.8 was employed to assess construct validity. Reliability was evaluated using composite reliability (CR) and average variance extracted (AVE). Six items with factor loadings below 0.50 were removed. The refined model demonstrated satisfactory psychometric properties, with all remaining items showing factor loadings above 0.50. CR values ranged from 0.770 to 0.808, and AVE values ranged from 0.504 to 0.529, indicating adequate convergent validity and internal consistency. **Conclusion:** The PHP scale is a valid and reliable instrument for assessing physical health perception among university freshmen. **Implications:** This scale can support early screening, research on student well-being, and the development of campus-based health promotion and preventive intervention programs.

Keywords: Confirmatory factor analysis; physical health perception; psychometrics; scale validation; university students

OR025

Improving Readiness to Change among Indigenous Risky Drinkers: A Single-Arm Trial of Culturally Adapted Brief Motivational Interviewing

RIFHAN RASULI¹, MARIAM MOHAMAD^{1*}, AIMI NADIRA MAT RUZLIN¹,
ENGKU MARDIAH ENGKU KAMARUDIN²

¹*Department of Public Health Medicine, Faculty of Medicine, Universiti Teknologi MARA, Sungai Buloh Campus, 47000 Sungai Buloh, Selangor, Malaysia*

²*Department of Counsellor Education and Counselling Psychology, Faculty of Educational Studies, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia*

**Correspondence: mariammd@uitm.edu.my*

ABSTRACT

Risky drinking remains a major public health challenge globally, with research indicating its disproportionate impacts on Indigenous communities. This study evaluated the preliminary efficacy of a culturally adapted brief motivational interviewing (MI) intervention – a collaborative, goal-oriented counselling approach designed to evoke motivation for change and resolve ambivalence – in addressing risky drinking among Indigenous adults in Malaysia. This single-arm trial involved 40 participants with Alcohol Use Disorders Identification Test (AUDIT-10) scores ≥ 8 recruited from three Indigenous Jakun settlements in Rompin, Pahang. The intervention comprised three face-to-face sessions delivered over one month, with readiness to change assessed using a validated Jakun-version Readiness to Change Questionnaire at baseline, immediately post-intervention, and at six- and 12-week follow-ups. The cohort was predominantly male (97.5%), aged 30-39 (40.0%), married (60.0%), had completed primary education (45.0%), were self-employed (77.5%), and all belonged to the low-income group. Beer was the main beverage consumed (92.5%). Generalised Linear Mixed Models using a multinomial distribution with a cumulative logit link function revealed a statistically significant main effect of time on the odds of transitioning to a more advanced stage of change ($F(3, 118) = 5.98, p < 0.001$). Pairwise comparisons relative to baseline demonstrated that immediately following the intervention, participants had significantly higher odds of advancing (odds ratio [OR] = 4.05; 95% confidence interval [CI] [1.26, 12.98], $p = 0.019$). This positive trajectory strengthened at six weeks (OR = 9.85; 95% CI [2.52, 38.43], $p = 0.001$) and 12 weeks (OR = 12.94; 95% CI [3.02, 55.54], $p < 0.001$) post-intervention. Sensitivity analysis using the baseline observation carried forward method for missing data confirmed the robustness of these findings ($F(3, 155) = 4.98, p = 0.003$). These results provide preliminary evidence that culturally adapted brief MI is a promising strategy for improving readiness to change among risky drinkers within Indigenous populations in Malaysia.

Keywords: Alcohol drinking; indigenous peoples; motivational interviewing; transtheoretical model

OR026

Development and Validation of a Questionnaire to Assess Nutritional Knowledge Related to Cognitive Function Among Malaysian Older Adults

HANIS MASTURA YAHYA^{1*}, TEO CI WEI¹, PONNUSAMY SUBRAMANIAM²

¹*Nutritional Science Programme, Centre for Healthy Ageing and Wellness, Faculty of Health Science, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Clinical Psychology and Behavioural Health Program, Centre for Healthy Ageing and Wellness, Faculty of Health Science, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

**Correspondence: hanis.yahya@ukm.edu.my*

ABSTRACT

Adequate knowledge of neuroprotective nutrition is essential for preserving cognitive health during ageing. However, there is a lack of validated tools specifically designed to assess nutritional knowledge related to cognition among the Malaysian elderly. This study aimed to develop and validate a reliable questionnaire to fill this gap. The questionnaire was developed by generating an item pool across seven domains, with items scored dichotomously. Content validity was assessed by a panel of six experts, and the Scale-Level Content Validity Index (S-CVI/Ave) was calculated. To evaluate reliability, a two-week interval test-retest was conducted among 108 older adults (mean age 69 ± 7.9 years). Statistical analyses included the Intraclass Correlation Coefficient (ICC), Bland-Altman Limits of Agreement (LOA), and Standard Error of Measurement (SEM%). The initial 30-item draft was refined to 20 items following expert review. The S-CVI/Ave was 0.92, indicating excellent content validity. Reliability analysis demonstrated high stability over time, with ICC values ranging from 0.78 to 0.88. The LOA plots showed strong agreement between assessments, and the SEM% was 4.38% (well below the 10% threshold), confirming the instrument's precision. The developed 20-item questionnaire is a valid and reliable tool for assessing nutritional knowledge related to cognitive function among Malaysian older adults. To enhance the instrument's discriminant power and mitigate the impact of guessing, incorporating a 'Not Sure' response category is recommended for future refinement.

Keywords: Aged; cognition; knowledge; Malaysia; validation study

OR030

Associations between Working Memory Performance and Prefrontal Activation in White Noise among Adolescents with Internet Gaming Addiction

NOOR FADZLIYANA MOHD FADIL^{1,2}, AHMAD NAZLIM YUSOFF^{1*},
MAZLYFARINA MOHAMAD¹, CHING SIN SIAU³, ELZA AZRI
OTHMAN⁴, NOOR ALAUDIN ABDUL WAHAB⁵, FAIZAH MOHD ZAKI⁶,
ZATI AMALINA MOHD ABDUL GHAFAR¹

¹*Centre for Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Medical Imaging Department, School of Integrative Medicine and Life Sciences, KPJ Healthcare University, Kota Seriemas, 71800 Nilai, Negeri Sembilan*

³*Centre for Community Health Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

⁴*School of Medical Imaging, Faculty of Health Sciences, Universiti Sultan Zainal Abidin, 21300 Kuala Nerus, Terengganu, Malaysia*

⁵*Center for Rehabilitation and Special Needs, Faculty of Health Science, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

⁶*Department of Radiology, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia*

*Correspondence: nazlimtrw@ukm.edu.my

ABSTRACT

White noise (WN) can enhance cognitive performance. Previous research demonstrated that WN increased auditory working memory (AWM) performance in healthy young adults during a backward word recall task (BWRT). However, the effect of WN on AWM among adolescents with internet gaming addiction (IGA) have not been studied. This study aimed to investigate the effects of WN at different intensity levels on AWM performance and brain activation among adolescents with and without IGA symptoms. Twenty-one adolescents (9 with IGA symptoms) participated in this study. They performed the BWRT during an fMRI scan in the presence of 45, 50, 55 and 60 dB intensity level WN. The fMRI data were processed using SPM12 to obtain brain activation (NAV) in the prefrontal cortex. Group differences and association between variables of interest were analysed using SPSS. No significant group difference in brain activation is observed at all intensity levels of WN ($p > 0.05$). However, BWRT performance was found to be negatively and significantly correlated with NAV for the right middle frontal gyrus ($\rho = -0.717$, $p = 0.03$) and right superior frontal gyrus ($\rho = -0.763$, $p = 0.017$) but only at 55 dB and in the IGA group. No significant correlation was found in the non-IGA group. Findings from this study suggest that AWM performance in IGA individuals is associated with altered recruitment of prefrontal regions under moderate auditory noise conditions. The negative correlation may reflect enhanced efficiency of neural processing within the dorsolateral prefrontal cortex during a cognitively demanding auditory tasks.

Keywords: Adolescent; brain; dB; gaming; white noise

OR033

Food Insecurity among Adults with Type 2 Diabetes in Malaysia: The Role of Socioeconomic Determinants

SHIVARANJANI MOHAN KUMAR¹, HARVINDER KAUR GILCHARAN SINGH^{1*}, TAN SHY PYNG², NORLAILA MUSTAFA³, SNIGDHA MISRA⁴, KANIMOLLI ARASU⁵, KANISWARI MUNUSAMY⁶

¹*Centre for Community Health Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Centre for Healthy Ageing and Wellness Studies, Faculty of Health Sciences, University Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

³*Department of Medicine, Universiti Malaysia Kebangsaan, 50300 Kuala Lumpur, Malaysia*

⁴*Jeffrey Cheah School of Medicine and Health Sciences, Monash University, 47500 Bandar Sunway, Selangor, Malaysia*

⁵*Division of Nutrition and Dietetics, School of Health Sciences, International Medical University, 57500 Kuala Lumpur*

⁶*Poliklinik Warga, Hospital Canselor Tuanku Mukhriz, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

**Correspondence: harvinder_kaur@ukm.edu.my*

ABSTRACT

Food insecurity is increasingly recognised as a significant social determinant of health that can have a negative impact on dietary habits and disease management in people with Type 2 diabetes mellitus (T2DM). In Malaysia, rising diabetes prevalence, with persistent socioeconomic inequalities, increases the risk of poor diet quality and limited nutrition security. This study examined the association between socioeconomic characteristics, lifestyle factors, diet quality, and food insecurity among adults with T2DM in Klang Valley. This cross-sectional study recruited 100 adults with T2DM from two health clinics. Information on sociodemographic and socioeconomic characteristics, lifestyle practices, and dietary intake was collected through structured interviews and medical record reviews. Food insecurity and diet quality was determined using validated questionnaires. Associations between variables were examined using Chi-square tests and logistic regression analyses. Participants' mean age was 58.5 ± 9.9 years, with females representing the majority (61.0%). More than half of the participants were from the B40 income group (54.0%). Overall, 35.0% of respondents experienced food insecurity. Among the reported indicators, 26.0% expressed concern about food scarcity due to financial limitations, while 17.0% were unable to afford nutritious food. A significant 91.0% of participants had poor or unsatisfactory diet quality. Logistic regression analysis revealed that individuals with lower educational attainment were significantly more likely to experience food insecurity (OR = 6.24; 95% CI:1.98–19.73; $p = 0.002$). Similarly, those who were unemployed (OR = 3.67; 95% CI:1.41–9.53; $p = 0.008$), belonged to the B40 income

group (OR=6.00; 95% CI:2.29–15.76; $p<0.001$), or received financial assistance (OR = 5.34; 95% CI:1.97–14.49; $p = 0.001$) were significantly more likely to experience food insecurity. However, diet quality was not significantly associated with food insecurity ($p = 0.912$). These findings highlight the important influence of socioeconomic vulnerability on food insecurity among adults with T2DM. Addressing broader social determinants of health and strengthening nutrition security initiatives may help improve diabetes management among vulnerable populations.

Keywords: Diabetes mellitus; diet quality; food insecurity; nutrition security; socioeconomic determinants

OR038

From Awareness to Action: Sexual Health Knowledge, Attitudes and Practices among Undergraduates at Universiti Kebangsaan Malaysia

VANITHA MARIAPPAN^{1,2*}, THEENISCHA MAGESWARAN¹, SURENDRAN RAJARATNAM³, ISMARULYUSDA ISYAK^{1,2}, KUMUTHA MALAR VELLASAMY⁴

¹*Nutritional Biomedical Science Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Centre for Research in Psychology and Human Well-Being, Faculty of Social Sciences & Humanities, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia*

³*Center for Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

⁴*Department of Medical Microbiology, Faculty of Medicine, Universiti Malaya, 50603 Kuala Lumpur, Malaysia*

**Correspondence: vanitha.ma@ukm.edu.my*

ABSTRACT

Sexually transmitted diseases (STDs) continue to pose a substantial public health concern among young adults, particularly in developing countries where awareness and preventive behaviours may be inconsistent. University students represent a population at heightened risk due to increased independence, exploration of relationships, and varying levels of sexual health literacy. This study aimed to assess the knowledge, attitudes and practices (KAP) related to STDs among undergraduate students at Universiti Kebangsaan Malaysia and to examine potential differences between science and non-science students. A cross-sectional survey was conducted involving 751 undergraduate students from 12 faculties. Data were collected using a self-administered questionnaire adapted from the validated International AIDS Questionnaire – English Version (IAQ-E) alongside established KAP instruments. Non-parametric statistical analyses were employed to evaluate group differences and explore correlations among knowledge, attitudes, and practices. Overall, 58.6% of respondents demonstrated good knowledge of STDs, 57.5% expressed positive attitudes towards sexual health and prevention, and 61.0% reported engaging in safe practices. Science students achieved significantly higher knowledge scores compared with non-science students ($U = 5277.5$, $Z = -5.822$, $p < 0.001$). However, no statistically significant differences were observed between the two groups in terms of attitudes or practices. Correlation analysis revealed a weak but statistically significant association between knowledge and attitudes ($\rho = 0.082$, $p < 0.05$). No significant relationships were found between knowledge and practices or between attitudes and practices. Despite moderate to high levels of knowledge and generally positive attitudes, the findings indicate that awareness does not necessarily translate into safer sexual behaviours. This disconnect highlights the limitations of knowledge-based approaches alone. Strengthening comprehensive, behaviour-focused sexual health

education, particularly for non-science students, may help bridge the gap between awareness and practice and contribute to more effective STD prevention strategies among university populations.

Keywords: Awareness; knowledge; prevention; sexually transmitted diseases; undergraduates

OR039

Plant Predominant Dietary Patterns and Cognitive Resilience in Breast Cancer Survivors: Translating Nutritional Science into Healthy Longevity

MOHD RAZIF SHAHRIL*, ZHANG JIE, NUR AFIFAH MURSYIDA
ZAUJAN, PONNUSAMY SUBRAMANIAM, SUZANA SHAHAR

Centre for Health Ageing and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia

**Correspondence: razifshahril@ukm.edu.my*

ABSTRACT

Cancer-related cognitive impairment accelerates neurocognitive ageing, posing a significant barrier to healthy longevity in breast cancer survivorship. While fundamental research links this decline to systemic inflammation and oxidative stress, translating these bench insights into bedside clinical applications remains underexplored. Dietary exposure bridges this gap, as plant-predominant diets provide bioactives targeting these pathophysiological pathways. To evaluate nutrition as a translational strategy, this cross-sectional study examined associations between empirically derived dietary patterns and cognitive function among 262 breast cancer survivors at Hospital Canselor Tuanku Muhriz, Malaysia, within the Cancer Survivors Trajectories of Ageing Research (C*STAR) study. Food frequency questionnaires and principal component analysis identified three dietary patterns: vegetable-rich, fruit-and-condiment, and ultra-processed and mixed. A comprehensive neuropsychological battery assessed verbal memory (Rey Auditory Verbal Learning Test), processing speed and attention (Digit Symbol Test; Trail Making Test Part A), working memory (Digit Span), and executive function (Trail Making Test Part B). A composite score captured global cognitive performance alongside domain-specific assessments. Multivariable linear regression models adjusted for sociodemographic and clinical variables revealed no significant associations between dietary patterns and individual cognitive domain scores (all $p > 0.05$). Greater adherence to the fruit and condiment pattern, characterised by higher intake of plant-derived foods and bioactives, was significantly associated with better composite cognitive performance ($\beta = 1.185$, $p = 0.039$). The vegetable-rich, ultra-processed, and mixed patterns showed no significant associations. Adherence to a plant-predominant dietary pattern is associated with better global cognitive performance in breast cancer survivors. Targeted non-pharmacological dietary interventions provide an evidence-based translational strategy to mitigate cognitive ageing and promote healthy longevity after cancer.

Keywords: Breast cancer; cognitive ageing; dietary pattern; survivorship

OR040

Simulation-Based Technologies for Patient Education in Medical Imaging and Radiotherapy: A Systematic Review

NUR ANIS IZZATI CHE MUT^{1,2}, HAIRIL RASHMIZAL ABDUL RAZAK³,
NORHAYATI MOHD ZAIN², RUKIAH A. LATIFF¹, MUHAMMAD
SAFWAN AHMAD FADZIL¹, NORULZAHRAH MOHD ZAINUDIN⁴,
MOHAMAD AMINUDIN SAID⁵, AMALINA FARHI AHMAD FADZLAH⁴,
NUR NAJIAH HAMZAINI¹, NOR ANIZA AZMI^{1*}

¹*Diagnostic Imaging and Radiotherapy Program, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Medical Imaging Department, School of Integrative Medicine and Life Sciences, KPJ Healthcare University, 71800 Nilai, Negeri Sembilan, Malaysia*

³*Faculty of Health and Life Sciences, University of Exeter, Exeter, Devon, EX1 2LU, United Kingdom*

⁴*Fakulti Sains dan Teknologi Pertahanan, Universiti Pertahanan Nasional Malaysia, 57000 Kuala Lumpur Malaysia*

⁵*Nuclear Medicine Department, National Cancer Institute, 62250 Putrajaya, Malaysia*

**Correspondence: noraniza.azmi@ukm.edu.my*

ABSTRACT

Patient education plays a crucial role in preparing individuals for medical imaging and radiation-related procedures. However, conventional education methods such as verbal explanations and printed materials may be insufficient to help patients understand complex procedures and unfamiliar clinical environments. Immersive technologies have been introduced to overcome these limitations. Thus, this review was conducted to explore the current applications of immersive and simulation-based technologies for patient education in medical imaging and radiation procedures, and to assess their impact on patients. The review synthesises findings from studies published between 2005 and February 2026 that were identified through Scopus, PubMed, and WoS databases using keywords related to immersive technology, patient education, medical imaging, and radiotherapy. A total of 20 studies met the inclusion criteria and have moderate to high quality assessments. The technologies used included virtual reality (VR), augmented reality (AR), mixed reality (MR), virtual tours, and video simulation. Based on thematic analysis, four main themes were identified: (i) improvement in patient knowledge and understanding; (ii) reduction of anxiety and psychological distress; (iii) enhancement of patient experience and preparedness; and (iv) usability and feasibility of immersive technologies. Across the reviewed studies, it is reported that immersive technologies improve patient understanding of radiotherapy, imaging procedures, and image-guided interventions. Several studies also demonstrated reductions in anxiety, fear, or distress before treatment or imaging procedures. However, the review found that most studies were concentrated in radiotherapy and radiation oncology settings, with comparatively fewer studies conducted in broader diagnostic imaging contexts including nuclear medicine. This highlights the need for future research to expand

immersive patient education into other modalities. In conclusion, immersive and simulation-based technologies represent effective and feasible strategies for patient education in medical imaging and radiation procedures, with strong potential to improve patient understanding, reduce anxiety, and enhance procedural preparedness.

Keywords: Education; immersive simulation; medical imaging; patient; radiotherapy

OR041

Protecting Vulnerable Workers in a Warming Climate: A Community Health Perspective on Occupational Heat Stress in Malaysia

DAYANA HAZWANI MOHD SUADI NATA*, NUR ADRIANA KHARUDIN,
PUTRI ANIS SYAHIRA MOHAMAD JAMIL

*Centre for Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia,
Kuala Lumpur, Malaysia*

**Correspondence: dayanahazwani@ukm.edu.my*

ABSTRACT

Occupational heat exposure is increasingly recognised as a critical public health and occupational safety concern, particularly in tropical regions where rising ambient temperatures and high humidity intensify heat strain among outdoor workers. Construction workers are among the most vulnerable occupational groups due to sustained physical exertion, prolonged sun exposure, and limited access to preventive heat mitigation measures. Beyond individual health consequences, heat stress may compromise workforce productivity, increase healthcare burden, and exacerbate socioeconomic vulnerabilities within affected communities. This study aimed to determine the prevalence and determinants of heat stress among construction workers in Malaysia and to generate evidence supporting community-oriented occupational health protection strategies. A cross-sectional study was conducted among 87 construction workers from three construction sites in Klang Valley and Johor Bahru using the validated Heat Strain Score Index (HSSI). Descriptive analysis was used to characterise worker demographics and heat-related symptoms, while one-way ANOVA and Pearson correlation analyses were applied to examine differences in heat strain scores and associated risk factors. A substantial proportion of workers reported heat-related illness symptoms, including mild headache (55.2%), muscle pain (54.0%) and dizziness (46.0%). Despite these symptoms, 45% of respondents were categorised within the HSSI Green Zone, indicating potential under-recognition of heat stress risk. Significant differences in HSSI scores were observed across construction sites ($F(2,84) = 15.0, p < 0.001$). Heat strain scores were also significantly associated with body mass index, household income, and hydration practices ($p < 0.001$). These findings underscore the need for strengthened heat stress awareness, improved hydration management, and integrated occupational heat mitigation strategies involving collaboration between researchers, industry stakeholders, and policymakers to enhance worker protection and community health resilience in the context of climate change.

Keywords: Climate change; community resilience; construction workers; occupational heat stress

OR042

Integrated Manual and Automated Image Analysis for Correlation Dielectrophoretic Quantification of Bacteria against Plate Count Enumeration

ARASH ZULKARNAIN AHMAD ROZAINI^{1,2}, MUHAMMAD AKMAL SUHAIMI¹, ABDULLAH ABDULHAMEED³, REVATHY DEIVASIGAMANI¹, NORAZIAH MOHAMAD ZIN², AMIN AHMAD KAYANI^{4,5}, MUHAMAD RAMDZAN BUYONG^{1*}

¹Centre Institute of Microengineering and Nanoelectronics, Universiti Kebangsaan Malaysia, Bangi 43600, Selangor, Malaysia

²Center for Diagnostic, Therapeutic and Investigative Studies Faculty of Health Sciences Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

³Center for Communication Systems and Sensing, King Fahd University of Petroleum & Minerals, Dhahran 31261, Saudi Arabia

⁴Functional Materials and Microsystems Research Group and the Micro Nano Research Facility, RMIT University, Melbourne VIC 3000, Australia

⁵ARC Research Hub for Connected Sensors for Health, RMIT University, Melbourne, Melbourne VIC 3000, Australia

*Correspondence: muhdramdzan@ukm.edu.my

ABSTRACT

Dielectrophoresis (DEP), an electrokinetic technique, offers a rapid alternative to conventional plate count enumeration for bacterial quantification. At an optimal operating frequency, bacterial cells migrate towards regions of high electric field gradients, enabling quantitative observation of particle accumulation within a defined region of interest near microelectrode edges. *Citrobacter freundii* was used as a model organism to evaluate DEP-based bacterial quantification under positive dielectrophoresis (pDEP) conditions. The optimal operating frequency was determined using COMSOL Multiphysics simulations and subsequently validated through experimental DEP observations. Simulations predicted 3 MHz as the optimal frequency, generating an electric field strength of $1 \times 10^5 \text{ V m}^{-1}$ and an estimated particle velocity of $\sim 5 \mu\text{m s}^{-1}$, promoting bacterial accumulation within the region of interest. A constant deposition methodology was implemented, where 1 μL of bacterial suspension was introduced onto the microelectrode surface for DEP analysis, while identical suspensions were plated on agar for conventional colony enumeration. Bacterial concentrations ranging from 10^5 to 10^2 CFU mL^{-1} were investigated to determine DEP-derived bacterial counts per mL. DEP-trapped bacteria were quantified using two image-based approaches. Microscopy images of the defined region of interest (ROI) near the electrode edges were analysed by manual counting and by automated particle detection using ImageJ, where image thresholding and particle analysis were applied to enumerate bacterial cells. Correlation with plate count enumeration was evaluated using linear regression analysis, demonstrating a strong positive relationship.

Automated image analysis exhibited higher correlation ($R^2 = 0.94$, $p < 0.05$) compared with manual counting ($R^2 = 0.86$, $p < 0.05$) across 10^2 – 10^4 CFU mL⁻¹, indicating improved robustness at higher bacterial densities. These findings demonstrate that DEP quantification at 3 MHz reliably reflects bacterial concentration and highlight the potential of DEP-based platforms for rapid quantitative microbial analysis.

Keywords: Automated; bacteria; dielectrophoresis; microbial analysis

OR043

Maternal Genetic Diversity of the Minangkabau Population Revealed by Targeted Mitochondrial Sequencing and Its Forensic Implications

TAUFIK HIDAYAT^{1,2*}, SERI MIRIANTI ISHAR¹, RUS DINA RUS DIN¹,
NUR MAHIA MD ISA³, NOR HAFIDZA HARON⁴, NOOR HAZFALINDA
HAMZAH¹

¹*Forensic Science Program, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Perpustakaan Tun Seri Lanang, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia*

²*Department of Forensic and Medicolegal, Faculty of Medicine, Universitas Andalas, Jalan Perintis Kemerdekaan number 94 Jati, Padang, Sumatera Barat 25127, Indonesia*

³*Jabatan Patologi & Mikrobiologi Veterinar, Fakulti Perubatan Veterinar, Universiti Putra Malaysia, 43400 UPM, Serdang, Malaysia*

⁴*Transfusion Medicine Unit, Pathology Department, Hospital Kajang, 43000 Kajang, Selangor, Malaysia*

*Correspondence: P114808@siswa.ukm.edu.my

ABSTRACT

Multiple human migrations have shaped Island Southeast Asia, but Austronesian expansion is the most significant. Despite this, the maternal genetic structure of many indigenous tribes remains poorly defined at high phylogenetic resolution. In a matrilineal society, the Minangkabau people of Indonesia offer a unique paradigm for studying the dynamic link between social structure and maternal genetic history. Seven primer sets targeting both control and coding regions were used to analyse mtDNA variation in 25 unrelated Minangkabau individuals. These primer sets targeted control (HVS-I and HVS-II) and phylogenetically informative mitochondrial DNA coding regions. After PCR amplification and Sanger sequencing, UGENE, Aliview and SnapGene were used for sequence alignment and polymorphism detection. Haplogroup assignment, diversity indices, genetic distance, and phylogenetic reconstruction were done using MITOMAP, DnaSP6, Arlequin 3.5, and MEGA 12. An analysis found substantial maternal genetic diversity. The research found 25 unique haplotypes in 25 individuals, resulting in a diversity of 1.000 ± 0.0113 . Every sample has a fixed 9-bp 8281-8289del deletion. This deletion distinguishes Asian haplogroups. Y2 (16%) and B4c (B4c1b2a2), both Austronesian lineages, dominated the haplogroup distribution, followed by M7c (12%). The profile also contained trace levels of East Asian (D5b and M9) and West Eurasian (H2a) lineages, indicating a complex history of regional mixing and transoceanic gene flow. These findings demonstrate that the Minangkabau maternal gene pool represents a genetically layered population shaped by migration, admixture, and matrilineal inheritance. Importantly, the high haplotypic diversity and absence of shared haplotypes enhance the forensic utility of mtDNA, particularly for human identification in degraded samples and disaster victim identification in disaster-prone regions.

Keywords: Forensic genetics; haplogroups; island Southeast Asian; Minangkabau; mitochondrial DNA

OR044

Association between Upper Limb Coordination and Grip Strength in Older Adults: A Cross-Sectional Study

WAROONNAPA SRISOPARB^{1,2*}, PAKWIPA JULPHUNTHONG¹, OLAN ISARIYAPAN^{1,2}, NICHAPA NONTACHAICHOOSIN¹, PLAIFA SAISUI¹, PANTITRA THUENPANOM¹

¹*Department of Physical Therapy, Faculty of Allied Health Sciences, Naresuan University, Phitsanulok, 65000 Thailand*

²*Exercise and rehabilitation sciences research unit, Faculty of Allied Health Sciences, Naresuan University, Phitsanulok, 65000 Thailand*

**Correspondence: waroonnapas@nu.ac.th*

ABSTRACT

Age-related declines in upper limb function, particularly coordination and muscle strength, may compromise everyday performance and functional independence over time. Understanding how these domains interact is important for informing strategies that support functional capacity across ageing trajectories. This study aimed to examine upper limb coordination and hand grip strength in older adults, and to explore their relationship. Data were collected from 84 community-dwelling older adults (13 males, 71 females; mean age 69.4 ± 10 years) who demonstrated adequate standing balance, were able to follow instructions, and completed all outcome measures. Upper limb coordination was assessed using the Plate Tapping Test (PTT) and hand grip strength was measured using a Jamar hydraulic hand dynamometer. The relationship between variables was analysed using Spearman's correlation coefficient. The mean PTT completion time for the dominant arm was 14.3 ± 5.42 seconds, and the mean dominant hand grip strength was 24.6 ± 5.61 kg. Males showed better performance than females in both measures. A statistically significant moderate negative correlation was observed between coordination performance and grip strength in the dominant hand ($r_s = -0.422$, $p < 0.001$), indicating that faster coordination was associated with greater strength. Given the moderate strength of the association, these measures may reflect partially distinct aspects of motor performance rather than a single underlying construct. This suggests that age-related changes in upper limb function may involve multiple interacting domains. Considering both coordination and strength in assessment and intervention planning may enhance the specificity of approaches aimed at maintaining upper limb function. Further research is warranted to explore strategies that specifically target upper limb coordination in older adults, alongside conventional strength-focused approaches, with potential implications for sustaining functional independence and supporting healthy ageing trajectories.

Keywords: Aged; arm activity; coordination; elderly; hand grip strength

OR045

Elucidating The Framework of Herbal and Dietary Supplement Consumption among Elderly People

NUR AIN MAHAT^{1*}, MAHANEEM MOHAMED²

¹*Department of Professional Nursing Studies, Kulliyyah of Nursing, International Islamic University Malaysia, 25200 Kuantan, Pahang, Malaysia*

²*Department of Physiology, School of Medical Sciences, Health Campus, Universiti Sains Malaysia, 16150 Kubang Kerian, Kelantan, Malaysia*

**Correspondence: nurainmahat@iium.edu.my*

ABSTRACT

The widespread use of herbal and dietary supplements (HDS) among elderly populations raises concerns regarding safety, appropriateness, and integration with conventional healthcare. Despite their perceived benefits, HDS consumption is often shaped by complex socio-cultural influences and occurs with limited clinical disclosure. This study aimed to (i) explore the practices and expectations of elderly individuals regarding HDS use, and (ii) develop and validate a comprehensive framework to support safe HDS consumption. A qualitative exploratory design was employed, involving in-depth interviews with 12 elderly participants to capture rich experiential data, followed by expert validation of the developed framework. Three overarching themes emerged from the analysis. The first theme, herbal use as a way of life, reflects the routine integration of HDS into daily health practices, driven by familiarity and perceived effectiveness. The second theme, health decisions shaped by heritage and faith, highlights the influence of cultural traditions, intergenerational knowledge, and religious beliefs in shaping health behaviours. The third theme, expectations of staying well in later life, underscores the motivation to maintain independence, prevent illness, and enhance quality of life. Across all themes, limited disclosure of HDS use to healthcare providers was consistently observed. Participants often viewed HDS practices as personal or culturally embedded knowledge rather than clinical information, increasing the risk of potential herb–drug interactions and compromised patient safety. Building on these findings, a conceptual framework was developed and validated by a panel of 8 experts in healthcare, pharmacy, and public health. The framework positions the elderly at the centre, supported by interconnected roles of family members, healthcare providers, and regulatory authorities such as the Ministry of Health and the National Pharmaceutical Regulatory Agency. Family members act as key influencers in decision-making, while healthcare providers are responsible for fostering open communication, promoting disclosure, delivering evidence-based education, and monitoring adverse outcomes. Regulatory bodies contribute through policy dissemination, public education, and enforcement, incorporating public feedback to ensure cultural relevance. Core elements of the framework include reliable advice, shared responsibility, collaborative governance, and effective communication. The expert validation process confirmed the framework’s relevance, clarity, and applicability in real-world settings. This study offers a theoretically grounded and practice-oriented model to enhance safe HDS consumption, strengthen healthcare communication, and inform policy development in ageing populations.

Keywords: Dietary supplement; elderly; herbal

OR046

Neuroprotective Effects of Intermittent Fasting against Inflammatory Markers on Stress Induced Rat's Hippocampus and Prefrontal Cortex

ABUBAKAR HABIBU BUNZA¹, KNS SIRAJUDEEN², NURUL AIMAN MOHD YUSOF¹, NUR ASYILLA CHE JALIL³, MOHD ASNIZAM ASARI^{1*}

¹Department of Anatomy, School of Medical Sciences, Health Campus, Universiti Sains Malaysia, 16150 Kubang Kerian, Kelantan, Malaysia

²Department of Basic Medical Sciences, Kulliyah of Medicine, International Islamic University Malaysia, Bandar Indera Mahkota, 25200 Kuantan, Pahang, Malaysia.

³Department of Pathology, School of Medical Sciences, Health Campus, Universiti Sains Malaysia, 16150 Kubang Kerian, Kelantan, Malaysia

*Correspondence: asnizam@usm.my

ABSTRACT

Intermittent Fasting enhances the antioxidant defence mechanisms in the brain by counteracting oxidative damage. It suppresses the pro-inflammatory nuclear factor kappa B (NF-κB) pathway and increases the activity of nuclear factor erythroid-2-related factor 2 (Nrf2), a master regulator of antioxidant defences, these declined the levels of inflammatory mediators such as TNF-α, IL-1β, and IL-6 and drives the expression of pro-inflammatory cytokines. The study aims to determine the neuroprotective effects of intermittent fasting on the brain's inflammatory markers in rats exposed to stress. This was an animal experimental study conducted in the Health Campus of Universiti Sains Malaysia. A total of 32 male Sprague-Dawley rats were divided into four groups each consisting of eight rats. Fasting groups fast for 1 month followed by 2 weeks of two interrupted stress protocols. The Control-Stress group underwent similar stress protocols. All animals were sacrificed, hippocampus and prefrontal cortex were extracted and homogenised. IL-6, IL-1β, TNF-α and IL-10 ELISA were conducted on the homogenate. Data collected were analysed using GraphPad Prism. There were significant statistical differences of IL-10 in the rat's hippocampus and IL-1β in the Prefrontal cortex but no significant statistical differences of IL-6 and TNF-α in the rat's Prefrontal cortex. Intermittent fasting has been found to improve neuronal survival and synaptic plasticity by reducing the levels of neuro-inflammatory mediators.

Keywords: Chronic stress; inflammatory markers; intermittent fasting; neuroprotection

OR047

A Cross-Sectional Study on the Impact of Eating Behaviors on Dietary Intake in Children with Autism Spectrum Disorder

AINA SHAFIQAH ZARISMAIL, NUR HANA HAMZAID*

Centre of Rehabilitation and Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

**Correspondence: hanahamzaid@ukm.edu.my*

ABSTRACT

Children with autism spectrum disorder (ASD) are commonly known to experience feeding-related problems such as food selectivity due to their sensitive sensory. This would lead to interruptive mealtime behaviors such as aggressive behavior as these children refuse to eat foods that are beyond their sensory needs. If not intervened earlier, these children are at risk of having poor nutritional status later in life. Therefore, this study aimed to identify the common mealtime behaviors exhibited in autism children and investigate the association between mealtime behaviors, autism-related symptoms and nutritional intake among autism children. This cross-sectional study was carried out on 51 children aged three to ten years old who have been diagnosed with ASD at Tunku Ampuan Besar Tuanku Aishah Rohani (HTABTAR). Structured questionnaires were administered to parents for demographic information, mealtime behavior (BAMBI) and challenging behavior in children (ASD-BPC). Anthropometric measurement was obtained to determine nutritional status based on WHO growth standards. WHO Anthro plus version 3.2.2 and SPSS version 29 were used to analyses data. The mean age of participants is 6.4 ± 1.73 years old, with 94% participants showed presence of feeding difficulties. A statistically significant strong association was observed between mealtime behaviors and autism-related symptoms ($r = 0.521$, $p < 0.001$). The acceptance of vegetables was statistically significantly to total calorie intake, $p < 0.05$. However, the ASD-BPC scores and acceptance rate of carbohydrate, protein and fruits do not statistically significantly predict the total calorie intake in children with ASD, $F(5,48) = 1.394$, $p > 0.05$, $R^2 = 0.134$. In conclusion, this study discovered the more severe autism-related symptoms are exhibited, it is more likely that the child would display disruptive mealtime behaviors. However, the acceptance of carbohydrate, protein, fat and autism-related symptoms are not strongly correlated with total energy intake among children with ASD.

Keywords: Autism spectrum disorder; autism-related symptoms; dietary intake; mealtime behavior; nutritional status

PP001

Design and Usability Evaluation of a Mobile Health Application for Childhood Stunting Prevention among Women of Reproductive Age: A Participatory Design Study

RASYIDAH GHAZALLI, WAN AZDIE ABU BAKAR*, NURUL HAZIRAH JAAFAR

Kulliyah of Allied Health Sciences, International Islamic University Malaysia, 25200 Kuantan, Pahang, Malaysia

**Correspondence: wazdie@iiu.edu.my*

ABSTRACT

Mobile phones have emerged as powerful tools for delivering health information to the public. However, limited locally developed mHealth interventions targeting stunting prevention among women of reproductive age exist in Malaysia. Therefore, the study aimed to develop and evaluate the usability of a mobile health application, Be Healthy, Let's Stop Stunting (BLESS), designed to deliver nutrition education targeting a triad of prospective groups of women of reproductive age (pre-conception women, pregnant women, and mothers). The study was carried out in iterative stages following the five-stage ADDIE instructional design model (Analysis, Design, Development, Implementation, and Evaluation) and employed purposive sampling for participant recruitment. First, a user needs assessment was conducted through semi-structured interviews and focus group discussions to gather opinions from women of reproductive age about the preferred design of the mHealth app. In the next stage, a prototype of the application was created and developed. The final stage involved user usability evaluation using a Malay version of the Mobile Application Usability Questionnaire. The needs assessments indicated that the preferred application design should prioritise user-friendliness, familiarity with the interface, and offline functionality. Thirty-seven women of reproductive age participated in the usability testing. The participants expressed high satisfaction with the application (mean score: 6.16 ± 0.47 out of 7.00). The BLESS mHealth application demonstrated high usability and the potential to improve nutritional knowledge among women of reproductive age and to serve as a preventive measure against stunting among children below 2 years old in Malaysia.

Keywords: ADDIE; childhood stunting; mHealth; nutrition education; women of reproductive age

PP002

Antibacterial Activity and Physicochemical Characterisation of *Centella asiatica* Ethanolic Extract Cream Formulations against Selected Pathogenic Bacteria

SITI KHADIJAH PADZIL*, AUF ALI, NURUL AINAA NAJLA YUSRI,
NOR IZZA KAMILIA NORSIDEK, ATHIRSHDA

KPJ Healthcare University, 71800 Nilai, Negeri Sembilan, Malaysia

*Correspondence: ucn.khadijah@kpju.edu.my

ABSTRACT

Centella asiatica (pegaga) is a traditional medicinal plant well-recognised for its wound healing and antimicrobial properties. Despite growing interest in herbal topical formulations, scientific evidence supporting standardised *C. asiatica* cream preparations remains limited. This study aimed to formulate, physicochemically characterise and evaluate the antibacterial activity of *C. asiatica* ethanolic extract cream formulations at 2.5% and 7.5% concentrations against *Staphylococcus aureus*, *Escherichia coli* and *Bacillus cereus*. Ethanolic extracts were prepared via cold maceration using 90% ethanol. Oil-in-water cream formulations were developed and subjected to physicochemical evaluation, including pH, washability, dye test, and organoleptic assessment over 30 days. Fourier Transform Infrared (FTIR) spectroscopy was employed to confirm the presence and stability of key bioactive compounds. Antibacterial activity was assessed using well diffusion, minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) assays, with gentamicin (0.1% w/w) as positive control. Both formulations demonstrated skin-compatible pH (4.79–5.36), stable oil-in-water characteristics, and acceptable physical properties. FTIR analysis confirmed the preservation of functional groups associated with triterpenoids and glycosides. The 7.5% formulation exhibited significantly greater antibacterial activity ($p < 0.05$), producing inhibition zones of 16.5–22.5 mm, with an MIC of 50 mg/mL against all tested organisms. No bactericidal effect was observed, indicating a bacteriostatic mode of action. In conclusion, *C. asiatica* ethanolic cream, particularly at 7.5% concentration, demonstrates promising potential as a stable topical antibacterial formulation compared to lower % concentration. These results indicate a concentration-dependent antibacterial effect. Further optimisation and clinical evaluation are recommended to enhance its therapeutic applicability.

Keywords: Antibacterial; *Centella asiatica*; cream formulation; FTIR; phytochemical screening

PP005

Phytochemical Screening and Antimicrobial Activity of *Catharanthus roseus*: Comparative Efficacy of Leaf, Flower and Root Extracts against *Escherichia coli* and *Candida albicans*

SITI KHADIJAH PADZIL*, AUF ALI, NIROSHA PALARAMAN, SANGEETA THAMORTHIRAN

School of Pharmacy, KPJ Healthcare University, 71800 Nilai, Negeri Sembilan, Malaysia

*Correspondence: ucn.khadijah@kpju.edu.my

ABSTRACT

Catharanthus roseus (Kemunting Cina) is well-established as the source of the anticancer. However, its antimicrobial potential across different plant parts remains underexplored. With antimicrobial resistance (AMR) driving demand for novel plant-derived therapeutics, this study aimed to evaluate and compare the phytochemical profiles and antimicrobial efficacy of ethanolic extracts from the leaves, flowers, and roots of *C. roseus* against *Escherichia coli* and *Candida albicans*. Plant materials were extracted by cold maceration using 95% ethanol for seven days. Phytochemical screening employed standard qualitative methods and antimicrobial activity was assessed by the agar well diffusion method at 200 mg/mL, with Amoxicillin and Griseofulvin as positive controls. Statistical significance was determined using the unpaired t-test. Extraction yields were highest in leaves (16.25%), followed by flowers (12.29%) and roots (5.60%). Alkaloids, carbohydrates, flavonoids, proteins, and terpenoids were present in all plant parts, while glycosides and saponins were absent in flowers. All extracts showed statistically significant antimicrobial activity ($p < 0.001$). The leaf extract exhibited the strongest antibacterial activity against *E. coli* (12.9 mm), followed by flowers (9.0 mm) and roots (6.1 mm). For antifungal activity against *C. albicans*, roots produced the largest inhibition zone (6.8 mm), followed by leaves (6.0 mm) and flowers (4.8 mm). These findings demonstrate plant part-specific antimicrobial potential: leaves showed superior antibacterial efficacy consistent with their complete secondary metabolite profile, while roots displayed the strongest antifungal activity, likely driven by saponins known to disrupt fungal membranes. The reduced bioactivity of flower extracts correlates with the absence of these key phytochemicals. Collectively, these results extend the therapeutic relevance of *C. roseus* beyond oncology and support further isolation and characterisation of its bioactive compounds to address the growing AMR crisis.

Keywords: Antifungal activity; antimicrobial activity; *Catharanthus roseus*; ethanolic extract; phytochemical screening

PP006

Beyond the Lead Apron: Modeling Lifetime Cancer Risk Estimates from Thyroid Exposure in Clinical Staff

NOR AIN RABAIEE^{1*}, NUR FASEHAH HANI SHARANI², SAYED INAYATULLAH SHAH², NOR AZIYATUL IZNI³, MUSTAFA SALIH AL MUSAWI⁴

¹Department of Radiology, Kulliyah of Medicine, International Islamic University Malaysia, 25200 Kuantan, Pahang, Malaysia

²Department of Diagnostic Imaging and Radiotherapy, Kulliyah of Allied Health Sciences, International Islamic University Malaysia, 25200 Kuantan, Pahang, Malaysia

³Centre of Foundation Studies, Universiti Teknologi MARA, Cawangan Selangor, 43800 Dengkil, Selangor, Malaysia

⁴Faculty of Medical Sciences, Jabir Ibn Hayaan University, Najaf 54001, Najaf Governorate, Iraq

*Correspondence: norainrabaiee@iiu.edu.my

ABSTRACT

In radiology, radiation protection often prioritises the torso via lead aprons, frequently neglecting the highly radiosensitive thyroid gland. A significant research gap exists in translating physical dose measurements into long-term biological risks, particularly concerning gender-specific sensitivities. Current safety protocols often rely on generalised models that may underestimate the risk for female personnel. This study aimed to quantify the lifetime attributable risk (LAR) of thyroid cancer among clinical staff during central venoplasty procedures to advocate for enhanced shielding protocols. Thyroid doses were measured using nanoDot dosimeters during a single central venoplasty procedure for four clinical roles: Radiologist, Medical Officer (MO), Radiographer, and Staff Nurse. Risk estimation was performed using the NCI Radiation Risk Assessment Tool (RadRAT v.4.3.1), utilising an acute exposure model based on a U.S. 2015–2019 population baseline. Lifetime risk was calculated by integrating gender, birth year, and specific organ dose (mGy). The quantitative modeling revealed a significant disparity between radiation dose and subsequent excess lifetime risk (ELR). The radiologist received the highest dose of 0.064 mGy, resulting in a mean ELR of 0.0101 per 100,000. In comparison, the MO and radiographer received lower doses of 0.033 mGy and 0.021 mGy, with mean ELR values of 0.00634 and 0.0059 per 100,000, respectively. Notably, a biological outlier was observed in the staff nurse despite receiving the lowest dose of 0.007 mGy. The mean ELR for staff nurses was 0.00935 per 100,000, nearly identical to the radiologist's risk. The findings reveal a profound gender-based disparity where female clinical staff face disproportionately higher ELR and baseline future risks, even when receiving up to nine times less radiation dose than male colleagues. Acknowledging the heightened radiosensitivity of the female thyroid is essential for healthcare institutions to align with ALARA principles and proactively mitigate the long-term cancer burden among interventional personnel.

Keywords: Excess lifetime risk; interventional radiology; occupational health; radrat; thyroid dosimetry

PP008

Human Health Risk Assessment Associated with Exposure to Per- and Polyfluoroalkyl Substances in Fine Particulate Matter (PM_{2.5})

NURALIA SYAHIDA MOHD SHARIFA¹, 'AISAMUDDIN FAIZ MOHAMAD NOR¹, MASITAH MOHAMAD¹, DIDI ERWANDI MOHAMAD HARON², MUHAMMAD IKRAM A WAHAB^{1*}

¹Centre for Toxicology & Health Risk Studies, Environmental Health & Industrial Safety Program, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²SUTRALAB, Centre of Research Facility, Department of Research Development, Deputy Vice Chancellor (Research & Innovation), Universiti Malaya, 50603 Kuala Lumpur, Malaysia

*Correspondence: ikram@ukm.edu.my

ABSTRACT

Fine particulate matter (PM_{2.5}) is a significant urban air pollutant capable of adversely affecting human health through inhalation exposure; however, studies on its role as a carrier for Per- and Polyfluoroalkyl Substances (PFAS) in Malaysia remain very limited. This lack of data creates a critical knowledge gap, particularly in assessing health risks associated with airborne PFAS exposure. Therefore, this study aimed to determine the concentration and distribution of PFAS bound to PM_{2.5} in the urban environment of Kuala Lumpur, to evaluate seasonal variations between the Northeast Monsoon and Southwest Monsoon, and to assess non-carcinogenic and carcinogenic health risks arising from inhalation exposure. PM_{2.5} sampling was conducted at a high-traffic urban location near Hospital Kuala Lumpur over a full annual monsoon cycle. PFAS compounds associated with PM_{2.5} were analysed using liquid chromatography–mass spectrometry, while seasonal differences were evaluated using the Mann–Whitney U test. Human health risk assessment was performed based on Lifetime Average Daily Dose (LADD), Hazard Quotient (HQ), and Cancer Risk (CR), supported by the Multiple-Path Particle Dosimetry (MPPD) model to evaluate PM_{2.5} deposition patterns within the human respiratory system across different age groups. The results demonstrated the consistent presence of PFAS bound to PM_{2.5} in ambient air, with dominance of short-chain compounds such as PFBA alongside detectable levels of legacy PFAS including PFOA, and compound-specific seasonal variability. Health risk assessment indicated that children are more vulnerable to non-carcinogenic risks, while carcinogenic risk associated with PFOA increased with age and cumulative exposure duration. This study contributes new scientific evidence on airborne PFAS in a tropical urban setting and provides essential baseline data to support the inclusion of PFAS in ambient air monitoring programmes and the strengthening of air pollution management policies and public health protection strategies in Malaysia.

Keywords: Fine particulate matter; polyfluoroalkyl substances

PP009

Exciting and Inhibiting Connections in Healthy Adult Brain: Applying Bayesian Inference on Resting-State fMRI Data Stratified by Brain Hemisphere, Gender and Handedness

SITI AISHAH SHUIB¹, AHMAD NAZLIM YUSOFF^{1*}, NOOR ALAUDIN ABDUL WAHAB², FAIZAH MOHD ZAKI³, ELZA AZRI OTHMAN⁴, MOHD IZUAN IBRAHIM¹, NURUL HANIM NASARUDDIN⁵

¹Center for Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Science, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Center for Rehabilitation and Special Needs Studies, Faculty of Health Science, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

³Department of Radiology, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia

⁴School of Medical Imaging, Faculty of Health Sciences, Universiti Sultan Zainal Abidin, Gong Badak Campus, 21300 Kuala Nerus, Terengganu, Malaysia

⁵Faculty of Cognitive Sciences and Human Development, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia

*Correspondence: nazlimtrw@ukm.edu.my

ABSTRACT

The human brain maintains a dynamic equilibrium between excitatory and inhibitory neuronal interactions. While a balance in the number of excitatory and inhibitory neurons is not strictly required for neural equilibrium, it remains unclear whether a similar principle applies to the distribution of exciting and inhibiting connections. Addressing this gap is clinically significant, as disproportionate excitatory–inhibitory connectivity has been increasingly implicated in the pathophysiology of neuropsychiatric and neurological disorders. This study investigated the distribution of exciting and inhibiting connections within the Default Mode Network during rest using resting-state functional magnetic resonance imaging (fMRI) data from 47 healthy young adults, with the primary objective of establishing a normative baseline against which disease-related connectivity imbalances can be identified. Dynamic Causal Modeling and Statistical Parametric Mapping (SPM) were employed to analyse effective connectivity among key DMN nodes, including the posterior cingulate cortex, medial prefrontal cortex, bilateral inferior temporal cortex, lateral parietal cortex, medial dorsal thalamus, and bilateral posterior cerebellum. A fully connected model was specified and inverted for each participant, with Bayesian Model Averaging applied at the group level. Results showed that all connections among DMN nodes exceeded a posterior probability of 0.9, indicating strong bidirectional interactions between regions, with inhibiting connections outnumbering exciting connections during rest. Analyses stratified by brain hemisphere, gender, and handedness further revealed apparent differences in the relative strength of excitatory-inhibitory connectivity, indicating

that these biological factors meaningfully shaped intrinsic connectivity patterns. Together, these findings provide a clearer understanding of DMN dynamics in healthy individuals and establish a standardised reference framework for detecting connectivity imbalances associated with disorders such as depression, schizophrenia, autism spectrum disorder, and Alzheimer's disease

Keywords: Adolescent; Bayesian; default mode network; exceedance; probability

PP011

Phytochemical Profiling and GC-MS Analysis of *Eleusine Indica* L. Gaertn Polar Extracts: Unveiling Novel Bioactive agents for Multiphase Wound Healing

MAZLYZAM ABDUL LATIF^{1,2*}, NUR SHAKILA GEAIN², ASMAH HAMID^{1,2}, FATINI NADIAH MOHAMMAD ZAMZAMI¹

¹Center for Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Biomedical Science Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia.

*Correspondence: mazlyzam@ukm.edu.my

ABSTRACT

Chronic wounds pose a major clinical challenge, underscoring the need for affordable, innovative wound-healing agents. *Eleusine indica* L. Gaertn, a common grassy weed used in local wound care, has been shown to have antimicrobial, anti-inflammatory, and antioxidant properties that support tissue repair. This study aimed to perform a qualitative phytochemical screening and quantitative gas chromatography-mass spectrometry (GC-MS) profiling of different polar extracts from *E. indica* leaves. The goal was to isolate and identify the specific bioactive components responsible for its healing effects. We extracted *E. indica* leaves using three polar solvents: methanol and ethanol through cold maceration, and water through both hot and cold infusion methods. The extracts were qualitatively screened for saponins, tannins, alkaloids, flavonoids, and triterpenoids. Afterward, GC-MS analysis was conducted to separate and identify their volatile bioactive constituents. Qualitative screening showed that alkaloids and flavonoids were present in all tested polar extracts. Saponins were only found in the aqueous extracts, while triterpenoids were exclusively isolated from the cold aqueous infusion. The GC-MS analysis identified a variety of bioactive compounds linked to faster wound healing. Notably, the methanol and ethanol extracts contained high levels of essential fatty acids, particularly 9,12,15-octadecatrienoic acid (up to 17.49%) and tetradecanoic acid (up to 15.02%), recognised for their strong anti-inflammatory and antimicrobial properties. Significant diterpenes, such as phytol, were identified, along with phytosterols like stigmasterol and gamma-sitosterol, which support antioxidant mechanisms and cell growth during tissue repair. The wide range of identified anti-inflammatory, antimicrobial, and antioxidant metabolites provides chemical support for the traditional use of *E. indica*. These findings underscore its strong potential for future clinical applications as new, commercially viable therapeutic agents targeting various stages of the wound-healing process.

Keywords: Bioactive compounds; *Eleusine indica*; gas chromatography-mass spectrometry; polar extracts; wound healing

PP012

High Bacterial Contamination Burden and MDRO Prevalence on Hospital Fomites: A Quantitative Study from a Tertiary Referral Hospital in Klang Valley Malaysia

JULIZA JUMARI^{1,3}, NORAZIAH MOHAMAD ZIN^{1*}, SUBASH NAMBIAR
SREE PATHI², SITI SHAHARA ZULFAKAR¹, NAURAH MAT ISA²,
NUR FAIZAH ABU BAKAR¹, SHIRLEY TANG GEE HOON¹, MOHD
YUSOF HAMZAH², CHEE LOON LEONG⁴

¹Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Radiation Processing Division, Malaysian Nuclear Agency, 43000 Kajang, Selangor, Malaysia

³Faculty of Health Sciences, MAIWP International University, 68100 Kuala Lumpur, Malaysia

⁴Infectious Diseases Unit, Department of Medicine, Hospital Kuala Lumpur, 50586 Kuala Lumpur, Malaysia

*Correspondence: noraziah.zin@ukm.edu.my

ABSTRACT

Healthcare-associated infections (HAIs) are serious threat to patient safety, yet the role of medical fomites as environmental reservoirs of HAIs, especially related to antimicrobial resistance (AMR) is often under-reported. This study investigated the prevalence of bacterial and AMR profiles on medical fomites across the medical ward in a tertiary referral hospital in Klang Valley, Malaysia. A quantitative study of cross-sectional environmental surveillance was conducted nine times between October 2024 and February 2025. 648 samples were collected randomly from five high-touch medical fomites namely blood pressure (BP) cuff, thermometer, stethoscope, glucometer and telephone across the tuberculosis ward, male general medical ward, high dependency ward, isolation ward, coronary care unit and cardiac rehabilitation ward. Bacterial contamination load on each fomite was determined using the standard plate count method and expressed as colony-forming units per item (CFU/item). Meanwhile, bacterial identification was performed by MALDI-TOF MS, and antimicrobial susceptibility testing was determined using VITEK® 2 analysis on selected isolates. Of whole samples, 98.1% showed significant microbial contamination. Statistical fomite analysis revealed that BP cuff had the highest bacterial contamination mean $\log_{10}$ (CFU/Items) ($M = 3.81$, $SD = 1.10$) than Stethoscope ($M = 3.18$, $SD = 0.81$), glucometer ($M = 2.58$, $SD = 0.66$), thermometer ($M = 2.52$, $SD = 1.02$), and telephone ($M = 2.58$, $SD = 0.91$), all $p < 0.05$. Bacterial contamination in the ward showed significant variation, $H(5) = 14.78$, $p < 0.05$ ($N = 648$). High dependency ward demonstrated highest contamination ($M = 119,300$, $SD = 789,133$, median = 1,500), while CRW had the lowest ($M = 6,205$, $SD = 24,043$, median = 450). Among 144 selected isolates, 20.8% ($n = 30$) were identified as AMR strains versus 79.2% as susceptible strains. The AMR strains were predominantly present in the high-dependency ward namely Multidrug-Resistant *Acinetobacter baumannii*, Methicillin-Resistant *Staphylococcus aureus* (MRSA), Extended-spectrum beta-lactamase *Klebsiella pneumoniae*, *Enterobacter cloacae* and *Pseudomonas sp.* Medical fomites in this hospital serve as critical reservoirs of bacterial contamination and a significant multidrug-resistant organism

burden, posing a high risk for cross-transmission. There is a vital need for enhanced stewardship and the integration of automated disinfection technologies to mitigate the HAIs, especially of AMR transmission in Malaysian hospitals.

Keywords: Antimicrobial resistance; fomite contamination; healthcare-associated infections; multidrug-resistant organisms

PP013

Genotoxicity Assessment of Buccal Cells among Municipal Workers Occupationally Exposed to Fogging Spray

NURUL FARAHANA KAMALUDIN^{1,2*}, QISTINA NAJIHA AHMAD
FAIZAL¹, NORMAH AWANG^{1,2}, SIEW EE LING³

¹Environmental Health & Industrial Safety Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

¹Center for Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

³ASASI Pintar Program, Pusat PERMATA@Pintar Negara, Universiti Kebangsaan Malaysia, 43600 Bangi, Malaysia

*Correspondence: nurulfarahana@ukm.edu.my

ABSTRACT

Pyrethroids, the primary class of insecticides used in mosquito fogging activities, are generally regarded as safer alternatives to older pesticides such as organophosphates. However, increasing evidence suggests that pyrethroid exposure may induce oxidative stress and DNA damage, raising concerns about potential health risks among workers involved in fogging operations. This study aimed to assess genotoxic damage in buccal cells among municipal workers exposed to insecticides during fogging activities. This cross-sectional study involved 30 respondents comprising 15 fogging operation workers (exposed group) and 15 office workers (control group) from a municipal council. A structured questionnaire was used to obtain information on sociodemographic characteristics, smoking status, and duration of occupational exposure to insecticides. Genotoxic damage was evaluated by determining the frequency of micronuclei (MN) per 1,000 buccal cells following acridine orange staining and fluorescence microscopy analysis. The results showed that MN frequency was significantly higher among workers in the exposed group compared to the control group ($p < 0.001$), indicating increased genotoxic damage associated with occupational exposure to fogging insecticides. Smoking status, age, and duration of exposure were also examined as potential contributing factors. A significant difference in MN frequency was observed between smokers and non-smokers within the exposed group ($p < 0.05$), while age did not show a significant association with MN frequency. Duration of exposure showed a weak but significant positive correlation with MN frequency ($R^2 = 0.072$, $p < 0.05$). Multiple linear regression analysis revealed that smoking status and duration of insecticide exposure were significant predictors of MN frequency, collectively explaining 65% of the variation observed ($R^2 = 0.65$, $p < 0.01$). These findings suggest that workers involved in fogging operations may face an increased genotoxic risk, highlighting the importance of continuous health monitoring and preventive occupational safety measures.

Keywords: Buccal cells; fogging; genotoxicity; insecticides; occupational exposure

PP014

Challenges in Accessing Hearing Healthcare Services among the Parents of Children with Cochlear Implants during the COVID-19 Pandemic

TANG ZHI LIM¹, CILA UMAT^{1,2*}, CHUN HONG GAN³, PEI-HUA CHEN⁴,
BEE SEE GOH^{5,6}

¹Center for Rehabilitation & Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Center for Ear, Hearing & Speech, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

³Clinical Psychology and Behavioural Health Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

⁴Speech and Hearing Science Research Institute, Children's Hearing Foundation, Taipei City 114, Taiwan

⁵Department of Otorhinolaryngology and Head and Neck Surgery, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia

⁶Hospital Canselor Tuanku Muhriz, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia

*Correspondence: cila@ukm.edu.my

ABSTRACT

The COVID-19 pandemic, a global event that led to a lockdown lasting around three years, has significantly impacted sectors such as the economy, education, and healthcare. This study aims to investigate the challenges parents face in accessing hearing healthcare services for children who received cochlear implants (CI) before and during the pandemic. Sixteen parents were interviewed to understand their experiences accessing hearing healthcare services during the COVID-19 pandemic. Children who received CI before and during the pandemic were categorised into pre-COVID-19 (n = 8) and COVID-19 (n = 9) groups, respectively. Three main challenges were identified among the parents. Firstly, difficulty in attending follow-up appointments was common for both groups of parents. Secondly, challenges related to performing online rehabilitation sessions at home were reported by parents from the pre-COVID-19 group, while time pressure for CI surgery was a stressor for parents in the COVID-19 group. However, interestingly, when comparing the timing of deafness diagnosis and CI-related variables of CI children between both groups, no significant differences were found ($p > 0.05$), indicating that the procedure was not affected by the pandemic. Although the parents reported numerous challenges in accessing hearing healthcare services, the comparative analysis revealed no significant differences in age at deafness diagnosis or age at CI implantation. This could be attributed to a standardised, rigorous protocol implemented at the tertiary hospital. In conclusion, while parents reported numerous challenges during the pandemic, the effects were not significantly different across this cohort, as they were alleviated by promptly receiving an appropriate telepractice service plan and by strictly following the established instructions. Nevertheless, the

pandemic has facilitated the rapid growth of telepractice services, enabling more families to access hearing healthcare in diverse ways.

Keywords: Children with cochlear implants; COVID-19; hearing healthcare services; parental challenges

PP015

Multiligamentous Knee Reconstruction All-inside Technique with Modified Universiti Kebangsaan Malaysia Internal Bracing Graft

**KESAVAN RAVINDRAN^{1*}, BADRUL AKMAL HISHAM MD YUSOFF²,
MUHAMMAD KARBELA REZA RAMLAN², NIK KAMARUL ARIF², ALIFF
OMAR², MOHAMAD AZWAN AZIZ²**

¹Department of Orthopaedic, Hospital Canselor Tuanku Muhriz, 56000 Cheras, Kuala Lumpur, Malaysia

²Arthroscopic and Sports Unit, Department of Orthopaedic, Hospital Canselor Tuanku Muhriz, 56000 Cheras, Kuala Lumpur, Malaysia

**Correspondence: kesharavz@gmail.com*

ABSTRACT

Multiligament knee injuries (MLKI) remain surgically challenging, with high risks of arthrofibrosis, graft failure, and tunnel-related complications. This technical note describes a single-stage, all-inside anterior cruciate ligament (ACL) and posterior cruciate ligament (PCL) reconstruction augmented with Modified Universiti Kebangsaan Malaysia Internal Bracing (MUIB) graft construct, combined with a Modified Larson posterolateral corner (PLC) reconstruction, aiming to enhance biomechanical stability while reducing tunnel widening and facilitating accelerated rehabilitation. All-inside PCL and ACL reconstruction was performed arthroscopically using retrograde socket preparation with FlipCutter technology. The MUIB graft technique incorporated FiberTape centrally within the allograft rather than externally augmenting the graft, thereby minimising windshield-wiper and bungee effects associated with tunnel widening. The PCL graft (approximately 80 mm) and ACL graft (within 62-67 mm depending on patient height) were secured using suspensory fixation devices with sequential tensioning. A Modified Larson PLC reconstruction was performed through an open approach using a single allograft to reconstruct the lateral collateral ligament (LCL) and popliteus tendon via anatomically positioned fibular and femoral tunnels. Graft fixation followed a defined sequence: PCL tensioned at 90° flexion with anterior drawer force, ACL in full extension, followed by simultaneous PLC tensioning. An accelerated rehabilitation protocol emphasising early range of motion, protected weight bearing, and early quadriceps activation was implemented. Internal bracing has demonstrated promising outcomes in MLKI reconstruction, with reported failure rates between 0% and 13.6%. However, conventional augmentation techniques may predispose to tunnel widening. By embedding the FiberTape within the graft substance, the MUIB construct aims to provide load sharing while mitigating tunnel-related complications. This approach offers anatomic restoration, reduced graft attenuation during the “killer turn” in PCL reconstruction, and potential for earlier functional recovery. Single-stage all-inside ACL-PCL reconstruction augmented with the MUIB graft and combined with modified Larson PLC reconstruction provides a biomechanically robust construct that may benefit in multiligament knee injuries

Keywords: All-inside reconstruction; internal bracing; multiligament knee injury; posterior cruciate ligament reconstruction; posterolateral corner reconstruction

PP016

Analytical Strategies for the Forensic Characterisation of E-Liquids: Matrix Mitigation and Sensitive Detection of Alkaloids and Additives

IFA SUFIEQAH MOHD ZULFIKAR¹, NUR SYAZWANIS NABILA CHE
MANSHOR¹, ROZAIMY ABDUL RAHIM², NOOR HAZFALINDA
HAMZAH², ATIKAH MOHD NASIR^{2*}

¹*Forensic Science Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Bangi, Selangor, Malaysia*

²*Centre of Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

**Correspondence: atikahnasir@ukm.edu.my*

ABSTRACT

This narrative review examines the development of analytical approaches used for the forensic characterisation of electronic cigarette liquids, with particular attention to matrix-related challenges and the detection of minor or undeclared constituents. Early analyses focused largely on nicotine quantification, whereas more recent work has expanded towards broader chemical profiling of alkaloids, flavouring agents, sweeteners, and synthetic additives. A major analytical difficulty in this field is the complex e-liquid matrix, in which high concentrations of propylene glycol and glycerol can suppress or distort the detection of target compounds. The review therefore considers sample-preparation strategies designed to reduce matrix interference, including freeze-out extraction, liquid–liquid extraction and solid-phase microextraction. It also evaluates the use of modern separation and detection platforms, particularly GC–MS/MS, LC–MS/MS and HILIC–MS/MS, for the targeted analysis of nicotine-related alkaloids, saccharides such as sucrose, and other additives of forensic or regulatory interest. In addition, the review discusses sensitivity-enhancement approaches in capillary electrophoresis, including field-amplified sample injection, which has enabled markedly improved detection of trace alkaloids such as pseudoephedrine and ephedrine. Taken together, these developments show how analytical chemistry has progressed from routine nicotine measurement to more refined forensic profiling of e-liquids. The review highlights the importance of method selection according to matrix complexity, analyte class and evidential purpose, and shows how robust analytical workflows can support product authentication, detection of undeclared substances, and more reliable regulatory assessment of electronic cigarette formulations.

Keywords: Alkaloids; analytical chemistry; electronic cigarette liquids; forensic characterisation; matrix effect

PP017

RAPD-PCR Profiling for Comparative Study of Low-Quantity Human DNA (Hair Shaft) and *E. coli* Banding Patterns

AINATUL IMANI, SERI MIRIANTI ISHAR*

Forensic Science Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

*Correspondence: seri.ishar@ukm.edu.my

ABSTRACT

Forensic analysis of human hair shafts remains one of the biggest challenges, as DNA is often limited and degraded and is frequently susceptible to microbial contamination. Traditional DNA typing is often inapplicable for such samples. For contaminated evidence, the RAPD approach offers a flexible and sequence-independent alternative for creating genetic profiles. This work assesses the applicability, reliability, and discriminatory power of RAPD in distinguishing human DNA profiles from typical microbial contamination. The study developed random DNA profiles using human hair shaft DNA and compared the resulting banding patterns generated by *E. coli* as a model for non-human contamination within the microbiome. To assess the efficiency of RAPD, the study employed an experimental approach to evaluate the influence of sample condition, primer choice, and PCR parameters on the quality and reproducibility of profiles obtained from both human hair samples and *E. coli*. In addition, to understand the historical application of RAPD and to confirm its potential as a preliminary screening tool in modern forensic science, relevant journal publications were reviewed and synthesised in this work. Compared to the DNA profile obtained from *E. coli*, RAPD generated non-overlapping and unique banding patterns for human hair shaft DNA, demonstrating its efficiency in discriminating between the species under investigation. In addition to illustrating RAPD's capability for species-specific exclusion of microbial contaminants, this successful discrimination addresses crucial gaps in the reproducibility validation of RAPD for human trace evidence processing. The results affirm that RAPD is a reliable, rapid, and cost-effective initial screening method for problematic trace evidence. Responding to the need, as highlighted in recent literature, for validated rapid screening techniques, this research offers standardised guidelines that may advance the adoption of RAPD, particularly in resource-limited forensic laboratories, to support investigations in which traditional methods may fail.

Keywords: Forensic DNA analysis; human hair shaft; microbial contamination; random amplified polymorphic DNA; species discrimination

PP018

Occupational Radiation Exposure among Medical Radiation Personnel at Teaching Hospital

MUHAMMAD ZHARIF ZAINAL¹, ABDUL KHALIQ MOHD SAPARUDDIN^{1,2}, SITI HAJAR ZUBER^{1,2}, IZA NURZAWANI CHE ISA^{1,2}, SITI FARIDZWANA MOHD RIDZWAN³, S. ELAVARASI SELVARAJAH³, MUHAMMAD SAFWAN AHMAD FADZIL^{1,2*}

¹*Diagnostic Imaging and Radiotherapy Program, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Center for Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

³*Department of Radiology, Hospital Canselor Tuanku Muhriz, 56000 Cheras, Kuala Lumpur, Malaysia*

**Correspondence: safwanfadzil@ukm.edu.my*

ABSTRACT

Occupational radiation exposure monitoring is a fundamental component of radiation protection programmes, ensuring regulatory compliance and minimising health risks among medical radiation workers. Continuous dose surveillance also provides an objective measure of the effectiveness of radiation protection policies implemented within clinical departments. This study evaluated occupational radiation exposure among 105 medical radiation workers in the Department of Radiology, Hospital Canselor Tuanku Muhriz, over a three-year period from 2022 to 2024. Personal radiation doses were monitored for effective dose (Hp (10)) and skin dose (Hp(0.07)) using Radiophotoluminescent Dosimeters, with monthly dose readings obtained via the Asahi Techno RPL Reader (FGD-650). Radiographers recorded the highest mean annual dose (1.10 ± 0.28 mSv), followed by radiologists (1.08 ± 0.35 mSv) and medical officers (0.76 ± 0.14 mSv). Nevertheless, all recorded doses remained well below the International Commission on Radiological Protection (ICRP) recommended occupational dose limit of 20 mSv per year. In 2023, a notable decrease in occupational exposure was observed across all professions as COVID-19 cases declined, healthcare operations stabilised, and procedural volumes returned to lower levels. Occupational radiation exposure among medical radiation workers in this centre remained within acceptable safety limits throughout the study period, indicating effective implementation of radiation protection measures. This study offers crucial baseline data that can guide the refinement of radiation protection policies, specifically tailored to the evolving demands of a teaching hospital. Continuous, evidence-based improvements are vital for sustaining high standards of protection and safeguarding the long-term health and safety of medical workers in clinical practice

Keywords: Medical radiation workers; occupational dose; radiation protection; radiophotoluminescent dosimeter

PP019

NeuroHabit© App: AI-Assisted Engineering and Validation of a Hierarchical Scoring Engine for Modelling Neurochemical Balance in Behavioural Interventions

AUF ALI^{1,2*}¹*School of Pharmacy, KPI Healthcare University, 71800 Nilai, Negeri Sembilan, Malaysia*²*Founder and Developer of NeuroHabit© app by Aufthority, Nilai, Negeri Sembilan, Malaysia***Correspondence: ucn.auf@kpju.edu.my*

ABSTRACT

Many Mobile Health (mHealth) applications fail to sustain long-term behavior change because linear scoring systems ignore the hierarchical nature of human neurobiology. NeuroHabit© moves beyond «streak-based» gamification by implementing a bio-mathematical engine that prioritises physiological recovery over simple task accumulation. This study describes the algorithmic framework of the application and the protocol for its clinical validation. Claude AI was utilised for the high-fidelity calibration of a proprietary algorithm featuring a multiplicative foundational gate, $rawPct \times (0.5 + 0.5 \times foundationRate)$, which scales habit efficacy against recovery baselines (e.g., sleep). A four-week pilot study is planned ($n = 30$), utilising a purposive sampling strategy to include both neurotypical and neurodivergent cohorts. Validation will be conducted via a structured questionnaire measuring: (i) Face Validity of the qualitative “Brain Phase” signal, (ii) Tracking Anxiety via an adapted Perceived Stress Scale (PSS-10), and (iii) Habit Automaticity via the Self-Report Habit Index (SRHI). Preliminary AI-assisted stress-testing has already verified the algorithm’s sensitivity to category neglect and foundational deficiencies. The subsequent human pilot study aims to demonstrate that the “Hidden Score” philosophy significantly reduces tracking-related anxiety compared to traditional binary trackers. It is hypothesised that prioritising “Weight 3” foundational behaviors will lead to higher SRHI scores, indicating superior habit automaticity. By integrating AI-driven technical calibration with a planned human-centered validation, NeuroHabit© proposes that «de-gamifying» mHealth through homeostatic modeling is a viable path for lifestyle medicine. This research offers a scalable framework to reduce digital fatigue while improving the biological accuracy of behavioral interventions.

Keywords: AI-assisted design; behavioural informatics; habit automaticity; mHealth validation; neurochemical homeostasis

PP020

***In vitro* Differential Responses of Human SH-SY5Y Neuronal Cells and Human U87 Astrocyte-Like Cells to Glutamate and Docosahexaenoic Acid**

**MUHAMMAD HANI RAHIMI RUSLEEN¹, NUR IZZATI MANSOR^{1*},
ZAINAH MOHAMED¹, ADILA A HAMID², NURUL HAFIZAH MOHD
NOR³, YOGESWARAN LOKANATHAN⁴, WONG YING PING⁵**

¹*Department of Nursing, Faculty of Medicine, Universiti Kebangsaan Malaysia, Cheras 56000, Kuala Lumpur, Malaysia*

²*Department of Physiology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Cheras 56000, Kuala Lumpur, Malaysia*

³*Institute of Islamic Civilization, Universiti Kebangsaan Malaysia, Bangi 43600, Selangor, Malaysia*

⁴*Department of Tissue Engineering & Regenerative Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia, Cheras 56000, Kuala Lumpur, Malaysia*

⁵*Department of Pathology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Cheras 56000, Kuala Lumpur, Malaysia*

**Correspondence: nurizzatimansor@ukm.edu.my*

ABSTRACT

Glutamate-mediated excitotoxicity is a major pathological mechanism in neurodegeneration, marked by overstimulation of receptors, resulting in neuronal injury or death. Astrocytes are essential for maintaining glutamate homeostasis and preventing neuronal damage. Furthermore, docosahexaenoic acid (DHA), an omega-3 polyunsaturated fatty acid, is widely known for its neuroprotective effects. Most *in vitro* studies have focused on neuronal cell models, while the comparative effects of glutamate and DHA in neuronal and astrocytic cells remain unexplored. This study aimed to evaluate the effects of glutamate and DHA on cell viability and morphology of neuronal and astrocytic cells. Human SH-SY5Y cells were differentiated using 10 μ M retinoic acid for 7 days to generate neuronal cells, while human U87 cells were used as astrocyte-like cells. Both cells were treated with varying concentrations of glutamate (0, 50, 75, 100, 150, 200 mM) or DHA (0, 50, 75, 100, 150, 200 μ M) for 24 hour. Following treatment, cell viability was determined by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay, and morphological changes were examined microscopically. Glutamate exposure reduced SH-SY5Y cell viability, with a gradual decline starting at 50 mM. Morphological analysis showed neurite damage at 100 mM, with drastic neuronal degeneration at higher concentrations (≥ 150 mM). In contrast, U87 cells maintained consistent viability across glutamate concentrations, although at ≥ 150 μ M altered U87 cell morphology and reduced cell viability. Similarly, DHA exhibited cell-type-dependent effects. SH-SY5Y cells remained healthy at 50–100 μ M, whereas DHA at ≥ 150 μ M was toxic to neuronal cells, resulting in neurite and structural damage. In U87 cells, DHA appeared to promote proliferation

at all concentrations, as indicated by increased cell viability, healthy morphology, and higher density cultures. These findings indicate that neuronal and astrocytic cells have different susceptibilities to glutamate and DHA, supporting the suitability of these in vitro models for neurotoxicity and neuroprotection research.

Keywords: DHA; excitotoxicity; neuroprotection; Sh-Sy5y; U87

PP022

Temporal Stability and Forensic Reliability of Lip Prints: A Systematic Review

ATIKAH MOHD NASIR¹, RABIATUL ADAWIYAH NOR HISHAM¹,
NUR MAHIZA MD ISA², NOOR HAZFALINDA HAMZAH^{1*}

¹*Forensic Science Program, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Faculty of Veterinary Medicine, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia*

**Correspondence: raviera@yahoo.com*

ABSTRACT

Lip prints have attracted increasing interest as a supplementary forensic identifier, particularly in cases where fingerprints or DNA are limited, degraded, or unavailable. Their practical value, however, depends not only on individual uniqueness but also on whether lip print patterns remain sufficiently stable overtime. This systematic review aimed to evaluate the available evidence on the temporal stability of lip prints and to consider their reliability for forensic application. A structured search was conducted across seven electronic databases for studies published between 2000 and 2024. The review followed PRISMA 2020 guidance and included longitudinal studies that assessed lip print patterns at more than one time point in human participants. Six studies involving a total of 637 individuals met the inclusion criteria. Follow-up periods ranged from six months to three years, and both manual classification and digital comparison approaches were reported. Overall, the findings suggest that major lip print features remain relatively stable over time, with reported consistency rates ranging from 73.8 to 100%. Where variation occurred, it was often associated with differences in recording technique, print pressure, environmental exposure, or observer interpretation rather than true anatomical change. Studies incorporating digital methods appeared to demonstrate better reproducibility than those relying solely on manual classification. Despite these encouraging findings, the current evidence base remains limited by small study numbers, methodological heterogeneity, inconsistent recording conditions, and the absence of a standardised analytical framework. Lip prints therefore show promise as a supplementary forensic tool, but the available evidence does not yet support their routine use as a standalone method of identification. Further longitudinal research using standardised and digitally supported methods is needed to strengthen the evidential basis for forensic practice.

Keywords: Cheiloscopy; forensic science; lip prints; temporal stability

PP023

Awareness on Artificial Intelligence and Future Career Prospect among Full Time Medical Imaging Students in a Private Institution: A Brief Study

NIK ROSLINDA NIK MAT*, WAN FARHANA ARISSA WAN ROSMADEY

KPJ Healthcare University, 71800 Nilai, Negeri Sembilan, Malaysia

**Correspondence: ucn.roslinda@kpju.edu.my*

ABSTRACT

Artificial intelligence (AI) is rapidly transforming medical imaging through advancements in diagnostic accuracy, image processing, and workflow optimisation. Despite this progress, the level of AI literacy and workforce readiness among future practitioners remains insufficiently defined. This study aimed to critically evaluate AI awareness among full-time diploma and bachelor medical imaging students at KPJ Healthcare University, Nilai, and to examine its association with educational level alongside perceptions of AI's impact on future career trajectories. A cross-sectional design was conducted using a validated self-administered questionnaire (Cronbach's $\alpha = 0.848$). A total of 135 students were recruited through purposive sampling. Data were analysed using descriptive statistics and non-parametric inferential tests, with statistical significance set at $p < 0.05$. The findings revealed that while a majority of students demonstrated high overall awareness (77.0%), this awareness was predominantly superficial, with limited depth in understanding key applications such as machine learning and AI integration across imaging modalities. No statistically significant difference was observed between diploma and bachelor cohorts ($p = 0.052$), suggesting that higher educational attainment does not necessarily correspond to enhanced AI literacy. In terms of perception, students exhibited cautious optimism, recognising AI as a tool for professional enhancement while simultaneously expressing uncertainty regarding its implications for job security and evolving clinical roles. These findings highlight a critical gap between awareness and applied competency, indicating a misalignment between technological advancement and current educational practices. The persistence of uncertainty further reflects limited curriculum integration of AI-related competencies. This study underscores the urgent need for structured, competency-based curriculum integration to enhance AI literacy and ensure workforce readiness in an increasingly AI-driven healthcare environment.

Keywords: AI literacy; artificial intelligence; curriculum integration; medical imaging education; student perception; workforce readiness

PP024

Longevity of Change: Preliminary Qualitative Findings on Sustained Health Practices from the AGELESS Programme

FATIN IZZATY MOHD SHAHRIN, NURUL HIDAYAH MD FADZIL,
DEVINDER KAUR AJIT SINGH, ARIMI FITRI MAT LUDIN, HASNAH
HARON, SUZANA SHAHAR*

*Centre for Healthy Ageing and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300
Kuala Lumpur, Malaysia*

**Correspondence: suzana.shahar@ukm.edu.my*

ABSTRACT

The AGELESS multidomain intervention was developed to improve physical, cognitive, nutritional, psychosocial, and metabolic-vascular outcomes among older adults with cognitive frailty. Although its short-term benefits have been well established, sustaining these health-promoting behaviors beyond the intervention period remains a critical challenge in aging populations. Evidence on the long-term continuation of such multidomain practices is limited, highlighting a significant gap in understanding their real-world sustainability. Therefore, this study aims to explore which intervention components are maintained over time, and to identify the factors that facilitate or hinder their continued adoption. A qualitative follow-up study was conducted 12 months post-intervention among older adults who completed the two-year AGELESS multidomain programme. Semi-structured interviews were conducted with eight purposively selected participants. Interviews were transcribed verbatim and analysed using thematic analysis with ATLAS.ti software, guided by six pre-specified domains (sustained practices, motivators, barriers, perceived benefits, impact on daily life, and recommendations for sustainability). Preliminary findings revealed a pattern of selective sustainability, where participants predominantly maintained simple, routine-based physical activities (e.g., walking, household tasks), while structured and cognitive components were not sustained. Continuation was driven by intrinsic, health-oriented motivations, particularly maintaining mobility and preventing discomfort. A major barrier was the inability to recall intervention content, compounded by the absence of post-programme support and resources. Participants adapted practices by integrating them into daily routines and modifying them according to physical capacity. Perceived benefits were described in functional terms, including improved mobility, reduced pain, and enhanced independence in daily activities. Explicit recommendations were limited, though findings suggest a need for ongoing reinforcement and support mechanisms. Sustainability of multidomain interventions among older adults with cognitive frailty is shaped by practicality, memory retention, and contextual integration. Interventions aiming to promote longevity should prioritise routine-compatible strategies and incorporate sustained support to enhance long-term impact.

Keywords: Cognitive frailty; multidomain intervention; older adults; qualitative; sustainability

PP025

The Effectiveness of Dynamic Strengthening Exercise Between Pain and Functional Activity for Mechanical Low Back Pain in Asia – A Systematic Review

SITI NORHIDAYAH SADIO, NURIN KHAIRUNISA AZMIRUDDIN

KPJ Healthcare University Nilai, Malaysia

**Correspondence: ucn.norhidayah@kpju.edu.my*

ABSTRACT

Mechanical low back pain (MLBP) is a prevalent condition in Asia, exacerbated by sedentary lifestyles, poor ergonomics, and occupational hazards. This systematic review evaluates the effectiveness of dynamic strengthening exercises (DSE) in reducing pain and improving functional activity among individuals with MLBP in Asia. A total of five peer-reviewed studies conducted between 2019 and 2024, involving 210 participants aged 20–60, were analysed. The interventions included lumbar stabilisation, spinal extension, and core strengthening exercises, with outcomes measured using the Visual Analogue Scale, Numerical Pain Rating Scale and Oswestry Disability Index. Results consistently demonstrated significant reductions in pain intensity and improvements in functional activity, outperforming passive modalities and general aerobic exercises. DSE effectively targets stabilising muscles such as the transversus abdominis, multifidus, and erector spinae, enhancing spinal alignment, trunk control, and postural stability. Additionally, combining DSE with ergonomic education further amplified therapeutic outcomes. The review highlights DSE as a cost-effective, accessible, and culturally adaptable intervention for MLBP management in Asia. However, limitations such as short intervention durations, small sample sizes, and lack of long-term follow-up underscore the need for further research. This study supports the integration of DSE into clinical practice and public health strategies to address the growing burden of MLBP in Asia.

Keywords: Dynamic strengthening exercises; functional activity; mechanical low back pain; pain

PP027

Cytotoxicity and Selective Anticancer Potential of Triphenyltin(IV) N-Methyl-N-Benzylthiocarbamate in Hepatocellular Carcinoma (HepG2) and Human Embryonic Kidney (HEK293) Cells

NORMAH AWANG^{1*}, PUTRI MALAYA TASLIM², NURUL FARAHANA KAMALUDIN¹

¹Center of Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Environmental Health & Industrial Safety Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

*Correspondence: norm@ukm.edu.my

ABSTRACT

Hepatocellular carcinoma (HCC) remains a major global health burden and is characterised by high mortality rates, therapeutic resistance, and limited responsiveness to conventional chemotherapy. Although agents such as 5-fluorouracil (5-FU) are clinically employed, their therapeutic utility is frequently compromised by systemic toxicity and acquired resistance. Therefore, the identification of novel chemotherapeutic scaffolds with enhanced selectivity and mechanistic precision is urgently required. Organotin(IV) compounds have emerged as promising antitumor candidates due to their pronounced cytotoxic potency, structural tunability, and ability to interact with critical intracellular targets. In the present study, the cytotoxic and selective anticancer potential of triphenyltin(IV) N-methyl-N-benzyl dithiocarbamate was systematically evaluated against human hepatocellular carcinoma (HepG2) cells and non-malignant human embryonic kidney (HEK293) cells. Cell viability was quantified using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reduction assay following 24-hour exposure, and half-maximal inhibitory concentration (IC_{50}) values were determined. Morphological assessments were conducted to characterise the mode of cell death. The compound exhibited marked cytotoxicity toward HepG2 cells, with an IC_{50} value of 0.70 μ M, while demonstrating comparatively lower toxicity in HEK293 cells (IC_{50} = 1.97 μ M). The calculated selectivity index ($SI = 2.81$) indicates preferential cytotoxic action toward malignant cells. Microscopic examination revealed classical apoptotic features, including cell shrinkage, chromatin condensation, membrane blebbing, and loss of adherence, suggesting activation of programmed cell death pathways rather than nonspecific necrosis. The enhanced anticancer activity may be mechanistically associated with the stability of Sn–N and Sn–S coordination bonds, which confer structural rigidity and facilitate sustained intracellular bioactivity. The lipophilic triphenyltin moiety likely enhances membrane permeability, promoting efficient cellular uptake and interaction with mitochondrial or nuclear targets, potentially triggering oxidative stress and apoptotic signaling cascades. Collectively, these findings demonstrate that triphenyltin(IV) N-methyl-N-benzyl

dithiocarbamate possesses potent and selective antiproliferative activity against hepatocellular carcinoma cells. The compound represents a promising organotin-based chemotherapeutic lead, warranting further investigation through mechanistic studies involving apoptosis-related biomarkers, cell cycle profiling, reactive oxygen species modulation, and comprehensive in vivo efficacy and toxicity evaluations.

Keywords: Apoptosis; HEK293; HepG2; MTT assay; organotin(IV); selectivity index, triphenyltin dithiocarbamate

PP028

Refractive and Visual Acuity Profiles in Keratoconus: A Retrospective Review at Universiti Kebangsaan Malaysia Optometry Clinic (2017–2024)

**BASHIRAH ISHAK*, HALIZA ABDUL MUTALIB, WAN MUHAMMAD
DZARRIDDIN WAN AZHAR, NURIN NADHIRAH MOHD YUSOFF,
NORSHAHIDATUL AKMA NAZARUDIN**

*Program of Optometry and Visual Sciences, Faculty of Health Sciences, Universiti Kebangsaan Malaysia,
50300 Kuala Lumpur, Malaysia*

**Correspondence: bashirah@ukm.edu.my*

ABSTRACT

Keratoconus is a progressive ectatic corneal disorder that causes marked changes in refractive error and deterioration of visual acuity. This study aimed to characterise the refractive error and visual acuity profiles of keratoconus patients according to disease severity and to evaluate the impact of optical interventions, including spectacles and contact lenses. A retrospective review was conducted at the Optometry Clinic, Universiti Kebangsaan Malaysia, involving 71 patient records collected between 2017 and 2024. Analysis showed no significant change in refractive error at the first follow-up ($p = 0.620$), whereas highly significant changes were detected at the second and third follow-ups ($p < 0.001$). Visual acuity differed significantly across severity levels ($p = 0.001$), with patients in the mild stage demonstrating the highest mean rank (48.10). In contrast, those in severe to most severe stages had substantially lower values (22.40–23.91). The Friedman test revealed a statistically significant improvement in visual acuity across the three visits ($\chi^2 = 99.852$, $df = 2$, $p < 0.001$), indicating that optical correction with spectacles or contact lenses plays a crucial role in enhancing visual acuity over time. This study provides a longitudinal clinical overview of keratoconus patients in an optometric setting, supporting more tailored management strategies based on disease severity. Additionally, the results highlight the need for improved electronic clinical documentation to support future research and patient monitoring. From a policy and practice standpoint, the findings reinforce the importance of early detection and continuous follow-up in keratoconus management, with implications for primary care guidelines and optometric training programmes.

Keywords: Keratoconus; refractive error; severity; visual acuity

PP029

Content Validity of a Newly Developed Paediatric Vision Screening Index

BAQIATU'L SABIQI 'ASSFI RAHMAT^{1,2*}, MD MUSTAFA MD-MUZIMAN-SYAH^{2,3}, NORAISHAH MOHAMED NOR⁴, THASHINI SANMUGAM⁵, NOOR FAKHIRA MUNEERA ABDUL HALIM²

¹Department of Optometry, Rehabilitation and Well-being, Faculty of Health and Life Sciences, Management and Science University, 40100 Shah Alam, Selangor, Malaysia

²Department of Optometry and Visual Science, Kulliyah of Allied Health Sciences, International Islamic University Malaysia, 25200 Kuantan, Pahang, Malaysia

³Children Health and Well-Being Research Group, Kulliyah of Allied Health Sciences, International Islamic University Malaysia, 25200 Kuantan, Pahang, Malaysia

⁴Department of Nutrition Sciences, Kulliyah of Allied Health Sciences, International Islamic University Malaysia, 25200 Kuantan, Pahang, Malaysia

⁵School of Graduate Studies, Management and Science University, 40100 Shah Alam, Selangor, Malaysia

*Correspondence: baqiatul_sabiqi@msu.edu.my

ABSTRACT

Question-based paediatric vision screening tools offer accessible, scalable alternatives to conventional screening approaches. However, ensuring robust content validity is essential to establish conceptual adequacy and representativeness of screening domains prior to further psychometric evaluation. This study aimed to evaluate the content validity of a newly developed paediatric vision screening index comprising parent- and child-reported versions in both English and Malay. A methodological validation study was conducted in two sequential stages: content validation of the original English version, followed by translation and content validation of the Malay version. Twenty-one subject matter experts, including ophthalmologists and optometrists actively involved in paediatric eye care, independently rated each item using a four-point relevance scale. Item-level Content Validity Index (I-CVI) and Scale-level Content Validity Index based on the average method (S-CVI/Ave) were computed. Items that failed to meet the recommended threshold values were either excluded or revised in response to expert feedback. For the English version, the parent questionnaire demonstrated I-CVI values ranging from 0.56 to 1.00 and S-CVI/Ave values between 0.81 and 0.91. A total of 21 items were excluded, and 8 items were revised. The child version showed I-CVI values ranging from 0.44 to 1.00 and S-CVI/Ave values between 0.84 and 0.94, resulting in the exclusion of 14 items and revision of 12 items. Following forward-backward translation and expert review, the Malay version demonstrated improved content validity. The parent version showed I-CVI values ranging from 0.82 to 1.00 and S-CVI/Ave values between 0.93 and 0.97, while the child version showed I-CVI values ranging from 0.82 to 1.00 and S-CVI/Ave values between 0.86 and 0.97. No further items were excluded in the Malay validation phase. The newly developed paediatric vision screening index demonstrated satisfactory content validity in both English and Malay versions after systematic expert evaluation and refinement.

Keywords: Content validity; paediatric vision screening; question-based

PP030

Knowledge and Attitudes of Early Childhood Development Practitioners Towards Childhood Hearing Loss and Hearing Services

XUE MIN GOH¹, RAFIDAH MAZLAN^{1,2*}, ONN WAH LEE^{1,2}, MASAYU DZAINUDIN³

¹Audiology Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Centre for Rehabilitation and Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

³Department of Early Childhood Education, Faculty of Human Development, Sultan Idris Education University, 35900 Tanjong Malim, Perak, Malaysia

*Correspondence: rafidahmazlan@ukm.edu.my

ABSTRACT

Early identification of childhood hearing loss depends not only on screening systems but also on the knowledge and attitudes of early childhood development (ECD) practitioners who observe children in everyday learning environments. This study examined levels of knowledge and attitudes toward childhood hearing loss among Malaysian ECD practitioners, the relationship between knowledge and attitudes, and demographic factors associated with these outcomes. A cross-sectional quantitative survey was conducted with 849 practitioners from public and private early childhood centers across Malaysia. Data were collected using an adapted self-administered questionnaire assessing knowledge and attitudes related to childhood hearing loss. Descriptive statistics summarised responses, while Spearman correlation and logistic regression analyses were used to examine associations and predictors. Overall, 66.4% of practitioners demonstrated high knowledge. Higher correct responses were observed in the Impact (86.7%) and Signs and Symptoms (83.5%) domains, while knowledge was lowest in Identification and Intervention (52.8%). Only 30.3% of participants felt equipped to support a child with hearing loss in the classroom. Positive attitudes were observed in 59.4% of practitioners; however, only 27.4% believed that children with hearing loss could attend mainstream classrooms. A weak but significant positive association was found between knowledge and attitudes ($p = 0.201$, $p < 0.05$). Gender, education level, and nature of employment predicted knowledge, while age and education level were associated with attitudes. ECD practitioners demonstrated basic knowledge of childhood hearing loss, but gaps remain in procedural knowledge, referral understanding, and confidence in classroom inclusion. Strengthened training, clearer referral guidance, and cross-sector collaboration between education and health services are needed to support early identification and inclusive practice in early childhood settings.

Keywords: Childhood hearing loss; early childhood workforce; early identification; health and education system; inclusive education; knowledge and attitudes; Malaysia

PP032

Predictors of Metabolic Dysfunction-Associated Fatty Liver Disease

SHANTHI KRISHNASAMY^{1*}, HO LI TAN¹, NORHASHIMAH NORSUDDIN¹, DEBORAH CHIA HSIN CHEW², NIK MADIHAH NIK AZIZ³, SYARATUL DALINA YUSOFF⁴, AZNIDA FIRZAH ABDUL AZIZ⁵

¹Center For Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Gastroenterology And Hepatology, Faculty of Medicine, Hospital Canselor Tuanku Muhriz, Universiti Kebangsaan Malaysia, 56000, Cheras, Kuala Lumpur, Malaysia

³Restorative Department, Faculty of Dentistry, Universiti Kebangsaan Malaysia, 50300, Kuala Lumpur, Malaysia

⁴Faculty of Pharmacy, 50300, Kuala Lumpur, Malaysia

⁵Department of Family Medicine, Faculty of Medicine, Hospital Canselor Tuanku Muhriz, Universiti Kebangsaan Malaysia, 56000, Cheras, Kuala Lumpur, Malaysia

*Correspondence: shanthikrishnasamy@ukm.edu.my

ABSTRACT

Metabolic dysfunction-associated fatty liver disease (MAFLD) is becoming increasingly prevalent due to the rising rates of overweight, obesity, and metabolic syndrome among Malaysian adults. This study aimed to assess the association between cardiometabolic factors, dietary intake, and oral health status with MAFLD among staff and students at the National University of Malaysia (UKM). A cross-sectional study was conducted among staff and students aged 18–65 years. Participants completed questionnaires on sociodemographic characteristics, medical history, physical activity, and oral health status, as well as a 3-day food diary to assess dietary intake. Anthropometric measurements were obtained, and liver stiffness and steatosis were assessed using transient elastography (FibroScan). Blood biomarkers, including liver function tests, full lipid profile, fasting blood glucose, and HbA1c, were also measured. Univariate and multivariate logistic regression analyses were performed to identify predictors of MAFLD, while Spearman's rank correlation was used to examine the associations between anthropometric measures, cardiometabolic factors, physical activity, dietary intake, and liver steatosis and stiffness. A total of 105 participants with a median age of 36.0 (IQR 16.0) years were included in the study. The overall prevalence of MAFLD among UKM adults was 19.0%. HbA1c (aOR = 15.36, 95% CI: 3.20–73.66, $p = 0.001$) and HDL-C (aOR = 0.044, 95% CI: 0.003–0.548, $p = 0.015$) were identified as independent predictors of MAFLD. Liver steatosis was strongly correlated with age, body mass index, waist circumference, body fat percentage, glycaemic indices, and triglyceride levels, and was inversely associated with HDL-C (all $p < 0.001$). Metabolic dysfunction, central obesity, and excess energy intake appear to be key drivers of MAFLD among staff and students at UKM, highlighting the importance of early identification of metabolic risk factors and caloric intake reduction in the prevention of MAFLD.

Keywords: Dietary intake; fatty liver; metabolic syndrome

PP034

Clinical Differentiation of Structural Language and Functional Communication Profiles in Post-Stroke Aphasia and Alzheimer's Disease Within a Multilingual Malaysian Cohort: A Retrospective Analysis

NUR FARAH FARHANA JAMIL*, HANI MADIHAH BAHARI,
NUR NAZIHAH HAZIZI

*Department of Rehabilitation Medicine, Hospital Al-Sultan Abdullah, Universiti Teknologi MARA, 42300
Puncak Alam, Selangor, Malaysia*

**Correspondence: nurfarah04@uitm.edu.my*

ABSTRACT

Communication impairment in older adults is frequently attributed to post-stroke aphasia (PSA) or Alzheimer's disease (AD), yet differentiating language-driven impairment from cognitively mediated communication decline remains clinically challenging, particularly in multilingual populations. This study compared structural language and functional communication performance between PSA and AD in a Malaysian tertiary rehabilitation cohort and examined whether multilingual status moderated these differences. A retrospective comparative chart review was conducted among patients aged 60 years referred for speech-language assessment between January 2023 and December 2025. Thirty patients with medically confirmed PSA (64.2 ± 7.8 years; 63% male) were compared with 35 patients with medically confirmed AD (72.6 ± 6.9 years; 68% female). Structural language performance was assessed using the Quick Aphasia Battery (QAB) total score. Functional communication was evaluated using the Derby Functional Communication Scale (DFCS) across Expression, Understanding and Interaction domains. Global cognition was measured using the Montreal Cognitive Assessment or Mini-Mental State Examination. Language history classified patients as monolingual or multilingual. Two-way factorial analysis of variance examined main and interaction effects, with effect sizes reported using partial eta squared (η^2_p) and Cohen's d. PSA demonstrated significantly lower QAB total scores than AD ($p < 0.05$; moderate-to-large effect size), indicating pronounced structural language impairment. Significant differences were observed across DFCS domains ($p < 0.05$; small-to-moderate effect sizes), with PSA showing reduced functional communication performance. AD exhibited significantly lower cognitive screening scores ($p < 0.05$; moderate effect size). Two-way analyses indicated significant main effects of diagnosis (η^2_p in the small-to-moderate range), while diagnosis-by-language interaction effects were small. These findings demonstrate statistically and clinically meaningful differentiation between PSA and AD within a multilingual Malaysian cohort and support integrating structural and functional measures to enhance diagnostic precision and rehabilitation planning in ageing populations.

Keywords: Alzheimer's disease; cognitive-communication disorders; functional communication; post-stroke aphasia; multilingualism

PP035

Comparative Study: Commercial Mineral Water as an Alternative Diluent for 10% Giemsa Staining in Malaria Diagnosis

AHMAD AZRI MOHAMAD^{1*}, ROHAYA ABDUL HALIM², MUHAMMAD HASIF WAQIF AMINUDDIN², SITI SAIYIDAH S.MONIL², NUR FARAH SYAFIQAL MOHD FADIL², NUR FARAHEEN SAHLI³

¹Department of Physical Therapy, Faculty of Allied Health Sciences, Naresuan University, Phitsanulok, 65000 Thailand

²Exercise and Rehabilitation Sciences Research Unit, Faculty of Allied Health Sciences, Naresuan University, Phitsanulok, 65000 Thailand

*Correspondence: ahmadazri.ilkkmkl@gmail.com

ABSTRACT

Malaria is a parasitic disease caused by *Plasmodium* spp. and transmitted through the bite of female Anopheles spp. mosquitoes. Microscopic examination of Giemsa-stained blood films remains the gold standard for malaria diagnosis as recommended by the World Health Organisation (WHO). Standard preparation of 10% Giemsa requires freshly prepared phosphate buffer at pH 7.2 to ensure optimal staining quality. However, buffer preparation requires distilled water, which is time-consuming and costly. Although tap water has been suggested as an alternative diluent, its inconsistent pH across different locations may affect staining quality. This study aimed to evaluate Mesra, a locally produced and cost-effective commercial mineral water brand that is accessible, as a potential alternative diluent for 10% Giemsa staining. A total of 52 positive malaria slides stored at -30°C were used. The slides were divided into two groups (n = 26 each). Each group was stained with 10% Giemsa prepared using either phosphate buffer (pH 7.2) or commercial mineral water with consistence pH 7.33. The staining results were evaluated by two screeners and the mean scores were recorded based on a scale 1 (poor) and 2 (good). The mean staining score for Giemsa prepared with commercial mineral water (1.85) was slightly higher than that prepared with phosphate buffer (1.79). However, statistical analysis revealed no significant difference between the two diluents (p = 0.639), indicating that commercial mineral water produces staining quality comparable to the standard phosphate buffer. Commercial mineral water is a viable, ready-to-use and inexpensive alternative to phosphate buffer for Giemsa staining. Its implementation can significantly simplify malaria microscopy in resource-limited settings without compromising diagnostic accuracy.

Keywords: Commercial mineral water; giemsa staining; malaria; microscopic diagnosis; phosphate buff

PP039

MyLangLab Narrative Database

NUREHAN AZHAR¹, TAN WONG JIN¹, YAZMIN AHMAD RUSLI^{1,2*}

¹*Speech Sciences Program, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Center for Rehabilitation & Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

**Correspondence: yazmin@ukm.edu.my*

ABSTRACT

Access to culturally and linguistically relevant language data has remained limited for speech-language therapists (SLTs) in Malaysia. Assessment and intervention practices have frequently relied on international language sample resources that do not fully reflect the multilingual and multicultural linguistic environment of Malaysian children. Rather than reflecting limitations in assessment accuracy, this situation highlights the need for innovative spaces that support the development of locally relevant assessment and intervention resources, as well as training materials for students and clinicians. To address this need, the MyLangLab Narrative Database project was conducted to develop a localised repository of narrative language samples to support teaching, research, and clinical training within the Speech Science Program at Universiti Kebangsaan Malaysia (UKM). The project focused on creating a prototype database that compiles narrative language data reflecting local linguistic contexts. The study adopted a qualitative design guided by the Analysis, Design, Development, Implementation and Evaluation model. During the analysis phase, existing international language databases were reviewed to identify best practices in database structure, accessibility, and data organisation. These insights informed the design of a blueprint and user manual outlining the database architecture and guidelines for data entry and retrieval. During the development phase, a working prototype was constructed incorporating a sample set of clinical narrative data. The prototype was then implemented through a pilot evaluation involving approximately 25 participants, including students and faculty members from the Speech Science Program at UKM. Participants interacted with the database and provided feedback regarding usability, functionality, and relevance for teaching and research purposes. User responses were analysed thematically to identify strengths, limitations, and areas for refinement. The outcome of the project was a functional prototype and a structured framework for a sustainable local language database. MyLangLab is expected to enhance access to culturally relevant language samples, support evidence-informed practice in speech-language therapy, and strengthen educational resources for future SLTs in Malaysia. Ultimately, the initiative contributes to the continued development of language assessment and intervention practices within the Malaysian context.

Keywords: Multilingual children; narrative; speech-language therapy clinical resource

PP040

High-Frequency 5G Radiofrequency Exposure Accelerates Testicular Ageing via Apoptosis in Male Rats

FARAH HANAN FATHIHAH JAFFAR¹, SYED MUHAMAD ASYRAF SYED TAHA¹, ATIKAH HAIRULAZAM¹, AINI FARZANA ZULKIFLI¹, MOHD FARISYAM MAT ROS¹, KHAIRUL OSMAN², ZAHRILADHA ZAKARIA³, MOHD AMYRUL AZUAN MOHD BAHAR⁴, SITI FATIMAH IBRAHIM^{1*}

¹*Department of Physiology, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia*

²*Forensic Science Program, Center for Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

³*Faculty of Electronics & Computer Technology and Engineering, Universiti Teknikal Malaysia, Durian Tunggal, 76100 Melaka, Malaysia*

⁴*Intel Microelectronics (M) Sdn. Bhd., Bayan Lepas Technoplex, Medan Bayan Lepas, Intel Corporation, 11900 Bayan Lepas, Pulau Pinang, Malaysia*

ABSTRACT

The testis is highly susceptible to radiofrequency radiation (RFR), and apoptosis is a key mechanism underlying testicular ageing. With the deployment of 5G networks operating at 3.5 GHz and 24 GHz, potential impacts on male reproductive health remain poorly understood. This study was conducted to investigate the effect of chronic high-frequency 5G exposure on apoptotic pathways in male rat testes. A total of eighteen (N = 18) male rats were randomly assigned to Control, 3.5 GHz, or 24 GHz exposure groups with n = 6 for each group. Exposure was conducted for 7 hours/day for 60 days. Testicular apoptosis was evaluated via caspase-3, p53, and Bax mRNA expression, and the apoptotic index was validated through the TUNEL assay. Findings showed that exposure to 3.5 GHz frequency caused significant upregulation of caspase-3 (0.814 ± 0.115) compared with the 24 GHz group (0.372 ± 0.050 , $p < 0.05$), indicating ongoing apoptosis in 3.5 GHz. Meanwhile, lower caspase-3 mRNA expression in the 24 GHz exposure suggests a possible adaptive stress response. The mRNA findings were consistent with the apoptotic index at 3.5 GHz (6.382 ± 0.677), which was significantly higher than that of the Control (0.434 ± 0.220 , $p = 0.0004$). Interestingly, p53 and Bax expression remained unchanged in both exposed groups, implying apoptosis occurred via alternative, non-canonical pathways. This observation may represent an atypical biological response to RFR exposure. In conclusion, chronic high-frequency 5G exposure induces frequency-dependent apoptotic changes in the testis, consistent with accelerated testicular ageing. Exposure at 3.5 GHz appears to cause a more pronounced effect on testicular apoptosis compared to 24 GHz. These findings underscore the need for further investigation and precautionary measures to protect male reproductive health amid widespread 5G deployment.

Keywords: 5G; caspase-3; testicular apoptosis

PP043

A Retrospective Study on Audible Contrast Threshold among Hearing Impaired Adults in Audiology Clinic: A Pilot Study

AINU MARDHIAH^{1,2}, CHONG FOONG YEN^{1*}, LEE ONN WAH¹

¹Center for Rehabilitation & Special Needs, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Global Precision Sales & Services Sdn Bhd, C-02-09 and C-03-09, Aurora Place, 50700 Kuala Lumpur, Malaysia

*Correspondence: oongyen.chong@ukm.edu.my

ABSTRACT

Despite advancements in Digital Noise Reduction (DNR) over the decades, speech-in-noise perception remains a major challenge for hearing aid (HA) users. The Audible Contrast Threshold (ACT) test provides an objective and language-independent estimate of speech-in-noise perception ability, and its results can be used to guide audiologists in personalising DNR settings in HAs. However, the ACT remains underexplored in clinical practice, particularly in Malaysia, and only limited studies are available globally. In this study, test duration, number of attempts, and participants' subjective feedback were collected at a private HA centre in Malaysia to evaluate its clinical practicality. A total of 25 patients ($n = 25$) with hearing loss (mean age = 58 years; range = 33–80 years) were recruited between November 2025 and March 2026, and 56% ($n = 14$) of them completed this test. The ACT was brief to administer, with a mean duration of 2.47 minutes ($SD = 0.81$; range = 1–4 minutes). Among participants who completed the test, nine required one attempt and five required two attempts. The findings showed that participants who were unable to complete the ACT had significantly poorer hearing thresholds ($M = 65.43$ dB HL, $SD = 16.94$) compared with those who completed the test ($M = 52$ dB HL, $SD = 16$), $t = 2.07$, $p = 0.05$. Some participants reported difficulty distinguishing between “siren-like” and “noise-like” stimuli, particularly those who were unable to complete the task. In addition, the analysis revealed a non-significant correlation between the better-ear Pure Tone Average (PTA) and ACT score ($r = 0.466$), possibly due to the small sample size ($n = 14$). In conclusion, these preliminary findings indicate that the ACT is practical to administer within a short clinical timeframe and demonstrates measurable associations with hearing thresholds.

Keywords: ACT; adults; DNR; hearing aids; practicality

PP044

Developing a Healthy Food Marketing Framework for Malaysian Youth through Stakeholder Collaboration and Participatory Action Research: A Multi-Phase Study Protocol

CHRISTINE JOAN¹, SHU CHIN TEH¹, MOHD JAMIL SAMEEHA^{1,2*},
ELAINE Q. BORAZON³, AVITA USFAR⁴, SIRINYA PHULKERD⁵, NISHA
ARUNATILAKE⁶, BEE KOON POH^{1,2}, ON BEHALF OF THE SHAPE ASIA
STUDY GROUP

¹*Centre for Community Health Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300
Kuala Lumpur, Malaysia*

²*Obesity-UKM Research Group, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia*

³*College of Social Sciences, National Sun Yat-sen University, Kaohsiung City 804, Taiwan*

⁴*Reconstra Utama Integra, Jakarta 12150, Indonesia*

⁵*Institute for Population and Social Research, Mahidol University, Phutthamonthon District, Nakhon Pathom
73170, Thailand*

⁶*Institute of Policy Studies, Colombo 07, 00700, Western Province, Sri Lanka*

**Correspondence: sameeha@ukm.edu.my*

ABSTRACT

The rising prevalence of childhood obesity in Malaysia highlights an urgent need for strategies that address obesogenic food environments and promote healthier dietary habits, particularly among children who are most vulnerable to unhealthy food marketing. Healthy food marketing (HFM) offers a promising approach by countering the dominance of unhealthy food promotions and encouraging healthier dietary behaviours. This study protocol is part of the SHAPE Asia (Malaysia) project and outlines a multi-phase approach to develop a contextually relevant, stakeholder-informed HFM framework to support youth in navigating food marketing and adopting healthier dietary practices. Phase 1 involves a bibliometric review to map global and regional evidence on HFM, with a focus on consumer empowerment approaches. This phase will identify research gaps, highlight effective communication channels, and contextualise Malaysia's current food marketing landscape. Phase 2 focuses on stakeholder mapping and shared priority setting to identify and engage policymakers, government ministries, civil society, international agencies, and businesses. Using a mixed-methods approach comprising focus group discussions and surveys, this phase will explore perceptions of unhealthy food marketing, barriers to HFM implementation, and opportunities to empower youth. It will generate shared understanding of priorities and foster readiness for collaborative action. Phase 3 employs Participatory Action Research, in which stakeholders co-design intervention strategies to improve food marketing practices within their organisations. Strategies will be piloted and refined through two iterative cycles of planning, action, and reflection. Stakeholder reflection sessions and feedback from youth as experiential stakeholders will guide refinement to ensure cultural relevance, feasibility, and sustainability within the Malaysian context. This study is expected to generate an actionable HFM framework tailored to Malaysia, strengthen multi-sector collaboration to transform

food marketing environments, and equip youth with the skills to adopt healthier lifelong dietary habits. The framework will also guide policy and stakeholder practices with locally relevant strategies.

Keywords: Adolescents; food environment; health promotion; healthy food marketing; participatory research

PP045

Caffeine Consumption Patterns among Malaysian University Athletes: Implications for Safe Intake and Community Health

SITTI JUNAINA MUSA¹, NEROSHA NAIR GUNASEGARAN¹, AHMAD ROHI GHAZALI², NOR FARAH MOHAMAD FAUZI^{1,5}, MOHD IZHAM MOHAMAD³, ABDUL HADI ABD RAHMAN⁴, NIK SHANITA SAFII^{1,5*}

Center for Community Health Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Center for Toxicology & Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

³Sports Nutrition Centre, National Sport Institute, 57000 Kuala Lumpur, Malaysia

⁴Center for Artificial Intelligence Technology, Faculty of Information Sciences and Technology, University Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

⁵Obesity-UKM Research Group, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

**Correspondence: nikshanita@ukm.edu.my*

ABSTRACT

Caffeine is one of the most widely consumed ergogenic aids in sport settings due to its ability to enhance performance and cognitive alertness. However, there is limited awareness of safe caffeine consumption among Malaysian university athletes and assessing their adherence to recommended intake limits. A cross-sectional study was conducted among 236 athletes (135 males, 101 females) using the validated Modified Caffeine Consumption Questionnaire (M-CCQ). Caffeine content of commonly consumed beverages was determined using High-Performance Liquid Chromatography (HPLC) to establish accurate intake estimates. Descriptive statistics were used to summarise consumption patterns, and Spearman's correlation analysis was performed to examine associations between beverage types. Overall, median caffeine intake was 243 mg/day (0–919.5) with 69.1% of athletes exceeding the recommended daily intake limit. Ready-to-drink Nescafé canned coffee was the most consumed product (34%) with an intake of 151 mg/day (118 - 200), followed by Nescafé 3-in-1 (33%) with 172 mg/day (86 - 258). Although ground coffee ranked third in consumption frequency, it contributed the highest median daily caffeine intake of 1056 mg/day (528-1056), representing the highest single-day intake observed. Notably, caffeine content per serving of several commercially available coffee products exceeded the recommended 50 mg/serving threshold. Moderate positive correlation was observed between intake of instant coffee mixes and canned coffee ($p = 0.46$, $p < 0.001$), suggesting concurrent consumption of multiple caffeinated beverages. The high prevalence of caffeine intake among university athletes raises public health concerns. Inadequate labelling of caffeine content in commercially available beverages may contribute to unintentional overconsumption, potentially increasing the risk of adverse effects such as sleep disturbances and impaired recovery. These findings highlight the need for greater awareness, improved product labelling, and targeted educational interventions to support safer caffeine consumption practices among athletes and promote community health.

Keywords: Athletes; caffeine; overconsumption; product labelling

PP048

Differentiation of *Escherichia coli* and *Staphylococcus aureus* in Mixed Suspensions Using Dielectrophoresis

AKMAL SUHAIMI¹, ARASH ZULKARNAIN¹, NORAZIAH MOHAMAD ZIN^{2*}, MUHAMAD RAMDZAN BUYONG¹

¹Institute of Microengineering and Nanoelectronics, Universiti Kebangsaan Malaysia, 43600 Selangor, Malaysia

²Center for Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Science, Universiti Kebangsaan Malaysia, 43600 Selangor, Malaysia

*Correspondence: noraziah.zin@ukm.edu.my

ABSTRACT

Rapid identification of bacterial species is important for clinical diagnostics and infection control. Conventional microbiological methods often require lengthy culture and biochemical analyses, which can delay pathogen identification. In this study, dielectrophoresis (DEP) was investigated as a label-free technique for differentiating bacterial species in mixed suspensions. Two clinically relevant bacteria, *Escherichia coli* and *Staphylococcus aureus*, representing Gram-negative and Gram-positive organisms, respectively, were used as model species. Bacterial suspensions were prepared in a medium with controlled conductivity (0.03 S/m) to ensure consistent DEP conditions. Mixed bacterial samples were prepared using different dilution ratios to vary the relative cell concentrations. In Mixture A, *E. coli* was diluted at 1:100 and *S. aureus* at 1:1000, while in Mixture B, the dilution ratios were reversed. DEP experiments were conducted at 30 kHz and 300 kHz based on previously characterised responses of the two species. Particle trajectories were recorded and analysed, and 100 particles were tracked for each experiment. In the *E. coli*-rich mixture, 86% of detected particles were identified as *E. coli*, while 14% were identified as *S. aureus*. Conversely, in the *S. aureus*-rich mixture, 64% of detected particles were identified as *S. aureus* and 36% as *E. coli*. These results demonstrate that DEP responses can be used to differentiate bacterial species in mixed suspensions based on their distinct electrokinetic behaviours. The findings highlight the potential of DEP as a rapid and label-free approach for bacterial differentiation, which may support faster pathogen screening in clinical diagnostics as well as monitoring of bacterial contamination in environmental and water samples.

Keywords: Bacterial differentiation; dielectrophoresis; *Escherichia coli*; label-free diagnostics; *Staphylococcus aureus*

PP050

PCOS Phenotype Predicts Follicular Development and Treatment Response: A Prospective 3D Transvaginal Ultrasound Study

M.L. DINESH^{1*}, B.R. SHASINDRAU¹, MOHD HAZEMAN ZAKARIA²,
EFA'HUZAIMAH AFFENDI¹

¹Department of Medical Imaging, KPJ Healthcare University, Kota Seriemas, 71800 Nilai, Negeri Sembilan, Malaysia

²Pusat Pengimejan Diagnostik Nuklear, Hospital Sultan Abdul Aziz Shah, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia

*Correspondence: dinesh@kpju.edu.my

ABSTRACT

Polycystic ovary syndrome (PCOS) remains under-characterised in Malaysian primary care despite being a leading cause of anovulatory infertility. This study aimed to determine the distribution of PCOS phenotypes among infertile Malaysian women, examine demographic associations, and evaluate phenotype-specific follicular development changes following infertility treatment using three-dimensional transvaginal ultrasound (3D TVS). A prospective observational cohort study recruited 122 married women aged 20-40 years with Rotterdam-diagnosed PCOS at a private clinic in Selangor, Malaysia between January 2023 and December 2024. Participants were classified into phenotypes A-D and grouped as classic (A/B) or non-classic (C/D). Baseline and post-treatment 3D TVS measured endometrial thickness, ovarian volume, follicle number, and largest follicle size. Phenotype A was the most prevalent (53.3%), followed by phenotype D (23.8%), phenotype B (14.8%), and phenotype C (8.2%). Classic PCOS accounted for 68.0% of participants. Women with classic PCOS were significantly older than those with non-classic phenotypes. A significant association between ethnicity and phenotype was observed, with all Indian participants presenting with classic PCOS. Following infertility treatment, both groups demonstrated significant reductions in endometrial thickness, ovarian volumes, and follicle counts ($p < 0.05$). However, women with non-classic PCOS showed greater improvements in endometrial thickness and left ovarian volume. Three-dimensional transvaginal ultrasound provides a reliable method for PCOS phenotyping and follicular monitoring in Malaysian primary care infertility settings. The observed phenotype-specific and ethnicity-related patterns highlight the potential for more personalised infertility management strategies in women with PCOS.

Keywords: Follicular development; infertility; polycystic ovary syndrome; PCOS phenotype; three-dimensional transvaginal ultrasound

PP051

Feasibility and Potential Effects of High-Intensity Interval Training Versus Moderate-Intensity Continuous Training Prehabilitation in Enhance Recovery After Surgery Pathways

SURIAH AHMAD^{1,2}, NOR AZURA AZMI^{1*}, MD ALI KATIJJAHBE^{2,3}, IAN CHIK⁴, SYARIFAH NOOR NAZIHAN SAYED MASRI⁵, MUHAMMAD FARIS HAIKAL ZULKIFLI AMIN¹, NURUL NADHIRAH RAHMAT¹, MD ALI NUR AYUB^{3,4}

¹Physiotherapy Programme, Centre For Rehabilitation and Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Department of Physiotherapy, Hospital Canselor Tuanku Muhriz, 56000 Cheras, Kuala Lumpur, Malaysia

³Heart and Lung Centre, Hospital Canselor Tuanku Muhriz, 56000 Cheras, Kuala Lumpur, Malaysia

⁴Department of Surgery, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia

⁵Department of Anaesthesiology and Intensive Care, Hospital Canselor Tuanku Muhriz, 56000 Cheras, Kuala Lumpur, Malaysia

⁶Surgery Department, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia,

*Correspondence: nazura.azmi@ukm.edu.my

ABSTRACT

Exercise-based prehabilitation improves outcomes in major abdominal surgery, but optimal intensity remains unclear. High-intensity interval training (HIIT) may offer a time-efficient alternative to moderate-intensity continuous training (MICT), yet evidence on its feasibility, safety, and effects within Enhanced Recovery After Surgery (ERAS) programmes is limited. To examine the feasibility of HIIT versus MICT prehabilitation and their potential effects on maximal inspiratory pressure (MIP) and exercise capacity in patients undergoing major abdominal surgery. This prospective, double-blinded pilot randomised controlled trial recruited 23 patients from the surgical and anaesthetic clinics scheduled for major abdominal surgery at Hospital Canselor Tuanku Muhriz, Universiti Kebangsaan Malaysia. Participants were randomised to HIIT (n = 13) or MICT (n = 10) groups, receiving supervised prehabilitation for four weeks preoperatively. Assessments occurred at baseline, post-intervention, early postoperative (days 1–7), 4–6 weeks, and 3 months postoperatively. Outcomes included post-operative pulmonary complications (PPC), length of hospital stay (LOS), MIP, and 1-minute sit-to-stand test (1MSTST). Data were analysed using repeated-measures ANOVA ($p < 0.05$). All 23 participants completed the study (mean age 61.3 ± 11.9 years) with 95.7% retention and high adherence (HIIT 100%, MICT 90% with RR = 1.11, 95% CI: 0.94-1.36). No adverse events occurred. PPCs were reported in 8% of HIIT and 0% of MICT participants (RR = 0.85, 95% CI: 0.952–1.164, $p = 0.17$). LOS was significantly shorter in MICT (14.1 ± 22.4 days) than HIIT (21.4 ± 37.6 days), $p > 0.05$. No significant time $\times$ group interactions were found for MIP ($F(2.50, 52.48) = 0.58$, $p = 0.602$).

or 1MSTST ($F(3.50, 73.48) = 1.17, p = 0.33$). Between-group effects were also non-significant for MIP ($F(1,21) = 3.56, p = 0.073$) and 1MSTST ($F(1,21) = 1.29, p = 0.27$). Post-hoc analyses showed MIP declined from post-intervention to 4–6 weeks ($p = 0.038$) and improved by 3 months ($p = 0.002$). 1MSTST decreased pre-discharge ($p < 0.001$) but exceeded baseline at 3 months ($p = 0.005$). HIIT and MICT were feasible, safe, and demonstrated comparable effects on inspiratory muscle strength and exercise capacity in ERAS perioperative prehabilitation. Larger trials are warranted to confirm these preliminary findings. The protocol of this study is registered in the Australia New Zealand Clinical Trial Registry (ANZCTR) with registration number ACTRN 12625000023459.

Keywords: Enhanced Recovery After Surgery; exercise capacity; high-intensity interval training; major abdominal surgery; prehabilitation

PP052

Zerumbone Attenuates Aluminium Chloride-Induced Neurotoxicity in an In Vitro Alzheimer's Disease Model

NURUL IZZATY MOHD AZIZNUDIN¹, ASMAH HAMID^{1*}, MAZLYZAM ABD LATIF¹, NURUL FARHANA JUFRI¹, FARAH WAHIDA IBRAHIM¹, FARAH SYAHIBAH MOHD HARIRI¹, KHAIRANA HUSAIN² MUHD HANIS MD IDRIS³

¹Center For Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Office of the Dean of Faculty of Pharmacy, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

³Integrative Pharmacogenomics Institute, Universiti Teknologi MARA, 42300 Puncak Alam, Selangor, Malaysia

*Correspondence: asmah0901@ukm.edu.my

ABSTRACT

Alzheimer's disease (AD) is a progressive neurodegenerative disorder characterised by oxidative stress that leads to neuronal damage and reduced cell viability. This study evaluates the protective and therapeutic effects of zerumbone, a bioactive sesquiterpene compound isolated from the rhizomes of *Zingiber zerumbet* (wild ginger) on the H19-7/IGF-IR hippocampal cell line exposed to aluminium chloride (AlCl₃) as an in vitro AD model. Cell viability was assessed using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. Zerumbone showed low toxicity, maintaining cell viability above 70% even at high concentrations (100 µg/mL). Accordingly, the 10% inhibitory concentration (IC₁₀) and 25% inhibitory concentration (IC₂₅) were determined to be 0.7 µg/mL and 2.3 µg/mL, respectively. In contrast, AlCl₃ caused significant cytotoxicity consistent with oxidative stress-induced neuronal damage, and the 30% inhibitory concentration (IC₃₀) was chosen to simulate early-stage AD. Cell proliferation was measured using the alamarBlue assay. Two main groups were studied: a protective group, where cells were pre-treated with zerumbone before 24-hour exposure to AlCl₃, and a therapeutic group, where zerumbone treatment was applied after 24-hour AlCl₃ exposure. Treatment effects were evaluated at 2, 4 and 8 hours, while cell proliferation was measured at 24, 48 and 72 hours. In the protective group, both IC₁₀ and IC₂₅ zerumbone significantly increased cell viability at 24 and 48 hours compared to the AlCl₃ group, with the strongest effect observed after 2 hours of pre-treatment. In the therapeutic group, zerumbone also improved cell viability, with the highest effect at 24 hours and a steady increase from 2 to 8 hours of treatment. In conclusion, zerumbone shows both protective and therapeutic effects against AlCl₃-induced neurotoxicity by enhancing cell viability and proliferation. These findings suggest that zerumbone has good potential as an antioxidant agent for the prevention and treatment of AD.

Keywords: Alzheimer's disease; antioxidant; neuroprotection; proliferation; zerumbone

PP053

Understanding of Healthy Food among Employees in Klang Valley, Malaysia: A Qualitative Study

MOHD KHAIRUDDIN NOOR KHALIB¹, ZAHARA ABDUL MANAF^{2*},
ARIMI FITRI MAT LUDIN³, SUZANA SHAHAR², IDA ROSNITA ISMAIL⁴

¹*Nutrition Science Programme, Centre for Healthy Ageing & Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Dietetic Programme, Centre for Healthy Ageing & Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

³*Biomedical Science Programme, Centre for Healthy Ageing & Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

⁴*Graduate School of Business, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia*

**Correspondence: zaharamanafi@ukm.edu.my*

ABSTRACT

Understanding how individuals conceptualise healthy food is important for designing effective nutrition interventions and improving dietary behaviours. However, little is known about how working adults in Malaysia understand and interpret healthy food. This study aim to explore the understanding of healthy food among office employee in Klang Valley, Malaysia. A qualitative study was conducted among 24 office employees in Klang Valley, Malaysia. Data were collected between March and June 2025, through six triadic group discussions (n = 18) and six in-depth interviews (n = 6). Participant were recruited from public and private sectors. Data were audio recorded, transcribed verbatim, and analyzed using inductive thematic analysis with using Atlas.ti version 25. Participants generally demonstrated a good understanding of healthy food concepts. Healthy food was commonly defined as balanced diet, portion control, maintaining regular meal patterns (avoid meal skipping and late-night eating), higher intake of fruits and vegetables, and food low in sugar, salt, fat, and oil. Participants also associated healthy food with healthier cooking methods, minimally processed foods, food hygiene, and home-cooked meals. However, several misconceptions were identified. Participants frequently perceived avoiding rice, consuming coconut milk-based dishes, fried and oily local traditional foods, and choosing bread or biscuits as inherently healthy, while healthy eating was often equated with weight loss. These findings suggest that employees understanding of healthy food reflects a combination of nutrition knowledge, personal beliefs, cultural interpretations, resulting inconsistencies in dietary understanding. Office employees demonstrated a generally good understanding of healthy food concepts, however several misconceptions may influence food choices. Targeted nutrition education is needed to improve food literacy and support healthier dietary decisions among working adults.

Keywords: Food literacy; healthy food; nutrition knowledge; office employee

PP054

Histomorphometric and Macroscopic Evaluation of the Wound Healing Potential of *Eleusine indica* (Linn.) Gaertn. in a Rabbit Model

FATINI NADIAH MOHAMMAD ZAMZAMI¹, ASMAH HAMID^{1,2},
MAZLYZAM ABDUL LATIF^{1,2*}

¹Center for Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia,
50300 Kuala Lumpur, Malaysia

²Biomedical Science Program, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala
Lumpur, Malaysia

*Correspondence: mazlyzam@ukm.edu.my

ABSTRACT

Chronic and acute wound management remains a major clinical challenge, prompting increased interest in ethnobotanical therapies. *Eleusine indica* (Linn.) Gaertn., commonly known as *rumpu sambau*, has long been used in traditional medicine for its anti-inflammatory properties; however, the biological mechanisms underlying its wound-healing potential remain insufficiently understood. This study evaluates the wound-healing efficacy of *E. indica* aqueous extracts in an in vivo rabbit model through macroscopic wound contraction and microscopic tissue remodeling analysis. Full-thickness cutaneous wounds (6 mm) were created on the dorsal region of New Zealand White rabbit using a biopsy punch. Animals were divided into five groups: 1.5% *E. indica* aqueous extract, 3% *E. indica* aqueous extract, positive control (Intrasite Gel), vehicle control (Carbopol gel), and an untreated control. Wound healing progression was monitored through daily wound contraction measurements. Histopathological evaluation was performed on tissue samples collected on days 7 and 14 using Hematoxylin and Eosin (H&E) stain and Masson's Trichrome (MT) stain. Wounds treated with *E. indica* aqueous extracts demonstrated faster contraction rates compared with the control groups. H&E staining revealed enhanced re-epithelialisation in the treated groups, characterised by organised squamous epithelial layers and the presence of active fibroblasts involved in extracellular matrix formation. Masson trichrome stain further indicated dense and well-organised collagen fiber deposition within the dermal layer, suggesting improved structural remodeling and reduced inflammatory cell infiltration during the healing process. These findings indicate that *E. indica* aqueous extract supports cutaneous wound healing by stimulating fibroblast activity and promoting collagen deposition. The results provide a physiological basis for further metabolomic investigations to identify the bioactive metabolites and molecular pathways responsible for these regenerative effects.

Keywords: Collagen deposition; *Eleusine indica*; fibroblast proliferation; histopathology; wound healing

PP055

Antiproliferative Activity and Apoptosis Induction of Triphenyltin(IV) Diisopropyl Dithiocarbamate in HepG2 Hepatocellular Carcinoma Cells

AMIRUL HAIQAL MOHD ROZAIMI^{1*}, NURUL FARAHANA KAMALUDIN¹, NORMAH AWANG¹, ASMAH HAMID¹, MUHD HANIS MD IDRIS²

¹Center for Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Integrative Pharmacogenomics Institute, University Teknologi MARA, 42300 Puncak Alam, Selangor, Malaysia

*Correspondence: P159549@siswa.ukm.edu.my

ABSTRACT

Hepatocellular carcinoma (HCC), the primary form of liver cancer, remains one of the leading causes of cancer-related mortality worldwide. In Malaysia, HCC ranks among the top ten most common cancers, highlighting the urgent need for more effective chemotherapeutic agents. Organotin(IV) derivatives have attracted considerable attention due to their potent cytotoxic activity against various cancer cell lines. Among these derivatives, triphenyltin(IV) diisopropyl dithiocarbamate emerges as a promising anticancer candidate with submicromolar potency. This study aimed to evaluate the cytotoxic activity and determine the mode of cell death induced by triphenyltin(IV) diisopropyl dithiocarbamate in the human hepatocellular carcinoma cell line, HepG2. Cytotoxicity was assessed using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay following 24-hour exposure of HepG2 cells to various concentrations of the compound. The half-maximal inhibitory concentration (IC₅₀) was calculated and compared with menadione, which was used as a positive control. To explore the mode of cell death, the Annexin V-FITC/PI staining assay was performed and analysed to distinguish the percentage of apoptotic from necrotic cell populations. Triphenyltin(IV) diisopropyl dithiocarbamate demonstrated notable cytotoxicity in HepG2 cells, with an IC₅₀ of $0.78 \pm 0.08 \mu\text{M}$, showing substantially higher potency than menadione (IC₅₀ = $13.00 \pm 1.56 \mu\text{M}$). The mode of cell death analysis revealed that, at the IC₅₀ concentration, triphenyltin(IV) diisopropyl dithiocarbamate induced significant apoptosis, with 83.33% of the cell population undergoing apoptosis. These results suggest that apoptosis is the main mechanism behind the cytotoxic effects of triphenyltin(IV) diisopropyl dithiocarbamate in HepG2 cells. In conclusion, triphenyltin(IV) diisopropyl dithiocarbamate demonstrates potent antiproliferative activity against HepG2 cells and induces apoptosis as the predominant mechanism of cell death. Its superior cytotoxic efficacy compared with the positive control highlights its potential as a promising anticancer candidate for further investigation in the development of novel therapeutic strategies for hepatocellular carcinoma.

Keywords: Apoptosis; cytotoxicity; cell death; HepG2; triphenyltin

PP056

Detection of Heavy Metals and Microplastics in Selected Vegetables From Markets and Supermarkets around Klang Valley

SHAFARIATUL AKMAR ISHAK*, MOHD SHAM OTHMAN

Centre of Toxicology And Health Risk, Faculty of Health Science, Universiti Kebangsaan Malaysia, 50300
Kuala Lumpur, Malaysia

*Correspondence: sheffiyisacc@gmail.com

ABSTRACT

This study investigated the presence of microplastics and heavy metals in selected vegetables like green coral lettuce (*Lactuca sativa*), broccoli (*Brassica oleracea var. italica*), pegaga (*Centella asiatica*), and water spinach (*Ipomoea aquatica*) obtained from markets and supermarkets around Klang Valley. Microplastics were analysed using stereomicroscopy and Attenuated Total Reflectance Fourier Transform Infrared Spectroscopy (ATR-FTIR), while heavy metals were quantified using Inductively Coupled Plasma Mass Spectrometry (ICP- MS). Unwashed pegaga, green coral lettuce, and broccoli showed the highest microplastic counts (1.33 particles g⁻¹), with levels decreasing significantly after rinsing. One-way ANOVA confirmed significant differences ($p < 0.05$) in microplastic content among vegetables. For heavy metals, unwashed water spinach had the highest Pb concentration (1.12 mg/kg), while washed water spinach and pegaga recorded the highest Cu (1.63 mg/kg), Zn (11.61 mg/kg), and Ni (0.43 mg/kg). Significant differences were found for Cd, Zn, Cu, and Ni ($p < 0.05$), but not for Pb and Cr. Most heavy metal levels were within FAO/WHO limits, except Pb in unwashed pegaga and water spinach. The results highlight contamination risks based on vegetable type and rinsing status.

Keywords: Heavy metals; microplastics; UKM; vegetables

PP057

Quantifying Dose Enhancement Factor by Gold Nanoparticles (AuNPs) in 6 MV and 10 MV Photon Energies Radiotherapy: A Monte Carlo Study with GATE/Geant4

SITI HAJAR ZUBER^{1*}, SAMSON DAMILOLA OLUWAFEMI^{1,5},
MUHAMMAD FAHMI RIZAL ABDUL HADI², RAIZULNASUHA AB
RASHID³, WAN NORDIANA W ABD RAHMAN⁴, MUHAMMAD
SAFWAN AHMAD FADZIL¹, AHMAD NAZLIM YUSOFF¹, OMOLARA
VICTORIA OYELADE⁶

¹Centre for Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti
Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²School of Physics, Universiti Sains Malaysia, 11800 Penang, Malaysia

³Centre for Diagnostic Nuclear Imaging, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

⁴Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Selangor, Malaysia

⁵Department of Physics, Faculty of Science, University of Abuja P.M.B. 117 Abuja, Nigeria

⁶Department of Physics, Faculty of Science, Bingham University, P.M.B. 005 Karu, Nigeria

*Correspondence: hajarzuber@ukm.edu.my

ABSTRACT

Gold nanoparticles (AuNPs) are emerging as promising radiosensitisers because of their ability to enhance local energy deposition during radiotherapy. This study quantified dose enhancement induced by AuNPs in megavoltage photon beams and examined its dependence on photon energy and nanoparticle size using validated Monte Carlo simulations based on GATE and Geant4. A clinical linear accelerator was modeled to simulate 6 MV and 10 MV photon beams. Beam quality was optimised by adjusting incident electron energies to 6.4 MeV and 9.8 MeV, respectively. The simulated percentage depth dose (PDD) curves were validated against reference clinical data using gamma index analysis (3%/3 mm), achieving passing rates of 98.0% for 6 MV and 94.74% for 10 MV, demonstrating excellent agreement. Spherical AuNPs with diameters of 1.9, 5.0, and 15.0 nm were embedded in a homogeneous water-equivalent phantom. Absorbed dose, dose enhancement factor (DEF), and radial energy deposition profiles were evaluated to characterise nanoscale dose amplification. Results showed strong size- and energy-dependent enhancement. Under 6 MV irradiation, the maximum DEF for 1.9 nm AuNPs was 5.73, decreasing with increasing particle size. At 10 MV, ultrasmall AuNPs exhibited even greater enhancement, with a peak DEF of 16.13 for 1.9 nm particles, whereas 15.0 nm particles showed minimal enhancement (DEF 1.13). Radial dose analysis revealed highly localised energy deposition within 10–20 nm of the nanoparticle surface, primarily driven by secondary electron emission. Larger nanoparticles demonstrated reduced enhancement due to increased self-absorption, while smaller particles produced higher surface doses and steeper dose gradients. Overall, AuNP-mediated dose enhancement in megavoltage radiotherapy depends strongly on nanoparticle size and photon energy. Ultrasmall AuNPs (<5 nm) provide superior

radiosensitisation, particularly at 10 MV, highlighting the importance of nanoscale optimisation and confirming the robustness of Monte Carlo modeling for nanoparticle-based radiotherapy research.

Keywords: AuNPs; dose enhancement factor; GATE/Geant4; Monte Carlo dosimetry; nanoscale radiosensitisation

PP058

Elucidating The Efficacy of a Novel Supplement Containing Standardised *Cosmos Caudatus* Extract for the Mental Well-Being in the Ageing Population with Mild Cognitive Impairment: Study Protocol for a Randomised Controlled Trial

SHARIFAH MIZA HASMADI¹, SUZANA SHAHAR^{1*}, YOU YEE XING¹,
HAMIDUN BUNAWAN², NOR FADILAH RAJAB¹, NIK RUZYANEI NIK
JAAFAR³, MAZLYFARINA MOHAMAD⁴

¹Center for Healthy Aging and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Institute of Systems Biology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

³Department of Psychiatry, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia

⁴Centre for Diagnostic, Therapeutic & Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

*Correspondence: suzana.shahar@ukm.edu.my

ABSTRACT

Introduction: The global prevalence of mild cognitive impairment (MCI) among adults aged 50 years and above is rising sharply, linked to a higher progression rate in developing dementia. *Cosmos caudatus* (CC), an indigenous herb rich in flavonoids such as quercetin and quercitrin, has demonstrated neuroprotective and cognitive-enhancing potential. This study employs a randomised, double-blind, controlled trial design with three study arms aiming to explore the effectiveness of a plant-based supplement containing CC extract on cognitive function, mood status, blood biochemical profile, and cognitive-related biomarkers. **Method/analysis:** Participants with Mild Cognitive Impairment (MCI) aged 50 years and above will be invited to participate in this study and thereafter will be randomly assigned to one of three groups: 250 mg CC, 500 mg CC, or a placebo for six months. The primary outcomes include changes in cognitive function and mood status. The secondary outcomes include changes in the blood biochemical profile and cognitive-related biomarkers supported by functional magnetic resonance imaging (fMRI). **Ethics and dissemination:** This study protocol is currently undergoing ethical review and will be approved by the Medical Research Ethics Committee of the National University of Malaysia. Study findings will be disseminated through peer-reviewed conference presentations and open-access publications.

Keywords: Biomarkers; cognitive function; *Cosmos caudatus*; fMRI; mild cognitive impairment

PP059

Neuroprotective and Therapeutic Potential of Zerumbone against Aluminium-Induced Neurotoxicity: A Behavioural Study in an Alzheimer's Rat Model

FARAH SYAHIBAH MOHD HARIRI¹, ASMAH HAMID^{1*}, FATINI NADIAH MOHAMAD ZAMZAMI¹, NURUL FARHANA JUFRI¹, MAZLYZAM ABDUL LATIF¹, FARAH WAHIDA IBRAHIM¹, KHAIRANA HUSAIN²

¹Center for Toxicology & Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Centre for Drug and Herbal Development, Faculty of Pharmacy, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

*Correspondence: asmah0901@ukm.edu.my

ABSTRACT

Aluminum accumulation in the brain is one of the significant environmental risk factors for Alzheimer's disease (AD), leading to oxidative damage and cognitive decline. Aluminum may act as a cross-linker, encouraging amyloid-beta proteins to clump together into the plaques that are the hallmark of Alzheimer's. Aluminum also contributes to the hyperphosphorylation of tau proteins, leading to the neurofibrillary tangles that disrupt nutrient transport within brain cells. Zerumbone (ZER), a bioactive sesquiterpene from the *Zingiber zerumbet* plant, has shown potent anti-inflammatory and antioxidant properties. Therefore, the objective of this study is to evaluate the neuroprotective and therapeutic effects of ZER on aluminum-induced memory impairment and behavioral changes in Wistar rats. Eighty male Wistar rats were co-administered aluminum chloride (AlCl₃) (25 mg/kg) and Zerumbone (10, 20, and 30 mg/kg) via oral gavage for 42 days (neuroprotective) and 56 days (therapeutic). Cognitive function was assessed using the Open Field Test (OFT), Y-Maze and Novel Object Recognition (NOR) test. Results indicate that aluminum exposure significantly impaired spatial working memory in the Y-Maze test and reduced the Discrimination Index in the NOR test. Treatment with ZER effectively rescued these cognitive deficits without affecting locomotor activity in the OFT. Statistical analysis confirmed a significant restoration of recognition memory in ZER-treated cohorts compared to the aluminum-only group. In conclusion, ZER demonstrates robust neuroprotective and therapeutic efficacy against aluminum-induced neurotoxicity. These findings suggest ZER as a potential candidate for mitigating heavy-metal-induced cognitive impairment in Alzheimer's disease.

Keywords: Aluminium chloride; neuroprotection; novel object recognition; spatial memory; zerumbone

PP060

Boosting Post-Stroke Recovery: Feasibility and Potential Benefits of Soft Robotic-Assisted Hand Training with Standard Therapy in Community-Dwelling Survivors

NIK INTAN SHAKINAH RAMLI, NOR AZURA AZMI*, MOHD AZZUAN AHMAD, NOR AZLIN MOHD NORDIN, LEE SHU JUIN

Physiotherapy Programme, Center for Rehabilitation and Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

**Correspondence: nazura.azmi@ukm.edu.my*

ABSTRACT

Soft robotic hand devices have emerged as an adjunct to conventional physiotherapy to enhance motor recovery. However, evidence on their feasibility, safety, and potential effects remains limited, particularly among Malaysian community-dwelling stroke survivors. This study aimed to evaluate the feasibility and potential effects of soft robotic-assisted hand training combined with standard therapy (SRAHT-ST) on upper limb function, lung function, and quality of life in post-stroke patients. This prospective single-blinded pilot randomised controlled trial recruited 10 community-dwelling stroke survivors from the Klang Valley. Participants were randomly allocated to SRAHT-ST ($n = 5$) or standard therapy (ST, $n = 5$). Both groups received therapist-supervised sessions over 8 weeks. Feasibility and safety were assessed, and outcomes included the Action Research Arm Test (ARAT), Box and Block Test (BBT), Fugl-Meyer motor (FMM) and sensory (FMS) assessments, lung function, and quality of life, measured at baseline, 4 weeks, and 8 weeks. Ethical approval was obtained from Universiti Kebangsaan Malaysia (JEP-2024–261), and the study was registered with the Australian New Zealand Clinical Trials Registry (ACTRN12624000923561). All participants completed the study (mean age 61.3 ± 11.9 years) with 81.3% retention and high adherence. No adverse events were reported. Two-way mixed ANOVA showed significant improvements over time across all outcomes ($p < 0.001$). Significant time $\times$ group interaction effects were observed for ARAT, FMM, FMS, BBT, handgrip strength and lung function ($p < 0.01$). Post-hoc analyses demonstrated greater improvements in the SRAHT-ST group from baseline to 4 and 8 weeks ($p < 0.05$). The SRAHT-ST protocol was feasible, safe and showed potential benefits for upper limb and lung function. Findings preliminarily support soft robotic-assisted training as an adjunct to rehabilitation and guide protocol refinement. Given the small sample size, results should be interpreted cautiously and adequately powered trial is warranted.

Keywords: Community-dwelling; lung function; soft robotic-assisted hand training; stroke survivors; upper-limb function

PP061

SMARTCAFF-i™: A Web-Based Innovation for Monitoring Caffeine Intake among Athletes

SITTI JUNAINA MUSA¹, AHMAD ROHI GHAZALI², NOR FARAH MOHAMAD FAUZI^{1,5}, MOHD IZHAM MOHAMAD³, ABDUL HADI ABD RAHMAN⁴, NIK SHANITA SAFII^{1,5*}

¹Center for Community Health Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Center for Toxicology & Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

³Sports Nutrition Centre, National Sport Institute, Kuala Lumpur, Malaysia

⁴Center for Artificial Intelligence Technology, Faculty of Information Sciences and Technology, University Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

⁵Obesity-UKM Research Group, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

*Correspondence: nikshanita@ukm.edu.my

ABSTRACT

Caffeine is widely consumed by athletes to enhance physical performance, endurance and alertness. However, excessive intake may lead to adverse physiological effects such as cardiovascular disturbances, sleep disruption and anxiety. In Malaysia, limited information on the local caffeine content data together with the lack of practical tools to monitor caffeine consumption, poses challenges for athletes and active individuals in managing their intake safely. Addressing this gap is important for promoting safer caffeine consumption behaviours among athletes and physically active individuals. This study aimed to develop and evaluate SmartCaff-i™, a web-based application designed to assist athletes in monitoring their caffeine intake. The development of SmartCaff-i™ was conducted through four research phases. Phase 1 (needs analysis) explored caffeine consumption patterns and identified commonly consumed caffeinated beverages among athletes. Phase 2 (laboratory analysis) determined the caffeine content of selected commonly consumed beverages to establish a reliable local caffeine database. Phase 3 (application development) involved the design and development of the SmartCaff-i™ web platform, integrating the caffeine database into an interactive monitoring system, followed by prototype validation. Phase 4 (usability evaluation) assessed the usability and practicality of the application as a personal monitoring tool for caffeine intake. The SmartCaff-i™ application enables users to record and monitor daily caffeine intake through features such as a caffeine diary, intake tracking and educational caffeine information on caffeine consumption. By integrating locally generated research data with digital health technology, the application supports informed decision-making and promotes safer caffeine consumption. SmartCaff-i™ bridges the gap between laboratory research and digital health while demonstrates how research-based innovation can contribute to improving caffeine intake management among athletes and physically

active individuals. The development of this platform highlights the importance of interdisciplinary research collaboration and technological innovation in creating practical digital health solutions for monitoring dietary intake.

Keywords: Athlete; caffeine intake monitoring; community health; digital health innovation; SmartCaff-i™

PP062

Temporal Dysregulation of β -Amyloid Clearance in the Hippocampal Microenvironment of an Alzheimer's Disease Rat Model

HILAL HAIQAL FAIZAL¹, FARAH WAHIDA IBRAHIM^{1*}, AMIRUL HAFIZ AHMAD ABDULLAH¹, NOR FADILAH RAJAB², ASMAH HAMID¹, NURUL FARHANA JUFRI¹, SITI FATHIAH MASRE¹

¹Center for Toxicology and Health Risk Studies, Faculty of Health Sciences, National University of Malaysia, 50300 Kuala Lumpur, Malaysia

²Center for Healthy Ageing & Wellness, Faculty of Health Sciences, National University of Malaysia, 50300 Kuala Lumpur, Malaysia

*Correspondence: farahwahida@ukm.edu.my

ABSTRACT

Alzheimer's disease (AD) is characterised by the accumulation of β -amyloid ($A\beta$), with impaired clearance increasingly recognised as a key contributor to disease progression. However, the temporal dynamics of $A\beta$ clearance within the hippocampal microenvironment remain inadequately understood. This study examined transcriptomic and biochemical changes in rats exposed to aluminium chloride ($AlCl_3$, 200 mg/kg/day) at D14 (acute), D28 (subacute), and D42 (subchronic) timepoints, each comprising control and induced groups. RNA sequencing ($n = 12$) generated high-quality data (average 7.0 Gb/sample, 98.2% mapping), identifying 18,570 expressed genes, while soluble $A\beta_{40}$ levels were quantified using ELISA. At D14, soluble $A\beta_{40}$ levels were markedly elevated, accompanied by increased expression of amyloidogenic genes (*Bace1*, *Psen1*) and clearance-related genes (*Ide*, *Lrp1*, *Scarb1*, *Aqp4*, *Ccl2*). This pattern suggests an early compensatory response, primarily mediated by intracellular degradation and transport mechanisms. Microglial activation (*Trem2*) remained limited, while reduced *Apoe* and *Mme* expression may indicate incomplete engagement of lipid-associated and extracellular clearance pathways. By D28, soluble $A\beta_{40}$ levels had normalised alongside reduced expression of both production- and clearance-related genes, although *Aqp4* remained elevated. Increased expression of *Cxcl10*, *Trem2*, *Gfap*, and *Anxa3*, together with modest *Mme* levels, is consistent with enhanced inflammatory signalling and microglial activation, occurring alongside declining clearance capacity. At D42, sustained upregulation of *Trem2* and *Apoe*, coupled with increased *Gfap* and reduced *Map2* expression, points towards persistent glial activation with emerging neuronal impairment. Concurrent reductions in *Aqp4* and *Lrp1*, alongside partial *Ide* recovery and modest *Mme* expression, may reflect disruption of glymphatic and transport-mediated clearance pathways. Reduced *Cxcl10* with sustained *Ccl2* suggests a shift towards a more chronic inflammatory state. In conclusion, $A\beta$ clearance exhibits a stage-dependent transition from early compensatory responses to progressive dysregulation, with altered $A\beta$ dynamics at later stages.

Keywords: Aluminium-induced rat model; Alzheimer's disease; astrocytes and microglia; β -amyloid clearance

PP063

Exercise Modulates Lymphocyte Populations in Aluminium-induced Immunotoxicity in Wistar Rats

NOR MALIA ABD WARIF^{1,2*}, VOON ZI XIAN¹, NUR AISYAH ANUAR³,
SITI KHADIJAH ISMAIL¹, ARIMI FITRI MAT LUDIN^{1,3}

¹*Biomedical Sciences Program, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Centre for Toxicology & Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

³*Centre for Healthy Aging & Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

*Correspondence: malia.warif@ukm.edu.my

ABSTRACT

Aluminium (Al) is a common environmental toxicant that causes oxidative stress and accumulates in tissues, resulting in the suppression of immune function. Although Al is known for neurotoxicity, its effects on the peripheral immune system are poorly characterised. Exercise has been suggested as a nonpharmacological therapy that might alleviate chemical toxicities, but its protective mechanisms against Al-induced immunotoxicity in lymphoid organs remain unknown. The aim of this study is to evaluate the protective effects of a 70-day structured high-intensity interval training (HIIT) treadmill regimen against chronic aluminium-induced immunotoxicity in the spleen, mesenteric lymph nodes (MLN), and cervical lymph nodes (CLN) of Wistar rats. Twenty-four male Wistar rats were randomised into four groups: Sedentary Control (DES), Exercise Control (DEH), Aluminium Sedentary (ES), and Aluminium Exercise (EH). Aluminium groups received AlCl₃ (200 mg/kg/day) orally for 70 days. Treadmill exercise began on day 28 for DEH and EH and continued until day 70. At the study's end, the rats were sacrificed and the spleen, MLN, and CLN were harvested for flow cytometry analysis. Results demonstrated that the DEH group significantly reduced the percentage of splenic CD3+CD8+ T lymphocytes ($p < 0.05$). Conversely, the ES group led to a notable decrease in CD3+CD4+FoxP3+ regulatory T cells within both mesenteric and cervical lymph nodes ($p < 0.05$). A similar reduction in this regulatory population was also observed in the mesenteric lymph nodes of the DEH group ($p < 0.05$). Our findings demonstrate that aluminium toxicity causes substantial immune dysfunction, which can be effectively countered through physical exercise. This intervention specifically preserves immune integrity by stabilising immunoregulatory mechanisms within the gut-associated lymphoid tissues.

Keywords: Aluminium; exercise; immunotoxicity; lymphocytes; wistar rats

PP064

Microbiological Safety of Street-Vended Martabak in Kuala Lumpur

NUR AFIFAH MOHAMAD AYOP, SITI SHAHARA ZULFAKAR*

Environmental Health and Industrial Safety Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

**Correspondence: sitishahara.zulfakar@ukm.edu.my*

ABSTRACT

Street food is widely consumed for its affordability, yet it remains a public health concern due to food safety risks linked to poor hygiene. This study aimed to determine bacterial contamination levels in martabak and its side dishes (pickled onion and curry) sold by street vendors in Kuala Lumpur. This study also assesses vendor hygiene status and examines the association between premise hygiene score and the bacterial contamination level. A cross-sectional study was conducted involving 20 samples of martabak and its side dishes from 20 vendors around Kuala Lumpur. Samples were analysed for total viable count (TVC), coliforms, *Escherichia coli*, *Staphylococcus aureus* and *Salmonella* spp. The vendor's premise hygiene score was evaluated using a validated 25-item observational checklist that includes personal hygiene, utensils and storage, handwashing facilities, surrounding area, and pest control components. Findings revealed that only one martabak sample was found to have the presence of bacteria with the TVC reading of 2.957 log CFU/g. In contrast, pickled onion showed a higher frequency of contamination, with the presence of bacteria detected in six samples (average TVC: 3.884 ± 1.087 log CFU/g). Curry samples demonstrated the most critical microbiological profile, with concurrent contamination across four parameters. Hygiene assessments classified 85% of vendors as having moderate hygiene levels, while only 15% were categorised as clean. Spearman's correlation analysis showed a significant moderate negative association between hygiene scores and TVC levels of the side dishes ($r_s = -0.411$, $p = 0.035$). In conclusion, while freshly cooked martabak was generally safe, the accompanying side dishes posed significant risks due to inadequate hygiene. Strengthening sanitation facilities, improving handling practices, and providing continuous training for vendors are essential measures to reduce foodborne disease risks among consumers in Kuala Lumpur.

Keywords: Bacterial contamination; food safety; hygiene practice; martabak; street food

PP065

Development of Fast-Dissolving *Piper betle* Herbal Oral Strips with Antibacterial Activity for Halitosis Management

NORAZALINA^{1*}, AFIQ AZIL¹, AHMAD FARID¹, NURUL AKMARINA²

¹School of Pharmacy, KPJ Healthcare University, 71800 Kota Seriemas, Nilai, Negeri Sembilan, Malaysia

²Department of Chemical Engineering, Faculty of Chemical and Process Engineering Technology, Universiti Malaysia Pahang Al-Sultan Abdullah, 26300 Gambang, Pahang, Malaysia

*Correspondence: norazalina@kpju.edu.my

ABSTRACT

Halitosis is a common oral health condition primarily caused by bacterial activity in the oral cavity, particularly *Streptococcus mutans* and *Staphylococcus aureus*, which produce volatile sulphur compounds responsible for unpleasant breath odour. Conventional oral hygiene products such as mouthwashes and chewing gums often contain synthetic chemicals that may cause irritation and provide only temporary relief. Therefore, this study aimed to develop a natural and portable oral care product by formulating fast-dissolving herbal oral strips containing *Piper betle* L. leaf extract. Fresh betel leaves were subjected to ethanol extraction using the cold maceration technique to obtain the crude extract. Phytochemical screening was conducted to identify bioactive compounds, followed by antibacterial evaluation using minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) assays against *S. mutans* and *S. aureus*. The extract was then incorporated into an oral strip formulation using the solvent casting method with suitable film-forming polymers, plasticisers, sweeteners, and flavouring agents. The results revealed the presence of bioactive phytochemicals including flavonoids, phenols, and terpenoids, which are known for their antimicrobial properties. Antibacterial testing demonstrated effective inhibition of oral pathogens with MIC and MBC values indicating strong antibacterial potential, and inhibition zones reaching up to 24 mm. The developed oral strips exhibited desirable pharmaceutical characteristics, including thin, flexible films, pH values ranging from 6.7 to 6.9, and rapid disintegration within 30 seconds when placed on the tongue. These findings indicate that *Piper betle*-based oral strips represent a promising natural therapeutic approach for reducing oral microbial load and managing halitosis. This innovation integrates traditional herbal medicine with modern drug delivery technology, offering a safe, convenient, and user-friendly oral hygiene product with potential applications in both pharmaceutical and personal healthcare sectors.

Keywords: Antibacterial activity; fast-dissolving oral thin films; *Piper betle* L.; halitosis; solvent casting method

PP066

Feasibility of Soft Robotic–Assisted Hand Training Combined with Standard Therapy in Community-Dwelling Stroke Survivors: A Pilot Randomised Controlled Trial

NIK INTAN SHAKINAH RAMLI, NOR AZURA AZMI*, MOHD AZZUAN AHMAD, NOR AZLIN MOHD NORDIN, LEE SHU JUIN

Physiotherapy Programme, Center For Rehabilitation And Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

**Correspondence: nazura.azmi@ukm.edu.my*

ABSTRACT

Soft robotic hand devices have emerged as a potential adjunct to conventional physiotherapy for post-stroke upper limb rehabilitation. However, evidence regarding their feasibility, safety, and preliminary clinical signals remains limited, particularly among Malaysian community-dwelling stroke survivors. This pilot study evaluated the feasibility, safety, and preliminary clinical signals of a soft robotic-assisted hand training protocol combined with standard therapy (SRAHT-ST) on upper limb function, lung function, and quality of life in post-stroke patients. In this prospective single-blinded pilot randomised controlled trial, 10 community-dwelling stroke survivors within the Klang Valley were randomised to either SRAHT-ST (n = 5) or standard therapy alone (ST, n = 5). Both groups received 24 therapist-supervised sessions over 8 weeks. Primary feasibility outcomes were retention rates, intervention adherence, and safety. Secondary outcomes included the Action Research Arm Test, Box and Block Test, Fugl-Meyer motor, Fugl-Meyer sensory, lung function, and quality of life (EuroQol), which were measured at baseline, 4 weeks, and 8 weeks. Exploratory two-way mixed ANOVA was used to examine trends in changes over time and between groups. Ten participants completed the intervention (mean age 61.3 ± 11.9 years), with 81.3% retention, high adherence, and no intervention-related adverse events. Exploratory analyses showed improvements over time across outcomes ($p < 0.001$) with trends toward greater improvements in upper limb outcomes and lung function in the SRAHT-ST group versus ST (time $\times$ group interaction, $p < 0.05$). Findings should be interpreted as preliminary estimates only. The SRAHT-ST protocol was feasible, safe, and acceptable, supporting progression to a definitive trial. Exploratory findings suggest potential signals of benefit for upper limb and pulmonary function and inform future trial design. No efficacy conclusions can be drawn due to the pilot nature and small sample size.

Keywords: Feasibility; post-stroke rehabilitation; soft robotic hand; upper limb function; upper limb recovery

PP067

Investigation of Personalised Post-Reconstruction Positron Range Correction in ^{68}Ga Positron Emission Tomography Imaging

RUKIAH A LATIFF^{1*}, STEVE MEIKLE^{2,3}, PETER KENCH²

¹Centre of Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Discipline of Medical Imaging Science, Sydney School of Health Sciences, Faculty of Medicine and Health, The University of Sydney, New South Wales 2006, Australia

³The Brain and Mind Centre, The University of Sydney, 94 Mallett St, Camperdown, New South Wales 2050, Australia

*Correspondence: rukiah@ukm.edu.my

ABSTRACT

Positron range blurring inherently limits the spatial resolution of Gallium-68 (^{68}Ga) Positron Emission Tomography (PET) imaging. This study aims to develop an iterative post-reconstructed positron range correction (PRC) algorithm and evaluate its impact on the quality of clinical PET images. An iterative post-reconstruction PRC algorithm was developed by comparing measured clinical PET images with simulated annihilation images generated via GATE Monte Carlo simulations, utilising patient-specific CT-derived attenuation and density maps. The method was evaluated retrospectively on 17 patients undergoing ^{68}Ga -DOTATATE PET scans. Two experienced, blinded nuclear medicine physicians reviewed standard PET (PET_0) and corrected (PRC_5) images side-by-side, assessing overall image quality, artifacts, and lesion preferences. Quantitative analyses evaluated image noise, standardised uptake value (SUV), contrast, and signal-to-noise ratio (SNR). Quantitatively, the PRC significantly improved lesion contrast ($p < 0.05$) and SNR ($p < 0.05$), and yielded increased radiotracer uptake ($p < 0.05$). The algorithm successfully preserved the normal physiological distribution of ^{68}Ga in the body, with only a marginal noise increment of approximately 2%. Despite quantitative enhancements, qualitative reader assessments showed a preference for uncorrected PET_0 images. PET_0 images achieved 94.1% agreement for “Excellent” overall quality, whereas PRC_5 achieved 58.8% agreement, typically categorised as “Satisfactory”. Reduced preference for PRC_5 was attributed to artifacts induced during the deblurring process, predominantly in the abdominal region (e.g., liver, bowel, spleen), which were caused by spatial discrepancies between the PET and CT attenuation maps due to respiratory and peristaltic motion. Personalised post-reconstruction PRC successfully improves quantitative metrics such as lesion contrast and SNR in ^{68}Ga imaging. However, clinical preference is negatively impacted by motion-induced artifacts. In artifact-free images, PRC is the preferred modality, warranting further research to mitigate attenuation map discrepancies and fully translate quantitative gains into clinical diagnostic confidence.

Keywords: Gallium-68; GATE Monte Carlo simulation; iterative post-reconstruction positron range correction

PP068

Preliminary Results of a Scoping Review on Teaching Older Adults How to Land Safely: Movement-Based Strategies to Reduce Fall-Related Injury Risk

NUR AIMUNI ABU HASSAN, SHAWN SOH LENG HSIEN, NORMALA MESBAH, DEVINDER KAUR AJIT SINGH*

Centre for Healthy Ageing & Wellness, Faculty of Health Science, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

**Correspondence: devinder@ukm.edu.my*

ABSTRACT

Falls are a leading cause of injury, disability, and loss of independence among older adults. Although conventional fall prevention programmes aim to reduce fall incidence, many falls remain unavoidable. Teaching safe falling or movement-based landing strategies may help reduce injury severity when falls occur; however, the current evidence base remains fragmented. This scoping review aims to map the existing literature on interventions that teach safe falling or landing strategies to older adults and examine their potential role in reducing fall-related injury risk. A scoping review was conducted in accordance with PRISMA-ScR principles. Electronic databases including PubMed, Scopus, Web of Science, PEDro and the Cochrane Library were searched. Studies involving adults aged ≥ 60 years and interventions explicitly teaching safe falling, fall-landing, or fall recovery strategies were considered. Data were extracted and synthesised descriptively. A total of 21 relevant studies were identified, including randomised controlled trials, feasibility studies, qualitative studies, protocols, and laboratory-based biomechanical studies. Common approaches included judo-inspired programmes, martial arts-based falling techniques, tuck-and-roll strategies, motor analogy training, and fall arrest programmes. Reported outcomes included reduced hip impact force, improved balance and strength, enhanced fall technique competence, increased confidence, and reduced fear of falling. However, many studies involved small samples, healthy participants, or simulated laboratory falls. Preliminary findings suggest that movement-based safe landing strategies are feasible and may reduce fall-related injury risk among older adults. Further high-quality trials involving real-world populations and long-term outcomes are needed to support clinical implementation.

Keywords: Fall-related injuries; falls prevention; injury prevention older adults; safe landing strategies

PP069

Knowledge, Awareness and Practice of Breast Self-Examination among Female Students at a Private Healthcare University in Nilai

NURUL MAWADDAH MOHAMMAD*, KHAIRUNAJWA MUHAMMAD
KAMALUDDIN

Department of Medical Imaging, School of Health Sciences. KPJ Healthcare University, 71800 Nilai, Negeri Sembilan, Malaysia

**Correspondence: ucn.mawaddah@kpju.edu.my*

ABSTRACT

Breast cancer remains a major global public health concern, with early detection playing a critical role in improving survival outcomes. Breast self-examination (BSE) is a simple, low-cost strategy that promotes breast awareness, particularly among young women. This study aimed to assess the association between demographic characteristics, knowledge, and awareness with BSE practice among female students at a private healthcare university in Nilai. A cross-sectional study was conducted among 351 female students using a structured online questionnaire assessing socio-demographic characteristics and levels of knowledge, awareness, and practice of BSE. Descriptive statistics and multiple logistic regression analyses were performed. The findings indicated that 56.3% and 55.8% of respondents had poor knowledge and awareness, respectively, while 54.6% demonstrated poor BSE practice. Although 82.3% had heard of BSE, only 20.5% performed it monthly as recommended. Common barriers included a lack of knowledge (48.9%), low perceived risk (45.7%), and forgetfulness (42.0%). Significant associations were observed between knowledge and practice ($p < 0.001$) and between awareness and practice ($p < 0.001$), whereas no significant association was found between demographic factors and practice. Notably, 85.2% of respondents expressed a willingness to attend future educational programs. In conclusion, despite moderate awareness of BSE, substantial gaps exist in knowledge and consistent practice. These findings highlight the need for structured, skills-based educational interventions to enhance competency and promote regular BSE practice, thereby strengthening early detection efforts and contributing to improved breast health outcomes.

Keywords: Awareness and practice; breast self-examination; breast cancer; early detection; knowledge; university students

PP071

In-Hospital Mortality of Infective Endocarditis in Malaysia: A Systematic Review and Meta-Analysis

MUHD NAJMI HAKIM ABDUL RANI¹, SUHAINI KADIMAN⁴, ZALEHA
MD ISA², AFIFAH MOHAMED³, TATY ANNA KAMARUDIN^{1*}

¹Department of Anatomy, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur
Malaysia

²Department of Public Health Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras,
Kuala Lumpur, Malaysia

³Centre for Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti
Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

⁴Department of Anesthesia & Clinical Research, National Heart Institute, Kuala Lumpur, Malaysia

*Correspondence: tatykamarudin@ukm.edu.my

ABSTRACT

Infective endocarditis (IE) remains a severe infection associated with substantial early mortality. However, in Malaysia, reported in-hospital mortality rates vary widely across single-centre retrospective studies (13–44.4%), and no prior systematic review has provided a pooled national estimate. This lack of a comprehensive national synthesis limits comparison with international data and constrains evidence-based clinical and policy decision-making. This systematic review and meta-analysis aimed to estimate the pooled in-hospital mortality rate of IE in Malaysia. It also sought to contextualise outcomes across healthcare settings, time periods, and prognostic factors, with the goal of establishing the first national benchmark to inform clinical practice, registry development, and health-system planning. Conducted in accordance with the PRISMA 2020 statement, a systematic search of three electronic databases (PubMed, Google Scholar and SciSpace) was performed. Eligible studies were observational studies reporting in-hospital mortality among adult patients (18 years) diagnosed with IE using the modified Duke criteria in Malaysian healthcare institutions and published in English. Of 255 records identified, seven unique Malaysian studies (eight mortality estimates comprising 382 adult patients, 2005–2023) were included. Using a random-effects model, the pooled in-hospital mortality rate was 33.2% (95% CI: 27.8–39.1%), with low between-study heterogeneity ($I^2 = 13.3\%$, $p = 0.326$). Mortality was higher in tertiary referral centres (30.8–44.4%) compared to a secondary-care hospital (13.0%). *Staphylococcus aureus* infection, heart failure, hemodynamic instability, renal dysfunction, and the absence of surgical intervention were identified as adverse prognostic factors. Overall, in-hospital mortality in Malaysia remains high, systemic, and exceeds international benchmarks. These findings support the urgent need to strengthen clinical pathways, promote earlier diagnosis, establish multidisciplinary endocarditis teams, and develop a national IE registry to guide health-system policy and enable meaningful international comparison.

Keywords: Infective endocarditis; in-hospital mortality; Malaysia; meta-analysis; systematic review

PP074

Lysosomal Impairment Drives Vacuolation but Not Significant Amyloid- β 40 Accumulation in Human Brain Endothelial Cells

NURUL FARHANA JUFRI^{1,2*}, NOR AZMA HANNAH MAZLAN², MOHD HAMZAH MOHD NASIR³, FARAH WAHIDA IBRAHIM^{1,2}, ASMAH HAMID^{1,2}, IFFAH NADIAH LAILI¹

¹Center for Toxicology & Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia,

²Biomedical Science Program, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300, Kuala Lumpur, Malaysia.

³Department of Biotechnology, Kulliyyah of Science, International Islamic University of Malaysia, Kuantan Campus, Jalan Sultan Ahmad Shah, Bandar Indera Mahkota, 25200 Kuantan, Pahang Darul Makmur, Malaysia.

*Correspondence: nurulfarhana@ukm.edu.my

ABSTRACT

Alzheimer's disease (AD) is characterised by progressive cognitive decline and is strongly associated with the accumulation of amyloid-beta ($A\beta$) peptides. While lysosomal dysfunction has been implicated in increased $A\beta$ 42 accumulation, its effects on $A\beta$ 40 dynamics and cytosolic vacuolation in brain endothelial cells remain unclear despite its significant role in forming $A\beta$ deposits and importance in the $A\beta$ 42/40 ratio. Thus, this study investigated the impact of chloroquine (CQ)-induced lysosomal dysfunction on cytosolic vacuolation and intracellular and extracellular $A\beta$ 40 levels in human brain endothelial cells (HBEC-5i). HBEC-5i cells were treated with 70.5 μ M CQ for 24 hours to induce lysosomal impairment. Morphological changes and cytosolic vacuolation were evaluated by inverted microscope and ImageJ analysis, while $A\beta$ 40 levels in cell lysates and supernatants were quantified using ELISA. CQ-treated cells exhibited significant morphological alterations, including cell shrinkage and prominent cytosolic vacuole formation. Quantitative analysis revealed a significant increase in vacuolation density and vacuole diameter compared to untreated controls ($p < 0.05$), indicating disrupted autophagy-lysosomal pathway function. Despite these structural changes, intracellular $A\beta$ 40 levels showed a slight, non-significant decrease, while extracellular $A\beta$ 40 levels demonstrated a non-significant increase following CQ treatment. These findings suggest that $A\beta$ 40 may be preferentially trafficked out of endothelial cells, potentially via transcytosis pathways, even under lysosomal dysfunction conditions. In conclusion, CQ-induced lysosomal impairment significantly alters endothelial cell morphology and promotes cytosolic vacuolation but does not markedly affect $A\beta$ 40 levels. The results highlight a possible differential handling of $A\beta$ 40 compared to $A\beta$ 42 and underscore the importance of endothelial transport mechanisms in AD pathology. Further studies are needed to elucidate the interplay between lysosomal dysfunction, $A\beta$ trafficking, and blood-brain barrier integrity.

Keywords: Beta amyloid; chloroquine; endothelial cells; lysosome; vacuoles

PP075

Assessing the Malaysia Mandarin Pediatric Speech Perception Semi-automated Test: Content Validity and User Confidence among Audiology Students and Professionals

FOONG YEN CHONG^{1,2*}, WAN NUR AFIZA SYAFIRA WAN MUHAMAD ZIN², TIAN KAR QUAR^{1,2}, CILA UMAT^{1,2}

¹*Centre for Rehabilitation and Special Needs, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Audiology Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

**Correspondence: foongyen.chong@ukm.edu.my*

ABSTRACT

This cross-sectional study evaluated the content validity, confidence, and perception ratings of the MyMaPS semi-automated test application and manual among student clinicians and audiologists. Validation occurred in two phases. First, six audiologists evaluated the adapted Content Validity Index (CVI) and the confidence and perception questionnaires based on relevance, simplicity, clarity and ambiguity on a 4-point scale. In the second phase, 29 student clinicians who had received clinical training for at least 1 semester, and three audiologists were recruited from the Audiology Program, Universiti Kebangsaan Malaysia (UKM). All participants answered the pre-training confidence and perception questionnaire and underwent an online briefing session, followed by face-to-face training sessions. The CVI Semi-MyMaPS and the post-training confidence and perception questionnaire were administered after the training. The results showed that the two questionnaires received good face validity ratings (scores of > 3). The MyMaPS semi-automated test application and test manual met the satisfactory level of CVI (scores of > 0.83). Wilcoxon signed-rank tests indicated that the post-training confidence and perception ratings towards the MyMaPS semi-automated test manual ($T = 351$, $z = -4.47$, and $p < 0.001$) and test application ($T = 300$, $z = -4.31$, and $p < 0.001$) were significantly higher compared to pre-training scores. The results showed that the participants were confident in conducting the MyMaPS test and found it very helpful to use in a clinical setting. These results suggested that the content of the MyMaPS semi-automated test manual and test application are valid to be used among audiology students and audiologists in Klinik Audiologi & Sains Pertuturan (KASP), UKM.

Keywords: Confidence and perception ratings; content validity; mandarin; speech perception

PP076

Satisfaction with Mobility Assistive Technology and Quality of Life among Lower Limb Amputees: A Preliminary Analysis

JIA TIAN LOO, SIAW CHUI CHAI*

Centre for Rehabilitation and Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

**Correspondence: sc.chai@ukm.edu.my*

ABSTRACT

Lower limb amputations are profound and life-altering events. Many lower limb amputees face limited access to mobility devices and report dissatisfaction, probably affecting their quality of life (QOL). Yet, research combining these factors remains lacking. This cross-sectional study examined the relationship between satisfaction with mobility assistive technology and QOL among lower-limb amputees. This study collected data using a self-developed demographic questionnaire, the Quebec User Evaluation of Satisfaction with Assistive Technology (QUEST 2.0), and the World Health Organisation Quality of Life Brief Version (WHOQOL-BREF) at the PERKESO Rehabilitation Center. Inclusion criteria included: amputees who are Malaysian citizens aged 18 years or older; with a minimum educational qualification of the Malaysia Lower Certificate of Education (LCE/SRP), Lower Secondary Assessment (PMR), or Form Three Assessment (PT3); have undergone lower limb amputation with a surgical wound that had healed up to four months post-amputation; and are using assistive technology specifically for mobility, either permanently or temporarily awaiting prostheses. This preliminary analysis included 42 participants (Men = 29; Women = 13) aged 48.45 (SD = 10.99) years, with an average duration of mobility device use of 39.82 (SD = 32.05) weeks. They scored an average total satisfaction score of 3.62 (SD = 0.77), with higher satisfaction in device-related components (Mean = 3.70, SD = 0.78). The psychological domain of QOL had the highest score (Median = 66.67, IQR = 54.17 – 79.17). In contrast, the social relationships domain scored the lowest (Mean = 59.92, SD = 20.92). Overall- or device-related satisfaction was not associated with QOL domains. Service-related satisfaction was moderately associated with the social relationships QOL domain ($r_s = 0.340$, $p = 0.028$). Although lower limb amputees report general satisfaction with their mobility devices, service-related issues and low social relationships QOL remain concerns, emphasising the need for improved service delivery and a client-centred approach to rehabilitation.

Keywords: Amputation; assistive technology; lower limb; mobility; quality of life

PP077

Looking is Not Seeing: Objective Gaze Analytics Reveal Limits of Self-Reported Attention in Digital Slit-Lamp Video Training

KHADIJA DAYAH ABDULLAHI, MIZHANIM MOHAMAD SHAHIMIN*,
BARIAH MOHD ALI

*Optometry and Vision Science Program, Centre for Community Health Studies, Faculty of Health Sciences,
Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

**Correspondence: mizhanim@ukm.edu.my*

ABSTRACT

Digital video instruction is central to contemporary clinical skills training in rehabilitation and health sciences. However, learner engagement in digital environments is typically evaluated using self-report surveys, which may not accurately reflect actual attentional behaviour. Eye-tracking technology offers an objective, real-time measure of gaze analytics for visual attention, yet its relationship to perceived engagement remains insufficiently examined in optometry education. This study investigated the relationship between objectively measured visual attention and self-reported attentional engagement during video-based slit-lamp training. Forty-four Year 3 optometry students at Universiti Kebangsaan Malaysia viewed slit-lamp instructional videos while eye movements were recorded using a screen-based eye-tracking system (Tobii Pro Spark). Objective gaze analytics attention was operationalised as dwell time percentage within expert-defined, clinically relevant Areas of Interest (AOIs). Perceived attention was measured immediately post-viewing using the Attention subscale of the Reduced Instructional Materials Motivation Survey (RIMMS). Spearman correlation was conducted, followed by linear regression modelling. Objective dwell time on critical AOIs ($61.65\% \pm 6.76$) was not significantly associated with self-reported attention (Spearman $\rho = 0.21$, $p = 0.18$). Regression analysis confirmed that gaze allocation did not significantly predict perceived attention, $F(1,42) = 1.36$, $p = 0.25$, accounting for only 3.1% of variance ($R^2 = 0.031$). Findings reveal a measurable dissociation between observed visual behaviour and learners' perceived attentional engagement during digital slit-lamp instruction. These results highlight potential limitations of reliance on self-report metrics in evaluating digital clinical education and support the integration of gaze analytics as an innovative tool for optimising instructional design. Embedding objective attention measures into digital health training platforms may enhance evidence-based development of clinical skills education in rehabilitation and health sciences.

Keywords: Digital learning analytics; eye tracking; optometry education; rehabilitation education; video-based training; visual attention

PP079

Prediction of MiR-155 as Biomarkers in Breast Cancer: A Systematic Review

XIAOHUA CHEN^{1,2}, MING THONG ONG^{1*}, HUAYUN QIU²

¹*Institute for Research in Molecular Medicine, Universiti Sains Malaysia, Pulau Pinang, 11800 USM, Pulau Pinang, Malaysia*

²*Medical College of Shaoguan University, Shaoguan City, Guangdong Province 512026, China*

**Correspondence: omt@usm.my*

ABSTRACT

Despite advances in the detection and treatment of breast cancer (BC), it remains a prevalent disease worldwide. miRNAs play an important role in cancer, and their presence may have several advantages in tumour initiation, dissemination, immune evasion, and drug resistance. In general, miR-155 is considered an oncogenic miRNA that promotes tumour growth, angiogenesis, and invasiveness in BC. Therefore, researchers have used it as a predictive biomarker and therapeutic target. However, its predictive value for BC patients remains to be fully elucidated. To address this issue, this systematic review aims to summarise the existing evidence and describe the predictive significance of miR-155 in BC pathology. All eligible studies were searched on PubMed and WOS databases by various search strategies. A total of 37 studies were fully examined from 640 potentially eligible articles, comparing tissue, circulating, and urinary miR-155 expression levels with clinical applications and clinicopathological features in BC patients. We found miRNAs isolated from blood or urine samples of patients diagnosed with BC may be as valuable as those from tissue samples. They could provide information on early diagnosis, disease progression, relapse, treatment response, and metastasis. Simultaneously, challenges that need to be evaluated to understand the ability of miR-155 to address real clinical needs were analysed, highlighting the consistency in translating this molecule into clinical practice. Our paper suggests that the predictive role of miR-155 in BC management requires further validation. This work has been registered in PROSPERO with registration number CRD420251001681 and date of registration 14th May 2025.

Keywords: Blood; breast cancer; miR-155; prediction; tissue

i-001

Next-Generation Targeted Chemotherapy: A Novel Selective Organotin (IV) Compound for Lung Cancer Treatment

NURUL AMALINA ABD. AZIZ¹, NORMAH AWANG^{1*}, NURUL FARAHANA KAMALUDIN¹, ASMAH HAMID¹, CHAN KOK MENG²

¹Center of Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Product Stewardship and Toxicology, Petroliaam Nasional Berhad, PETRONAS Twin Towers, KLCC, 50088 Kuala Lumpur, Malaysia

*Correspondence: norm@ukm.edu.my

ABSTRACT

Lung cancer remains one of the leading causes of cancer-related mortality worldwide. Although cisplatin-based chemotherapy is widely used, its clinical effectiveness is limited by severe systemic toxicity and the emergence of drug resistance. Therefore, the development of safer, selective, and mechanism-driven anticancer agents is urgently needed. This innovation aims to develop and mechanistically validate a novel organotin (IV) N-alkyl-N-benzylthiocarbamate compound as a selective therapeutic candidate against human lung carcinoma (A549) cells. Nine organotin (IV) derivatives were synthesised via an efficient in situ approach (32–89% yield) and structurally characterised using elemental analysis, fourier-transform infrared spectroscopy (FT-IR), nuclear magnetic resonance (NMR) spectroscopy and single-crystal X-ray crystallography. Cytotoxicity was evaluated using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assays, while apoptosis and mechanistic pathways were investigated through annexin V–fluorescein isothiocyanate/propidium iodide (V-FITC/PI) staining, tetramethylrhodamine, ethyl ester (TMRE), dihydroethidium (DHE), comet assay, antioxidant inhibition and caspase activation analyses. This innovation introduces a rationally designed diphenyltin (IV) framework incorporating dual phenyl groups within a thiocarbamate system to enhance cancer-cell selectivity, improve apoptotic targeting efficiency and reduce nonspecific toxicity compared with conventional cisplatin therapy. Unlike cisplatin, which is commonly associated with systemic toxicity and resistance, the synthesised compound demonstrated selective cytotoxicity through reactive oxygen species (ROS)-mediated intrinsic mitochondrial apoptosis, representing a strategic advancement over platinum-based chemotherapy. Seven compounds demonstrated strong cytotoxicity (IC₅₀: 0.52–3.9 µM). The lead compound achieved 60% apoptosis at IC₅₀ (1.86 µM), with significantly higher selectivity toward cancer cells (selectivity index >6; *p* < 0.05) compared with normal cells. Mechanistic validation confirmed significant ROS-mediated DNA damage, mitochondrial membrane depolarisation, and activation of caspase-9 and caspase-3 pathways (*p* < 0.05). These findings highlight its potential as a safer and more targeted alternative anticancer therapy. The compound also demonstrates strong potential for patent protection, preclinical advancement, and pharmaceutical collaboration. Future

studies should focus explicitly on comprehensive in vivo efficacy, toxicity, pharmacokinetic, and therapeutic validation to support its translation into a next-generation targeted anticancer drug.

Keywords: Mitochondrial apoptosis targeting; organotin (IV)-based anticancer agent; selective lung cancer therapy; structure–activity relationship

i-003

Eco-Friendly Ultrasound Coupling Medium Based on Acacia Gum: A Sustainable Alternative to Synthetic Carbomer

KUSNANTO MUKTI WIBOWO^{1*}, NAWANG ESTIANI¹, WINI MELA SAPUTRI¹, FATIATUN FATIATUN^{2,3}

¹Department of Medical Electronics Engineering, Faculty of Health Sciences, Universitas Muhammadiyah Purwokerto, Purwokerto, Indonesia

²Department of Physics Education, Faculty of Education and Teaching, Universitas Sains Al-Quran, Wonosobo, Indonesia

³Department of Physics, Faculty of Science and Mathematics, Universiti Pendidikan Sultan Idris, 35900 Tanjung Malim, Perak, Malaysia

*Correspondence: kusnantomuktiwibowo@ump.ac.id

ABSTRACT

The ultrasonography (USG) gel is an essential coupling medium in diagnostic imaging; however, most commercial formulations still rely on synthetic polymers such as carbomer 940, which are non-biodegradable and potentially contribute to long-term environmental pollution. This study introduces “AcuGel,” a novel eco-friendly USG coupling gel formulated from acacia gum (gum arabic) as a sustainable and biodegradable alternative for greener medical applications. The innovation emphasises the replacement of petroleum-based synthetic gelling agents with a naturally derived biopolymer while maintaining the fundamental physical properties required for ultrasound imaging. Five formulations containing different concentrations of acacia gum (2%, 5%, 15%, 20% and 50%) were prepared with glycerin, propylene glycol, methylparaben, propylparaben and triethanolamine. The developed gels were evaluated through organoleptic observation, pH measurement, viscosity analysis, spreadability testing, and Fourier Transform Infrared (FTIR) spectroscopy to investigate their physicochemical characteristics and functional group compatibility. The organoleptic evaluation demonstrated favorable texture, color uniformity, and physical stability across several formulations. The pH values ranged from 6.5 to 8.0, while viscosity values varied between 60 and 2,400 cP. Spreadability measurements under 50–100 g loads ranged from 11 to 15 cm, indicating good surface coverage capability for ultrasound procedures. FTIR analysis confirmed the presence of characteristic polysaccharide functional groups of acacia gum, including hydroxyl (O–H) and carbohydrate-related bonds, demonstrating successful incorporation of the natural polymer within the gel matrix without significant chemical degradation. The findings highlight the strong potential of AcuGel as a sustainable alternative to conventional synthetic USG gels. This innovation supports environmentally responsible healthcare practices, aligns with the green hospital concept, and offers a promising pathway toward biodegradable medical consumables for future diagnostic applications.

Keywords: Acacia gum; biodegradable; natural formulation; USG gel

i-004

A Single-Stacked Prognostic-Predictive Model for Clinical Risk Stratification of *Staphylococcus aureus* Infection

ZUN LIANG CHUAN*, CHAI JIAN TAY

Universiti Malaysia Pahang Al-Sultan Abdullah, 26300 Kuantan, Pahang, Malaysia

*Correspondence: chuanzl@umpsa.edu.my

ABSTRACT

Staphylococcus aureus (*S. aureus*) infection remains a major cause of morbidity and mortality among burn patients. Collecting time-to-event data for infection risk analysis is often costly, labor-intensive, and logistically challenging due to prolonged patient follow-up and incomplete records. This project develops an innovative single-stacked prognostic-predictive model to identify statistically significant risk factors and confounders for *S. aureus* infection. By eliminating the need for longitudinal time-to-event data, the proposed model reduces follow-up burden, minimises data collection cost, and enables more practical implementation in resource-limited clinical settings. A historical imbalanced dataset of burn patients, comprising 18 demographic and clinical risk factors, was analysed. The CLR-BLR prognostic-predictive model integrates conventional logistic regression (CLR), automated stepwise feature selection, and Bayesian logistic regression (BLR). Its performance (accuracy = 91.56%, recall = 0.7708, precision = 0.9487, and F1-score = 0.8506) and interpretability were benchmarked against CLR, Support Vector Machine, C5.0 and Classification and Regression Tree. The proposed model outperformed all benchmarks, identifying prophylactic antibiotic treatment, infection-related timing, and antibiotic timing as key risk factors, while race, gender, respiratory tract burns, and flame burn etiology were identified as confounders. This model also quantified individual prediction uncertainty through 95% credible intervals. These findings, consistent with current clinical evidence, provide actionable insights for infection prevention and management in burn patients. The model shows potential for integration into hospital risk assessment and clinical decision-support systems to support early identification of high-risk patients without requiring time-to-event data collection. This study aligns with Sustainable Development Goals (SDGs), particularly SDG3 (Good Health and Well-Being) and SDG9 (Industry, Innovation and Infrastructure), by promoting innovative, data-driven healthcare approaches. Future studies should compare the predictive performance and feature consistency between the proposed CLR-BLR prognostic-predictive model and AI-based algorithms, utilising repeated stratified 10-fold cross-validation to evaluate robustness across different random seeds and data partitions.

Keywords: Burn patients; infection prevention; prognostic-predictive model; risk stratification; *Staphylococcus aureus*

i-005

The Read Emotion and Learn to Regulate (REAL™) Module: A Culturally Responsive Skill-Based Mental Health Module for Malaysian Adolescents

**NIK RUZYANEI NIK JAAFAR*, HAJAR MOHD SALLEH SAHIMI, AIMI
NUR ATHIRA PUTRI BAHARUDDIN, NUR HUSNA ISMAIL, NORSINAR
ADILUN**

*Department of Psychiatry, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur,
Malaysia*

**Correspondence: ruzyanei@hctm.ukm.edu.my; njruzyanei@gmail.com*

ABSTRACT

The rising prevalence of anxiety and depression among Malaysian adolescents underscores the need for preventive interventions that move beyond awareness and actively build emotional competencies. This study aimed to evaluate the REAL™ (Read Emotion and Learn to Regulate) module, a culturally responsive, school-based Social Emotional Learning (SEL) programme designed to strengthen emotion-regulation skills among lower-secondary students. The REAL™ module employs a 12-session, weekly, competency-based framework developed through structured expert validation, integrating culturally grounded principles of emotion identification, cognitive reappraisal, behavioural self-regulation, and adaptive coping through guided reflection and experiential learning. Its novelty is enhanced by the MVR-REAL™ Virtual Reality component, which is integrated in the module to provide a private, controlled practice space that addresses key limitations of traditional SEL, such as limited behavioural rehearsal and reduced contextual transfer. It also facilitates Malaysian adolescents who may benefit from learning these skills in a more personal and culturally comfortable setting. The effectiveness of the REAL™ module was tested in a two-arm randomised controlled trial involving 226 adolescents aged 13–14 years from four secondary schools in Selangor, with a waitlist control group. Significant improvements were observed in emotional regulation ($p < 0.001$), anxiety ($p < 0.001$), and depression ($p < 0.001$), with sustained anxiety reduction at six-month follow-up. No adverse events were reported. REAL™ offers a scalable, curriculum-compatible preventive mental health solution with strong empirical support and clear potential for school-based implementation and digital commercialisation.

Keywords: Adolescent mental health; emotion regulation; randomised controlled trial; Social Emotional Learning; virtual reality

i-006

AgeCARE-X: An Intelligent Machine Learning Framework for Predicting Accelerated Ageing in Breast Cancer Survivors and Cancer-Free Women

MOHD RAZIF SHAHRIL^{1*}, NUR AFIFAH MURSYIDA ZAUJAN¹, AZIZI ABDULLAH², SUZANA SHAHAR¹

¹Centre for Health Ageing and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Centre for Artificial Intelligence Technology, Faculty of Information Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

*Correspondence: razifshahril@ukm.edu.my

ABSTRACT

AgeCARE-X is a breakthrough predictive modeling framework engineered to transform the mitigation of biological aging in breast cancer survivors. Its core innovation lies in a quadrant-based visual interpretive structure designed to serve as a transparent decision-support tool. This layout immediately translates complex data into actionable insights, allowing users to visualise the tug-of-war between “decelerating” protective factors (e.g., nutrient adequacy) and “accelerating” risk factors (e.g., radiotherapy impact and stress). By prioritising this visual clarity, AgeCARE-X turns “black-box” machine learning into a clear, evidence-based roadmap for personalised longevity. To power this interface, the framework employs a high-precision computational pipeline. By integrating XGBoost with SMOGN for data balancing and Particle Swarm Optimisation for refinement, AgeCARE-X achieves superior predictive accuracy over conventional models. This architecture analyses multidimensional inputs including lifestyle, clinical history and functional fitness, to predict telomere length shortening, bridging the gap between high-level data science and practical health applications. This innovation carries immense commercial potential within the digital health and personalised survivorship sectors. By identifying modifiable behaviors, it provides a scientific foundation for interventions aimed at extending biological health span. Future research will leverage this framework to develop an AI-driven advocacy chatbot. This interface will deliver real-time, evidence-based guidance directly to survivors, democratising access to aging-mitigation strategies and empowering patients to take an active role in their long-term recovery.

Keywords: Accelerated ageing; breast cancer survivorship; framework; machine learning, telomere length

i-007

The Sound of Learning: A Phoneme-Based Book with App Support for Early Literacy

SITI ATIYAH ALI^{1*}, ZAINE ZULKIFLI², NURFAIZATUL AISYAH AB AZIZ¹,
JANET ANN FERNANDEZ², MOHAMAD HARDYMAN BARAWI¹

¹*Cognitive Science Programme, Faculty of Cognitive Science and Human Development, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia*

²*Psychology Programme, Faculty of Cognitive Science and Human Development, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia*

**Correspondence: asatiyah@unimas.my*

ABSTRACT

This project develops a multisensory learning tool in the form of a printed storybook and a mobile application to support children with reading difficulties, particularly those who struggle to connect written letters (graphemes) with their corresponding sounds (phonemes). Many children with learning difficulties, such as dyslexia, experience challenges in phoneme-grapheme mapping, which is essential for reading development. The developed book and application integrate visual and auditory elements to enhance learning through a multisensory approach, allowing children to learn through both seeing and hearing. The features include audio-supported storytelling with synchronised word highlighting, interactive sound buttons that allow children to hear word pronunciation, and a structured progression from simple to more complex word forms. Unlike existing phonics applications that mainly focus on screen-based activities, or traditional reading interventions that rely mostly on printed materials and teacher guidance, this tool combines physical reading materials with digital multisensory support. The word content in the book was obtained through the screening of words used in multiple storybooks for children aged 3 to 7 years old. Simple words with two to three phonemes were identified and included in this phoneme-based book. The suitability of the words used, and the usability of the application were tested among ten kindergarten children aged 4 to 6 years old. In addition, two kindergarten teachers provided feedback on the usability of the application and commented on its ease of use and user-friendliness. The product is still being developed and improved by collecting feedback from children and educators. Optional bilingual support is also included to accommodate multilingual environments such as Malaysia. Grounded in educational and cognitive research, this tool is expected to improve phonemic awareness, support early reading skills, and increase children's confidence in reading. The developed book and application can serve as practical resources for schools, therapy centres, and parents, while also providing a model for the future development of inclusive learning materials for children with special learning needs.

Keywords: Children; early literacy; multisensory learning; phoneme-grapheme

i-009

Skinova™: Translational Development of a *Centella asiatica*-Based Topical Antibacterial Cream

SITI KHADIJAH PADZIL*, ABDUR RAHMAN AUF ALI, NURUL AINAA
NAJLA YUSRI, NOR IZZA KAMILIA NORSIDEK, ATHIRSHDA

KPJ Healthcare University, Kota Seriemas, 71800 Nilai, Negeri Sembilan, Malaysia

*Correspondence: ucn.khadijah@kpju.edu.my

ABSTRACT

The increasing demand for plant-based dermatological products necessitates the development of scientifically optimised and standardised herbal formulations. *Centella asiatica* is widely recognised for its wound healing and antimicrobial properties; however, limited studies emphasise formulation robustness alongside antibacterial efficacy. This study aimed to develop and comparatively evaluate 2.5% and 7.5% *C. asiatica* ethanolic cream formulations with emphasis on physicochemical performance and functional activity. Ethanolic extracts were prepared via cold maceration and incorporated into oil-in-water cream systems. Formulations were evaluated for organoleptic properties, pH, viscosity, washability, homogeneity, emulsion type (dye test), and phase stability over 30 days. Antibacterial activity was assessed against *Staphylococcus aureus*, *Escherichia coli*, and *Bacillus cereus* using well diffusion and minimum inhibitory concentration (MIC) assays. Both formulations demonstrated favourable physicochemical properties, with skin-compatible pH (4.79–5.36), good washability, and stable oil-in-water characteristics. The 7.5% formulation exhibited improved viscosity and spreadability compared to the 2.5% formulation, although minor phase separation was observed at day 30. Antibacterial testing revealed a concentration-dependent effect, with the 7.5% cream producing larger zones of inhibition (16.5–22.5 mm) and an MIC of 50 mg/mL across tested organisms, while the 2.5% formulation showed comparatively lower activity. No bactericidal effect was observed. In conclusion, the 7.5% *C. asiatica* formulation demonstrates superior functional and physicochemical performance, supporting its potential as a stable, plant-based topical antibacterial product. This study highlights formulation optimisation as a key step toward translational and commercial readiness.

Keywords: Antibacterial; *Centella asiatica*; formulation optimisation topical cream; physicochemical evaluation

i-010

EcoDish+: A Sustainable and Impactful Health Innovation in Antibacterial Dishwashing Soap from Recycled Cooking Oil, *Padina* spp. and Aloe Vera

FATTAH FAZEL¹, AFIQ AZIL², NUR BATRISYA², NURUL HANNANI²,
NORAZALINA MOHAMAD ZAH^{2*}

¹Research Management Department, KPJ Healthcare University, Kota Seriemas, 71800 Nilai, Negeri Sembilan, Malaysia

²School of Pharmacy, KPJ Healthcare University, Kota Seriemas, 71800 Nilai, Negeri Sembilan, Malaysia

*Correspondence: norazalina@kpju.edu.my

ABSTRACT

Improper disposal of used cooking oil contributes significantly to soil and water pollution, while conventional dishwashing liquids often contain harsh chemicals that may irritate the skin. Sustainable alternatives are needed to address both environmental and health concerns. This study developed EcoDish+, an antibacterial liquid dishwashing soap formulated from recycled cooking oil and enriched with natural extracts of *Padina* spp. and Aloe Vera. The process involved purification of waste oil, saponification, and incorporation of bioactive agents selected for their antibacterial, soothing, and moisturising properties. Phytochemical screening was conducted to identify active compounds, and antibacterial activity was assessed using the well diffusion assay against *Escherichia coli* and *Staphylococcus aureus*. Phytochemical analysis identified the presence of flavonoids, phenolics, and fixed oils, compounds associated with antimicrobial activity. The formulated soap demonstrated antibacterial properties, with inhibition zones of 22 mm against *S. aureus* and 15 mm against *E. coli*, indicating stronger activity against Gram-positive bacteria. Quality and safety evaluation further confirmed the suitability of EcoDish+ for consumer use. The formulation exhibited a near-neutral pH of approximately 7.5, meeting standards. Foam stability testing recorded an initial foam height of 100 mm, retained at 90 mm after five minutes, placing it well within the acceptable range of 13–220 mm and demonstrating cleaning performance. In addition to these physicochemical properties, the formulation exhibited moisturising benefits, making it safer for frequent household use. EcoDish+ represents an integration of recycled waste oil with marine and plant-derived bioactive ingredients, offering dual benefits of environmental sustainability and health protection.

Keywords: Aloe vera; antibacterial dishwashing soap; *Padina* spp.; recycled cooking oil; sustainable health innovation

i-011

BETFresh: Fast-Dissolving Herbal Paper Breath Strips Containing *Piper betle* L. Extract for Natural Oral Care

AHMAD FARID, NUR BATRISYA, NURUL HANNANI, AFIQ AZIL,
NORAZALINA MOHAMAD ZAH*

School of Pharmacy, KPJ Healthcare University, Kota Seriemas, 71800 Nilai, Negeri Sembilan, Malaysia

*Correspondence: norazalina@kpju.edu.my

ABSTRACT

Halitosis, or bad breath, is a widespread oral health issue that can disrupt social confidence and overall well-being. Conventional oral care products often rely on synthetic ingredients that provide only temporary relief and may cause irritation or dryness. To address these limitations, this study introduces BETFresh, a fast-dissolving herbal breath strip formulated with *Piper betle* L. extract as a natural antibacterial and deodorising agent. The innovation integrates traditional herbal medicine with modern strip technology to deliver a safe, portable, and user-friendly oral hygiene breath strip. *Piper betle* leaves were extracted using ethanol and water, followed by phytochemical screening that confirmed the presence of flavonoids, phenols, terpenoids and other bioactive compounds known for antimicrobial activity. Antibacterial evaluation against common oral pathogens, including *Streptococcus mutans* and *Staphylococcus aureus*, demonstrated significant inhibition, with minimum inhibitory concentration and minimum bactericidal concentration values of 1.25 mg/mL and 2.5 mg/mL, respectively. The ethanolic extract exhibited stronger antibacterial performance and was selected for strip formulation. Using the solvent casting technique, the extract was incorporated into thin polymeric films to produce fast-dissolving oral strips. BETFresh strips were evaluated for pharmaceutical properties, showing uniform thickness (0.20–0.25 mm), near-neutral pH (6.7–6.9), and rapid disintegration in under 30 seconds. These characteristics ensure both safety and convenience, while the antibacterial activity confirms dual functionality, instant breath freshness and microbial protection. The novelty of BETFresh lies in its sugar-free, halal-friendly design and its ability to combine herbal bioactivity with a modern dosage form. With low-cost formulation, scalable manufacturing, and strong consumer appeal, BETFresh demonstrates significant commercialisation potential for distribution through pharmacies, clinics, and e-commerce platforms. This innovation represents a promising step toward natural, effective, and accessible oral care breath strips.

Keywords: Fast-dissolving oral strips; halitosis management; herbal oral care; natural breath freshener; *Piper betle* L.

i-012

UroScan AI: An AI-Enhanced Camera-Based Urine Strip Detection System for Home and Clinical Applications

AHMAD LUQMANULHAKIM SUNAWARI, SHAHRINA ISMAIL, AZIRA KHALIL*

Faculty of Science and Technology, Universiti Sains Islam Malaysia, 71800 Nilai, Negeri Sembilan, Malaysia

**Correspondence: azira@usim.edu.my*

ABSTRACT

The increasing prevalence of chronic diseases such as diabetes and kidney disorders highlight the urgent need for accessible, reliable and user-friendly diagnostic tools. Conventional urine strip testing, while widely used, relies heavily on subjective visual interpretation, making results prone to human error and environmental variability. This project introduces UroScan Artificial Intelligence (AI), an AI-enhanced, camera-based urinalysis system designed to transform traditional strip testing into a standardised, intelligent, and digitally integrated health screening solution. The system utilises advanced image processing techniques, including region-of-interest segmentation, Red-Green-Blue (RGB) extraction, and CIEDE2000 ΔE colour difference analysis, to quantitatively evaluate multiple biochemical parameters using a standard smartphone camera without additional hardware. An integrated AI-powered conversational module further enhances the system by providing contextual interpretation, personalised health insights, and guided user interaction. Performance validation demonstrated an overall accuracy of 86.75%, with high sensitivity and strong classification reliability, indicating its effectiveness as a point-of-care screening tool. UroScan AI not only reduces interpretation variability but also enables digital record tracking for longitudinal health monitoring. By bridging the gap between conventional testing and intelligent diagnostics, this innovation improves early detection, empowers users with actionable insights, and supports proactive healthcare management. With scalable deployment through mobile platforms, clinical integration, and industry partnerships, UroScan AI presents strong potential as a next-generation digital health solution in the evolving MedTech landscape.

Keywords: Artificial intelligence; image processing; mobile health; point-of-care testing; urine strip detection

i-019

ClinicaGuide.com: An AI-Powered Clinical Decision Support System for Evidence-Based Allied Healthcare Practice

SAIFUL IZZUAN HUSSAIN^{1,2*}, SHAMSUDDIN SALLEH²

¹*Department of Mathematical Sciences, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia*

²*Scancer AI Sdn Bhd, Akademia Siber Teknopolis, Universiti Kebangsaan Malaysia, 43600 UKM Bangi, Selangor, Malaysia*

**Correspondence: sih@ukm.edu.my*

ABSTRACT

Healthcare professionals often face challenges in accessing timely, evidence-based clinical information during patient care, leading to decision delays and potential suboptimal outcomes. Traditional methods of searching medical literature are time-consuming and may not provide actionable guidance at the point of care. Therefore, innovative digital solutions are needed to support rapid clinical decision-making. This innovation aims to develop ClinicaGuide.com, an artificial intelligence (AI)-powered clinical decision support platform that delivers instant, evidence-based answers to allied health professionals. The system was designed with natural language processing capabilities, a comprehensive medical knowledge base, and an intuitive interface to provide real-time clinical guidance tailored to user queries. This promotes efficient information retrieval while ensuring accuracy and reliability. A pilot feasibility study was conducted to assess usability, response accuracy, and user satisfaction among healthcare practitioners. The novelty of this innovation lies in its AI-driven search architecture, localised content relevance, and accessibility across clinical and community settings. Unlike conventional medical databases, it is user-friendly, rapid, and scalable for diverse healthcare environments. The expected impact includes improved clinical decision-making, enhanced healthcare efficiency, reduced information-seeking time, and better patient outcomes. Commercially, the platform has strong potential for implementation in hospitals, rehabilitation centres, private clinics, and academic institutions, offering a scalable solution to democratise evidence-based practice in allied healthcare.

Keywords: Allied health; artificial intelligence; clinical decision support; digital health innovation, evidence-based practice

i-020

Development of Inventory of Communication Need

AUNI SYAFIQAH ROSLAN, SUSHEEL KAUR DHILLON JOGINDER
SINGH, FATIMAH HANI HASSAN*

Department of Speech Science, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

**Correspondence: fatimahhani@ukm.edu.my*

ABSTRACT

Aphasia disrupts all aspects of language comprehension and expression, which often compromises aphasia survivors' (AS) quality of life and daily functioning. Currently, Malaysia lacks a dedicated tool to document these challenges from the AS perspective, which leaves their specific communication needs under-researched. This study aims to develop and validate the Inventory of Communication Needs (ICON), an aphasia-friendly survey designed to explore the met and unmet communication needs of AS from their own perspective. A mixed-methods design was conducted in three stages. Stage 1 involved survey development based on 12 studies, resulting in 52 items presented in Malay and English using aphasia-friendly principles such as simple language, visual supports, clear layout, and flexible response time. Stage 2 involved validation by nine validators (speech-language therapists, AS, and caregivers) to assess item suitability, language comparability, and images using the Item Content Validity Index (I-CVI) and Kappa statistics. Stage 3 evaluated feasibility and reliability with six AS of varying severity, including refinement of administration procedures. The final ICON consists of 41 items, with strong content validity (I-CVI = 0.88) and excellent agreement (Kappa = 0.89). ICON is the first tool to empower AS to express their own needs, reducing reliance on caregivers and supporting greater participation. It is a bilingual and culturally relevant tool developed for the Malaysian context. The survey uses aphasia-friendly design features to support individuals across different severity levels and provides a structured way to identify communication needs for better intervention planning. Suitable for hospitals, rehab centers, and universities, ICON is a vital resource for healthcare professionals and researchers. Its potential as a digital tool will broaden patient-centered care in Malaysia, facilitating more precise intervention planning and evidence-based rehabilitation programs.

Keywords: Aphasia; aphasia-friendly; aphasia tool; communication need; survey development

i-021

Designing a Mobile Application for Older Adults with Dementia: Insights from Expert Perspectives

MUHAMMAD REDZUAN BIN MUHD^{1,2}, DEVINDER KAUR AJIT SINGH¹,
LEIGH HALE³, NORMALA MESBAH^{1*}

¹*Centre for Healthy Aging and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Physiotherapy Unit, Hospital Sultan Haji Ahmad Shah, 28000 Temerloh, Pahang, Malaysia*

³*School of Physiotherapy, University of Otago, Dunedin 9016, New Zealand*

**Correspondence: normala.mesbah@ukm.edu.my*

ABSTRACT

Mobile applications are increasingly vital in modern healthcare, yet digital tools rarely account for the specific cognitive and physical challenges faced by older adults living with dementia. This study sought to bridge this gap by establishing a multidisciplinary framework for a dementia-specific mobile health application. We conducted an interactive focus group discussion with an expert clinical panel, including a geriatrician (n = 1), specialised nurses (n = 2), occupational therapists (n = 2), and physiotherapists (n = 3), to capture frontline insights on clinical needs and practical design features. Using conventional content analysis, the qualitative data yielded five core thematic categories: user interface design, core functional requirements, clinical benefits, real-world implementation barriers, and future prototype enhancements. The panel emphasised that to combat cognitive deterioration, the interface must prioritise minimal text, high-contrast visual cues, and culturally tailored elements. Crucially, the innovation integrates tailored physical exercises, safety guidelines, and active caregiver support modules, such as task reminders and tracking, into a single ecosystem. This design directly addresses current market gaps by shifting from passive tracking to active, home-based rehabilitation and continuity of care. While implementation barriers like digital literacy were identified, the panel recommended automated tracking and motivational features to ensure long-term adherence. Ultimately, this research provides an actionable, clinician-vetted blueprint that enhances the feasibility, clinical safety, and user acceptance of digital interventions. By translating geriatric physiotherapy principles into a user-friendly digital interface, this application holds high commercialisation potential as a scalable, reimbursable digital health tool for home care interventions, rehabilitation centers, and family caregivers navigating dementia management.

Keywords: Dementia; digital health intervention; mobile application; older adults

i-022

CHEL 2.0 ACETA: Portable Hydrogel-Based Sensing for Detection of Chemical Contaminants and Pharmaceutical-Related Responses in Water

XU HOE CHUNG*

School of Pharmacy, IMU University Malaysia, Bukit Jalil, 57000 Kuala Lumpur, Malaysia

**Correspondence: 00000046746@student.imu.edu.my*

ABSTRACT

Water contamination by pharmaceutical and chemical residues presents a growing environmental and public health concern, particularly in low-resource settings where conventional analytical instrumentation is limited. The objective of this study is to develop a portable, low-cost hydrogel-based sensing platform capable of qualitative detection with real-time digital signal tracking of aqueous chemical contaminants. CHEL 2.0 ACETA utilises iron(III)-loaded hydrogel beads embedded with a colourimetric indicator system that undergoes a visible colour transition upon interaction with oxalic acid as a model analyte. The hydrogel matrix enables diffusion-driven and chelation interaction while maintaining structural integrity. To enhance analytical performance, the system is integrated with a microcontroller-based colour sensor that records real-time colour change signals. A pH sensing module is also incorporated to correlate acidity changes with observed colourimetric responses. Data are transmitted to spreadsheet software for real-time graphical visualisation of Red, Green and Blue (RGB) signals and pH trends. Additionally, preliminary observations reveal unexpected buoyancy and sinking behaviour of hydrogel beads in the presence of paracetamol, suggesting potential polymer-solute interactions that may influence hydrogel physical properties. CHEL 2.0 ACETA integrates hydrogel chemistry with digital analytics for qualitative detection with real-time signal monitoring of chemical contaminants. The system's modular design, low-cost materials, and real-time data capability highlight its potential for field-based environmental monitoring and preliminary pharmaceutical contamination screening. The platform may be developed into a portable analytical device for water quality assessment in resource-limited environments.

Keywords: Colorimetric detection; environmental contamination; hydrogel-based sensor; portable analytical device; real-time chemical monitoring

i-023

Augmented Reality in Dental Hygiene: Bridging Technology and Patient Acceptance

ANISYA HANUM OTHMAN¹, NIK RAHAYYU NIK ZULKIFELI^{2*},
NAZURAH NIK EEZAMMUDDEEN², AZWIN ASSILAH KAMARUDDIN²,
AIEMEEZA RAJALI²

¹Postgraduate Student, Faculty of Dentistry, Universiti Teknologi MARA, 47000 Sungai Buloh, Selangor, Malaysia

²Faculty of Dentistry, Universiti Teknologi MARA, 47000 Sungai Buloh, Selangor, Malaysia

*Correspondence: nikrahayyu7877@uitm.edu.my

ABSTRACT

Augmented reality (AR) has emerged as an educational technology in healthcare, offering immersive and interactive learning experiences that may enhance patient understanding regarding the care of dental prostheses. The successful implementation of AR depends largely on the patients themselves. To assess the acceptance of dental patients regarding the use of AR in dental prostheses care education. A qualitative study was conducted using in-depth interviews with dental patients from Universiti Teknologi MARA Sungai Buloh. The selection of participants with prostheses was done by the convenience sampling method. The novelty of this innovation lies in the integration of a prostheses care book with AR technology compared to a traditionally static print learning resource, transforming it into an interactive platform. Unlike conventional dental guidebooks that rely solely on two-dimensional illustrations, the AR book allows patients to visualise three-dimensional oral anatomy and hygiene instruction in real time through mobile devices. A total of ten adults, aged up to 69 years old, were interviewed in this study. The thematic analysis of their responses revealed several major patterns and insights. Patients preferred an interactive, easy-to-use, AR-based educational format. They perceived this approach as beneficial for improving their understanding of prostheses care instructions, particularly in visualising and following the required steps more effectively. Nevertheless, some older participants hesitated as they need guidance from others due to limited familiarity with digital tools. Although patients view AR as a helpful teaching method for prostheses care, significant barriers exist concerning accessibility and digital skills for widespread adoption. Future research should focus on developing user-friendly AR applications tailored to patient needs. Embedding AR into the dental guidebook introduces a next-generation learning platform. This book will incorporate AR features to make it an interactive educational experience.

Keywords: Augmented reality guidebook; dental prostheses care; patient education

i-024

DengarTrack: A Digital Workflow Innovation to Reduce Loss-To-Follow-Up in Universal Newborn Hearing Screening

SITI AMINAH KAMALUDIN^{1*}, MUHAMMAD UWAIS ZULKARNAIN²,
ONN WAH LEE¹, RAFIDAH MAZLAN¹

¹*Audiology Programme, Centre for Rehabilitation & Special Needs Studies, Faculty of Health Sciences,
Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Faculty of Artificial Intelligence, Universiti Teknologi Malaysia, 54100 Kuala Lumpur Malaysia.*

**Correspondence: p153232@siswa.ukm.edu.my*

ABSTRACT

Universal newborn hearing screening (UNHS) enables early detection of congenital hearing loss, but its benefits depend on timely follow-up; infants diagnosed and enrolled in intervention within the recommended window achieve language outcomes comparable to those of hearing peers, while delays cause largely permanent harm. In Malaysian public hospitals, UNHS still relies on paper records, spreadsheets, and informal communication channels. This fragmented workflow has contributed to a mean loss-to-follow-up rate of 23% across 48 hospitals-over four times the ≤5% benchmark established by the Joint Committee on Infant Hearing (JCIH, 2007), whose 2019 statement retains this priority through process safeguards and 1-3-6 timeliness goals. This innovation aims to develop, implement, and evaluate DengarTrack, a digital workflow platform supporting real-time tracking and coordinated follow-up. Developed with clinical and user input and aligned with Ministry of Health reporting requirements, DengarTrack combines four capabilities in one platform: case-level follow-up tracking, automated loss-to-follow-up alerting based on the JCIH risk indicators, parent-facing multilingual communication, and national registry integration via Health Level Seven Fast Healthcare Interoperability Resources (HL7 FHIR R4). It features Quick Response (QR) code-based newborn identification, a real-time dashboard, role-based access, and offline capture, and runs on Android devices already in use in hospitals. Planned evaluation comprises usability testing in selected public hospitals followed by a quasi-experimental study targeting a 30% relative reduction in loss-to-follow-up, alongside improved timeliness and user adoption. Its novelty is that, to our knowledge, no published system in a low- or middle-income setting unites all four capabilities, making DengarTrack first-in-class in this context. The expected impact includes reduced loss to follow-up, earlier intervention, lower staff workload, and stronger data governance. Commercially, DengarTrack is positioned for adoption within the Ministry of Health hospital network, offering a scalable, standards-based model transferable to other low- and middle-income settings.

Keywords: Digital health; loss-to-follow-up; mHealth; Malaysia; universal newborn hearing screening

i-025

MotherCare Connect: A Family-Centered Digital Platform for Continuous Postnatal Mental Health Monitoring in Malaysia

SIEW EN CHENG*, ZAHRA SOFEA IDONGESIT

Jeffrey Cheah Sunway Medical School, Sunway University, Bandar Sunway, 47500 Petaling Jaya, Selangor, Malaysia

**Correspondence: 24049017@imail.sunway.edu.my*

ABSTRACT

A total of 6.8–27.3% of Malaysian mothers are affected by postnatal depression (PND), yet many remain underdetected due to fragmented psychosocial support within current postnatal care pathways. Critically, PND does not follow a predictable onset timeline. Over 57% of women with depressive symptoms at 9–10 months postpartum were asymptomatic at earlier screenings. This exposes a fundamental limitation of existing care models, highlighting that single-point assessments leave much of the postpartum year unmonitored. This gap represents a significant public health challenge, directly impacting SDG3 (Good Health and Well-being) and SDG4 (Quality Education). MotherCare Connect is an integrated digital health platform designed to shift postnatal care from one-time screening to continuous, preventive monitoring. It integrates five evidence-based components: longitudinal mental health tracking using the Edinburgh Postnatal Depression Scale (EPDS) along with simple mood, sleep, and stress weekly postpartum assessments; automated referral pathways to licensed therapists; evidence-based parenting education; moderated peer-support communities; and paternal inclusion through validated EPDS screening, as paternal depression affects up to 50% of fathers whose partners have PND, amplifying family-wide risk. The platform is built on a tri-sector partnership model aligned with SDG17, integrating hospitals (clinical governance and referral networks), industry partners (sustainable funding through sponsorship), and academia (healthcare students and professionals delivering community workshops and peer-support through supervised training and volunteering). This model ensures clinical credibility and financial viability. By digitising postnatal care, it reduces unnecessary healthcare utilisation and associated carbon emissions. A Phase 2 AI-driven risk stratification module is planned to enable early identification of high-risk mothers, positioning MotherCare Connect as a scalable innovation at the intersection of maternal mental health, digital health equity, and environmental sustainability. Priced at RM120 lifetime membership with instalment options and income-based financial assistance, MotherCare Connect is designed for equitable access.

Keywords: Continuous monitoring; digital health platform; health equity; maternal mental health; postnatal depression

i-026

MedTech's Digi-Screening Software Update

SHIYIN PANG, YASWINII A/P SURESH KUMAR*, KIT LING LIM,
FAUSTINA SARAN D'CRUZ, SHARWEENA A/P MURALITHARAN

*Jeffrey Cheah Sunway Medical School, Sunway University, Bandar Sunway, 47500 Petaling Jaya, Selangor,
Malaysia*

**Correspondence: 24096646@imail.sunway.edu.my*

ABSTRACT

Feedback from patients in health screening at private hospitals include long waiting times. This reduces patient well-being (SDG 3 - Good Health and Well-being) as it causes overcrowding and increases the risk of contracting diseases. We at MedTech propose an innovation to the SunMed app to guide patients to the screening station with the shortest queue. Nurses scan the patient's individual QR code (available in their app) when entering a station to check the patient in, giving immediate access to the patient's digital record. Upon completing the test, completion is submitted by the stationed nurse and updates the particular patient's app. It then automatically provides the patient with the next station, the waiting time, and the number of patients in queue. Firstly, optimised queuing systems prevent overcrowded waiting areas, reducing the risk of transmitting infectious diseases. Furthermore, wait time reduction reduces anxiety in patients (SDG 3). Secondly, this innovative software increases resource efficiency as nurses do not have to look for patients in the numerous waiting rooms (SDG 9 - Industry, Innovation and Infrastructure). Additionally, digital ticketing queues will reduce paper consumption. As workload on healthcare workers at health screening stations can be reduced due to digital guidance for patients, they could be relocated to more demanding departments to reduce strain on others in sectors such as the Emergency Department. Finally, patients won't need to return at a later date due to long queues for stations such as CT scans and MRIs. This will decrease the carbon emissions caused by commuting to the hospital. Hence, the overall carbon footprint of the hospital will also be reduced (SDG 13 - Climate Action). Since this is only a modification to the pre-existing software, implementation cost would be low. In increasing health screening efficiency, overall community satisfaction would elevate.

Keywords: Digital healthcare; healthcare innovation; health screening; patient flow optimisation

i-027

BattCycle Vault: An Innovative and Sustainable Partitioned System for Battery Waste Management and Environmental Education

**ATIKAH MOHD NASIR*, NOR AYUSLINAWATI CHE SIDIK,
KHAIRUNNISA ZAIDAN, MIRA HAZIRA ZAID, SERI MIRIANTI ISHAR**

Forensic Science Programme, Centre for Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

**Correspondence: atikahnasir@ukm.edu.my*

ABSTRACT

The rapid proliferation of electrical and electronic equipment has catalysed a corresponding surge in electronic waste (e-waste), rendering it one of the fastest-growing and most hazardous waste streams globally. Under Malaysia's Environmental Quality (Scheduled Wastes) Regulations 2005, batteries are strictly classified as scheduled waste. Improper disposal of used batteries poses a severe environmental threat, leading to toxic chemical leaks, such as mercury and lead, and causing critical, long-lasting soil and water pollution. To address this escalating challenge, the primary goal was to develop the BattCycle Vault, aiming to establish a highly responsible battery-waste management system while fostering comprehensive environmental education. Replacing traditional, flimsy cardboard collection boxes, this initiative utilises a highly durable, portable plastic toolbox. Beyond physical collection, it operates an extensive educational campaign driven by the unique motto, "Don't Bin It, BattCycle It!" utilising targeted posters and campus exhibitions to proactively educate the university community. The core innovation lies within its uniquely designed partitioned storage system. This structural improvement effectively separates single-use primary batteries from rechargeable secondary batteries, an advancement that ensures much safer handling and rigorously prevents dangerous reactive chemical leaks during storage. The BattCycle Vault successfully increases community awareness while building a proactive green culture. By streamlining the transportation process to certified recycling centers, it facilitates organised, efficient recycling. This initiative helps recover valuable metals, significantly mitigates toxic pollution, and can reduce greenhouse gas emissions by up to eighty percent compared to producing new batteries from raw materials. Ultimately, the BattCycle Vault delivers a high-impact, easily implementable, and scalable model for sustainable institutional waste management. Offering significant cost and time savings, it presents exceptional commercial viability for widespread adoption across educational institutions, corporate offices, and municipalities globally seeking highly efficient, turnkey sustainability solutions for effectively handling hazardous e-waste and fostering enduring, eco-friendly daily community behaviors.

Keywords: Battery; e-waste; environmental education; recycle

i-028

MAVeRT™: A Clinician-Monitored Digital Vestibular Rehabilitation Platform for Accessible Home-Based Allied Health Care

NUR DINI IZZATI KHAIROL ANUR¹, NOR HANIZA ABDUL WAHAT^{1*},
FADILLA' ATYKA NOR RASHID²

¹Centre for Rehabilitation and Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Centre of Artificial Intelligence Technology, Faculty of Information Science & Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

*Correspondence: nor_haniza@ukm.edu.my

ABSTRACT

Introduction: Vestibular disorders commonly cause dizziness, imbalance, and an increased risk of falls, significantly affecting daily functioning and quality of life. Access to structured vestibular rehabilitation therapy (VRT) remains limited due to reliance on supervised clinical sessions and resource constraints. Digital health technologies offer opportunities to improve accessibility through remote and home-based rehabilitation delivery. **Objectives:** This innovation aimed to develop, validate, and pilot the Malaysian Adult Vestibular Rehabilitation Therapy (MAVeRT™) platform as a clinician-monitored, home-based digital VRT system with remote supervision and adherence tracking. **Methods:** The platform was developed from scratch using a multi-phase research-to-innovation approach. Evidence-based vestibular rehabilitation exercises were produced as guided instructional videos. Content validation by rehabilitation experts (mean age: 39.17 years; mean professional experience: 13.33 years) yielded Item Content Validity Index (I-CVI) values ranging from 0.83 to 1.00 and a Scale Content Validity Index average (S-CVI/Ave) of 0.96 for scripts and 1.00 for videos. A mobile prototype integrating validated videos and a digital diary was evaluated for usability, achieving a mean 'Skala Kebolehgunaan Aplikasi Mudah Alih' (SKAMA) score of 71.25 (SD = 19.63), exceeding the benchmark score of 68. The prototype was subsequently enhanced into a fully functional platform integrated with a clinician dashboard website. **Novelty:** MAVErT™ integrates validated exercise content, patient self-monitoring, and clinician dashboard monitoring within a single ecosystem. Distinctive features include AI-assisted movement accuracy detection, real-time motion capture feedback, automated compliance monitoring, smart reminders, and session-resume functionality. **Impacts:** MAVErT™ enhances access to home-based VRT with clinician oversight, reduces clinical burden, shortens waiting times, improves patient adherence, and enables data-driven patient management through continuous monitoring and analytics. **Commercialisation Potential:** MAVErT™ is scalable as a subscription-based digital health solution for hospitals, audiology and physiotherapy clinics, and rehabilitation centers, with opportunities for institutional licensing and tele-rehabilitation integration.

Keywords: Digital health; mobile health application; tele-rehabilitation; vestibular rehabilitation therapy

i-029

CT-Trans: Bridging Healthcare Innovation and Engineering Partnerships Towards Global Transformation in Specimen Transportation

SITI NOOR AISYAH MD HUSSIN*, SHAHRIL AZIAN MASROM, NUR AZLIN MUSTAPHA, NUR EYUNI MOHD SALLEH, NOR ZAHIDI MOHD KODORI, NORHUDA JAAFAR, NURHIDAYAH MD ZAN, NURHASLINDA AMBO

Tibi/Leprosy Unit, Disease Section, Johor Bahru Public Health Laboratory, Tampoi, 81200 Johor Bahru, Johor, Malaysia

**Correspondence: cnoeye@yahoo.co.uk*

ABSTRACT

Johor Bahru Public Health Laboratory (JBPHL), the tuberculosis (TB) reference laboratory in Johor, identified persistent sputum specimen leakage rates of 1.9% (2019) and 1.2% (2020), compromising diagnostic quality, biosafety, and operational efficiency. To address this, the Container Transportation System (CT-Trans) was developed as a simple yet transformative solution to ensure upright, stable, and secure sputum specimens during transportation. CT-Trans is a low-cost (≈RM0.10/unit), high-impact innovation made from recycled plastic materials, aligning with sustainable healthcare practices and the 3R (Reduce, Reuse, Recycle) concept. Each unit accommodates up to four specimens, minimising movement and preventing leakage during transit. The project aimed to reduce leakage rates to below 1% in accordance with Malaysian Society for Quality in Health (MSQH) standards. Implemented in phases (pilot in 2021, statewide expansion in 2022, and culture-building phase from 2023), a cross-sectional analysis (2021–2025) demonstrated a consistent, sustained reduction in leakage rates from 0.65% (2021) to 0.085% (2025), surpassing national benchmarks. Operational costs were reduced from RM5000 to RM200, with reduced specimen rejection, minimised repeat patient sampling, and improved biosafety compliance. Novelty lies in its frugal, sustainable innovation approach, converting waste materials into a practical laboratory solution with immediate scalability. Unlike conventional transport methods, CT-Trans integrates cost-efficiency, environmental sustainability, and workflow optimisation into a single intervention. Collaboration with a university mechanical engineering department further advances product robustness and resilience, demonstrating the power of global and interdisciplinary partnerships. From a commercialisation perspective, CT-Trans demonstrates high scalability and replicability across healthcare facilities, particularly in resource-limited settings. Its low production cost, ease of assembly, strong performance outcomes, and ongoing engineering optimisation position it for mass production, intellectual property development, and global adoption, advancing healthcare innovation.

Keywords: Container Transportation System; global partnership; leakage; sustainable healthcare; tuberculosis

i-030

Intervention for Older Malaysians Living with Knee Osteoarthritis and Diabetes Mellitus

SUMAIYAH MAT^{1*}, MALAK OMAR ABBAAS ALABABSEH^{1,2}, AZIANAH MOHAMAD IBRAHIM¹, DEVINDER KAUR AJIT SINGH¹, ON BEHALF OF THE I-MOAD RESEARCH GROUP

¹Centre for Healthy Ageing and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, J50300 Kuala Lumpur, Malaysia

²Physiotherapy Programme, Faculty of Allied Medical Sciences, Isra University, Amman, Jordan

*Correspondence: sumaiyah.mat@ukm.edu.my

ABSTRACT

Knee osteoarthritis (KOA) commonly coexists with type 2 diabetes mellitus (T2DM) in older adults, leading to compounded functional limitations, reduced physical activity, and poorer self-management outcomes. Integrated care interventions for this multimorbidity remain limited, particularly in low-resource settings. This study aimed to develop a theory-informed, multifaceted intervention, the Intervention for Older Malaysians Living with Knee Osteoarthritis and Diabetes Mellitus (i-MOAD), to support coordinated, patient-centred self-management in Malaysian primary care. Guided by the UK Medical Research Council framework and the Theoretical Domains Framework, the development process incorporated evidence synthesis, stakeholder consultation, qualitative interviews with older adults, healthcare providers and physiotherapists, mapping to Behaviour Change Techniques, and refinement using Acceptability, Practicality, Effectiveness, Affordability, Safety and Equity (APEASE) criteria. Findings identified key barriers including limited knowledge, low self-efficacy, pain-related avoidance, reduced motivation, limited social support, and structural constraints in primary care delivery. These determinants informed a multidomain intervention comprising KOA and diabetes self-management education, physical activity and exercise guidance, dietary and weight management advice, self-monitoring tools, exercise logs, and optional WhatsApp-based reinforcement. Although this intervention-development study did not generate inferential p-values or confidence intervals, evidence was strengthened through descriptive APEASE stakeholder evaluation. Of 26 stakeholders attending the final meeting, 20 completed the APEASE questionnaire; healthcare providers rated the intervention highly for acceptability (88.9%), perceived effectiveness (88.3%), safety (86.7%), practicability (82.8%), equity (82.2%), and affordability (79.2%). The intervention may be sustainable because it is low-cost, culturally adapted, booklet-based, supported by simple digital tools, and designed for integration within existing primary healthcare workflows. Future research should evaluate feasibility, adherence, implementation outcomes, and preliminary effectiveness in a pilot trial.

Keywords: Complex intervention; feasibility; knee osteoarthritis; type 2 diabetes mellitus

i-031

THABAT: The Home-based Activity for Balance and Training Program to Prevent Falls in Older Jordanians with Knee Osteoarthritis

ANAN AL-SHDIFAT^{1,2}, SUMAIYAH MAT^{1*}, DEEPASHINI HARITHASAN¹,
ALIA ALGHWIRI³, DEVINDER KAUR AJIT SINGH¹

¹Centre for Healthy Ageing and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia

²Department of Rehabilitation Sciences, Faculty of Applied Medical Sciences, Al al-Bayt University, Mafraq, Jordan

³Department of Physiotherapy, School of Rehabilitation Sciences, The University of Jordan, Amman, Jordan

*Correspondence: sumaiyah.mat@ukm.edu.my

ABSTRACT

Falls are a major concern among older adults and are associated with reduced independence, mobility limitations, and poor quality of life. Older adults with knee osteoarthritis are at increased risk of falls due to pain, muscle weakness, impaired balance, and joint stiffness. The Home-based Activity for Balance and Training (THABAT) program was developed through a multi-phase process that included a systematic review and meta-analysis, a cross-sectional study, and qualitative interviews with older Jordanian adults. Findings from these phases were integrated and refined through an expert e-Delphi consensus process, resulting in consensus on the structure, content, and delivery format of a structured home-based intervention booklet. The 12-week feasibility implementation is currently being conducted among 24 older adults aged 60 years and above, randomised into an intervention group and a control group, with 12 participants in each group. The intervention group receives the THABAT booklet, which includes strengthening, balance, and flexibility exercises, together with education on fall prevention, knee pain management, and adaptation of daily activities, while the control group receives conventional physiotherapy exercises for knee osteoarthritis. Sessions are performed two to three times per week for 45–60 minutes, with monthly follow-up to monitor adherence and support participants. Feasibility outcomes include attendance, adherence, and participant satisfaction, while clinical outcomes include fear of falling using the Arabic Falls Efficacy Scale-International, self-reported falls, Timed Up and Go, Two-Minute Walk Test, 30-Second Sit-to-Stand Test and Knee Injury and Osteoarthritis Outcome Score quality of life and activities of daily living subscales. As the study is ongoing, expected outcomes include improved feasibility, acceptability, balance, mobility, physical function, and self-management. THABAT represents an innovative, culturally adapted intervention with potential clinical and community impact for older adults with knee osteoarthritis in Jordan.

Keywords: Falls prevention; feasibility study; home-based exercise; Jordan; knee osteoarthritis

i-032

Hawdy: A Culturally Tailored Video-Based Pelvic Health Innovation for Older Jordanian Women with Urinary Incontinence

ROWA AL MOMANY¹, SUMAIYAH MAT^{1*}, SAAD AL-NASSAN^{2,3},
NORMALA MESBAH¹, DEVINDER KAUR AJIT SINGH¹

¹*Physiotherapy Program and Centre for Healthy Ageing and Wellness, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Department of Physiotherapy, Faculty of Applied Medical Sciences, The Hashemite University, Zarqa 13133, Jordan*

³*Department of Physiotherapy, Faculty of Allied Medical Sciences, Al-Ahliyya Amman University, Amman 19328, Jordan*

**Correspondence: sumaiyah.mat@ukm.edu.my*

ABSTRACT

Urinary incontinence is a prevalent and under-discussed condition among older women, often negatively affecting quality of life, confidence, daily activities, and social participation. In Jordan, help-seeking is frequently delayed due to embarrassment, stigma, limited awareness, and the belief that urinary incontinence is a normal part of aging. Hawdy was developed through three sequential evidence-informed phases: a scoping review of home-based pelvic health interventions, a modified e-Delphi study with Jordanian physiotherapy experts, and focus group discussions with women from the target population. The scoping review identified key components and delivery strategies of effective home-based pelvic health programs and was published in a peer-reviewed journal. The modified e-Delphi study established expert consensus on the essential components of a culturally appropriate home-based rehabilitation program for older Jordanian women with urinary incontinence and was also published in a peer-reviewed journal. Focus group findings further highlighted the importance of privacy, cultural sensitivity, accessibility, simple Arabic communication and self-management support in pelvic health interventions for older women. Hawdy integrates multiple conservative pelvic health components, including pelvic floor muscle training, bladder training strategies, behavioral and lifestyle modification, pelvic health education, breathing and postural exercises, and adherence support. Its home-based delivery allows women to pause, replay and practice the program independently at their own pace, supporting privacy, engagement, accessibility, and long-term self-management. To our knowledge, Hawdy is among the first Arabic video-based pelvic health interventions developed specifically for older women in Jordan using a structured, evidence-informed, and user-centered design process. Hawdy offers a scalable, low-cost, and clinically grounded innovation with potential for wider implementation in community and healthcare settings to improve access to conservative urinary incontinence care, reduce stigma, and empower older women to manage pelvic health confidently at home.

Keywords: Hawdy; Jordan; older women; pelvic health; urinary incontinence; video-based intervention

i-033

Development of a Bio-Based Head Phantom for Dosimetry in Brain Radiotherapy

SITI HAJAR ZUBER^{1*}, MUHAMMAD FAHMI RIZAL ABDUL HADI²,
NURUL HASHIKIN AB. AZIZ², MOHD FAHMI MOHD YUSOF³,
MOHD ZAHRI ABDUL AZIZ⁴

¹*Centre for Diagnostic, Therapeutics and Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*School of Physics, Universiti Sains Malaysia, 11800 Penang, Malaysia*

³*Faculty of Nursing and Health Sciences, Universiti Islam Melaka, 78200, Melaka, Malaysia*

⁴*Pusat Perubatan USM, Universiti Sains Malaysia, 13200 Penang, Malaysia*

**Correspondence: hajarzuber@ukm.edu.my*

ABSTRACT

An innovative bio-based head phantom for brain radiotherapy has been developed using Rhizophora wood bonded with soy flour and lignin, marking a significant advancement over conventional commercial phantoms. This study aims to investigate the integration of two organic adhesives and surface coatings in enhancing the mechanical and dosimetric properties of the phantom across various energy ranges, and to demonstrate its feasibility in clinical dosimetry for brain radiotherapy. To further support the dosimetric evaluation, a dedicated Monte Carlo simulation module using GATE (Geant4 Application for Tomographic Emission) was developed. The GATE module incorporates the detailed geometry, material composition, and density characteristics of the bio-based phantom, allowing for accurate simulation of clinical treatment beams and verification of dose delivery. Technically, the bio-based phantom presents several advantages. It is engineered with a compact, low-gap slab structure, minimising air pockets to ensure dosimetric uniformity. Its density closely replicates that of human cranial tissues, enabling accurate simulation of photon attenuation and scatter, which is critical for optimising treatment planning and delivery. The measured percentage depth dose (PDD) of the head phantom was in agreement with the experimental PDD of water, with overall discrepancies of 0 to 8.7% at depths ranging from 1.0 to 15.0 cm, with the average thermoluminescence dosimeter dose recorded in the planning target volume being 283.13 cGy, which is within $\pm 5.6\%$ of the prescribed dose. Compared to standard commercial phantoms, the proposed bio-based phantom offers a more cost-effective alternative due to the use of readily available renewable materials and simplified fabrication processes, thus demonstrating strong commercialisation potential for oncology and radiotherapy applications. This affordability may benefit hospitals, oncology and radiotherapy units, academic institutions, and research centers with radiation applications. The phantom's lightweight construction also improves ease of handling and positioning during radiotherapy procedures, enhancing workflow efficiency in clinical settings. Beyond its technical merits, the phantom embraces sustainable innovation. The use of renewable, biodegradable materials such as Rhizophora wood, soy flour, and lignin not only

reduces environmental impact but also aligns with eco-conscious practices in medical research and device fabrication. Together with the Monte Carlo GATE module, this work offers a comprehensive solution that supports both clinical accuracy and environmental responsibility, setting a new benchmark in the development of medical phantoms for radiotherapy dosimetry.

Keywords: Bio-based; dosimetry; Monte Carlo; phantom fabrication; radiotherapy

i-034

Enhancing Breast Cancer Screening Awareness through Virtual Reality Simulation

MAZLYFARINA MOHAMAD^{1*}, HARYANI HARUN², NUR DINI HISYAM^{1*}

¹*Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Woodlands Health Campus, 737628 Singapore*

**Correspondence: mazlyfarina@ukm.edu.my*

ABSTRACT

Mammography is a crucial diagnostic tool for breast cancer screening; however, the procedure frequently induces psychological distress, anxiety, and discomfort, largely due to patients' lack of procedural knowledge. To address this educational gap, virtual reality (VR) has emerged as a highly advantageous platform, offering an immersive and safe environment to familiarise women with the screening process. This study investigated the efficacy of VR as a patient education tool, specifically assessing its impact on procedural understanding, anxiety reduction, and overall patient experience. A total of 40 consenting participants engaged in a simulated VR mammography screening session. Pre- and post-intervention surveys utilising a 5-point Likert scale were administered to evaluate shifts in perceived knowledge and satisfaction. Spearman's correlation analysis revealed a highly significant improvement in participants' procedural knowledge following the VR intervention ($p < 0.001$). Crucially, the immersive educational experience not only enhanced comprehension but also successfully alleviated patient anxiety. These findings highlight the distinct advantages of VR as an innovative, patient-centered educational tool that effectively demystifies breast cancer screening and promotes greater awareness and compliance. Commercially, this immersive VR innovation demonstrates strong potential for implementation in hospitals and community-based awareness programs, providing a scalable and user-centered solution to enhance women's experiences during mammography screening.

Keywords: Breast cancer screening; mammography; patient education; virtual reality

i-035

Development of a “Tummy Trail”: A Nutrition Intervention and Comprehensive Feeding Module for Special Needs Children

NUR FATIHAH JAFFRI¹, RAJA NUR SAHIRAH RAJA SULAIMAN¹,
NURUL HIDAYAH NASARUDIN¹, NUR HANA HAMZAID^{2*},
NURUL HIDAYAH MD FADZIL¹, NURUL HUSNA SHUIB³

¹*Department of Dietetics, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Centre for Rehabilitation & Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

³*Department, Yayasan Insan Istimewa Selangor, 40200 Shah Alam, Selangor, Malaysia*

**Correspondence: hanahamzaid@ukm.edu.my*

ABSTRACT

Nutritional challenges among special needs children, including picky eating and selective food behaviors, significantly impair growth, development, and overall quality of life among children with autism spectrum disorder, global developmental delay, Down syndrome, and attention deficit hyperactivity disorder. A needs assessment at a therapy centre, Yayasan Insan Istimewa Selangor, identified major challenges related to limited dietary variety, poor nutritional status, picky eating and selectivity, alongside gaps in knowledge regarding feeding strategies, growth monitoring, and nutrition management. Therefore, an innovative module, Tummy Trail, introduces a comprehensive feeding module and nutrition intervention package designed to improve food acceptance, reduce picky eating behaviors, and enhance nutritional outcomes among children with special needs. This package integrates three synergistic, complementary components that collectively support child engagement, parent education, and intervention guidance. First, an interactive Sensory-Based Food Exploration Game Kit uses storytelling, sensory play, and educational games. Second, Bite by Bite: Food Chaining Diary is an interactive parental guidance diary incorporating food chaining techniques and the sequential oral sensory approach. Third, the Panduan Pemakanan Anak Autisme module provides visual-based nutrition education, nutrition requirement guidance, and evidence-based feeding strategies tackling behavioural feeding issues. The novelty of this innovation lies in its holistic and progressive integration of child-centric playful engagement, evidence-based parental guidance, and specialised support, offering a scalable solution for various early intervention settings. Expected impacts include improved food acceptance, enhanced dietary diversity, better nutritional status, and enhanced caregiver confidence in managing feeding challenges among children with special needs.

Keywords: Comprehensive feeding module; educational games; food chaining; nutritional challenges; picky eating; special needs children

i-036

Malaysia's First Direct Observed Treatment Innovation for Leprosy Patients in Pekan District, Pahang

SURENDER BENEDICT^{1*}, WONG CHIN MUN², MAZLINA MOHTAR³,
MASRI QUAK SAIFUDDIN¹

¹Pekan District Health Office, 26600 Pekan, Pahang, Malaysia

²Pahang State Health Department, Bandar Indera Mahkota, 25582 Kuantan, Pahang, Malaysia

³Hospital Sultan Haji Ahmad Shah, 28000 Temerloh, Pahang, Malaysia

*Correspondence: suryender1@gmail.com

ABSTRACT

Leprosy is a chronic infectious disease caused by *Mycobacterium leprae*. It remains a challenge to eliminate local transmission due to poor treatment adherence within a long duration, poor understanding of infection, stigma, and lack of treatment facilities in remote areas. Incomplete treatment leads to Grade 2 deformity. Challenges in confirming treatment success among leprosy patients highlighted the need for a structured treatment-monitoring system in Pekan district. Pekan District Health Office implemented the direct observed treatment (DOT) approach used in tuberculosis (TB) patients for leprosy since 2023, following the directive from the Pahang State Health Department. This innovation is the first prototype in Malaysia used for defining leprosy treatment success, a revolution in leprosy disease management. It aims to ensure successful treatment outcomes and enable early detection of possible medication side effects. Treatment adherence monitoring was implemented through a structured book modified from TBIS10E, called LIS10E. Leprosy patients were monitored via DOT daily by identified social leaders/caretakers, weekly by community nurses in health clinics, and monthly by the medical team in Treatment Centers. Milestones of treatment adherence are rewarded with incentives. This innovation achieved a 95.8% treatment adherence success rate in 2024–2025. The success was evaluated through clinical improvement, reduction in the number of defaulters, negative slit skin smear repeated results, and leprosy incidence reduction among contacts in endemic areas. Operational challenges include logistical barriers in accessing remote areas for weekly follow-up by healthcare personnel and cooperation from stakeholders. The innovation demonstrated treatment success for leprosy, while preventing complications such as medication allergy, Grade 2 deformities, and disease transmission. An estimated RM4,000,000 could potentially be saved annually from workforce productivity loss and job incapacity stemming from delayed diagnosis/incomplete treatment, projected based on 200 new cases reported yearly in Pahang. Therefore, the program is recommended for nationwide expansion due to its effectiveness, sustainability, cost-efficiency, and high treatment-adherence success rate.

Keywords: Direct observed treatment; leprosy; *Mycobacterium leprae*; treatment adherence

i-037

Mobile Intravenous Administration Platform: Redefining Patient Mobility and Safety in Intravenous Therapy

SYAFIQ YUSUF*, RAJA MUHAMMAD 'IFFAT SYAMIM RAJA AHMAD
SHALABY, SYED AHMAD YUSOF SYED MUHAMMAD, NORAZALINA
MOHAMAD ZAH, IZWAK IKMAL ISHAK

KPJ Healthcare University, Kota Seriemas, 71800 Nilai, Negeri Sembilan, Malaysia

**Correspondence: syafiqapeyy@gmail.com*

ABSTRACT

As standard practice, conventional IV poles are used everywhere, with many known flaws such as compromising comfort and increasing the risks of falls and treatment disruption. There exists a desire for higher quality care regarding IV therapy, especially for ambulatory patients traveling to and from healthcare facilities in discomfort. In hopes of developing a Mobile Intravenous Administration Platform (MIVAP) with the aid of a small liquid pump to enhance mobility over a duration of 20 minutes, meanwhile not sacrificing the existing benefits of current IV practices. A patient-first thought process was implemented, focusing on practical use, ergonomic design, and constant refinement. The device was developed using a vest, durable pockets attached with adjustable straps, and quality handwork to securely hold standard IV bags (300–500 mL). The design was kept familiar to ensure compatibility across diverse healthcare settings, including wards, outpatient care, and home-based therapy, to reduce rejection of new systems. The innovation's novelty lies in replacing the traditional stationary IV pole with a wearable vest with an onboard pump that increases flow rate. This design allows for the independence of the patient without risking therapy. By enabling hands-free mobility, the device reduces fall risks, minimises IV line dislodgement, enhances patient dignity, and allows for patient freedom of movement. For healthcare workers such as nurses, they no longer need to accompany patients during patient visits within hospital premises. Upon completion, expect improved patient safety, faster recovery through increased mobility, and increased patient independence with this project. With an estimated production cost of RM70 per unit, the solution is highly cost-effective and scalable. Its commercialisation potential is strong, supported by dual business models (direct distribution and subscription rental), targeting hospitals, clinics, and home care markets. This project aims to be a precedent for novel patient-centric IV therapy.

Keywords: Intravenous therapy; medical device innovation; patient mobility; patient safety; wearable IV system

i-038

Translating *Alocasia denudata* Engler into a Commercial Botanical Platform for Advanced Diabetic Wound Care

MAZLYZAM ABDUL LATIF^{1,2*}, ASMAH HAMID^{1,2}, MOHD ZULASYRAF
MOHD ZAKI¹, SITI AISYAH ARSHAD¹, FATINI NADIAH MOHAMMAD
ZAMZAMI¹

¹Center for Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia,
50300 Kuala Lumpur, Malaysia

²Biomedical Science Program, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala
Lumpur, Malaysia

*Correspondence: mazlyzam@ukm.edu.my

ABSTRACT

The escalating economic burden of chronic diabetic foot ulcers necessitates scalable, advanced therapeutic solutions. We present a novel therapeutic gel derived from *Alocasia denudata* Engler, engineered to transition from a naturally occurring botanical into a highly marketable, evidence-based biomedical product targeting the lucrative clinical wound care sector. The formulation's efficacy was validated through dual-model testing. In vitro assays utilising human dermal fibroblasts (HDF-a) assessed cell viability via 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assays, alongside scratch migration assays and RT-PCR gene expression profiling. In vivo wound healing profiles were evaluated using 150 diabetic-induced male Wistar rats randomly distributed into five treatment groups (n = 30/group). Six full-thickness dorsal excisional wounds (6 mm diameter) were created per animal. At strategic intervals (Days 1, 3, 6, 10, and 14), six rats per group were sacrificed (n = 6/interval) to monitor macroscopic contraction, tissue biochemistry (total protein, cathepsin B, uronic acid, hexosamine), and histology. Our proprietary formulation demonstrated remarkable therapeutic potential. In vitro human cell models confirmed the extract is entirely non-cytotoxic while significantly upregulating core regenerative genes (TGF- β , PDGF- β , VEGF- α , COL-III, COL-IV). This multi-target efficacy is driven by a standardised profile of bioactive compounds, predominantly 9-octadecanamide (Z)- and 9,12-octadecadienoic acid (Z,Z)-methyl ester, tumerone, ar-tumerone and curlone, which were identified by GC-MS, alongside high potassium (19.84%), which is essential for tissue repair. Crucially, in vivo trials of the 10% topical gel achieved rapid wound closure, robust collagen deposition, and profound anti-inflammatory action via Cathepsin B suppression. Moving beyond traditional alternative medicine, this research establishes a fully validated botanical platform. The *A. denudata* 10% gel offers a highly efficacious, biochemically defined, and cost-effective therapeutic solution optimised for commercial product development.

Keywords: *Alocasia denudata* Engler; bioactive compounds; diabetic wound healing; topical therapeutics; translational biomedicine

i-039

Translating Traditional Herbs to Modern Therapeutics: Multi-Targeted Neuroprotective Efficacy of *Zingiber zerumbet* against Alzheimer's Disease

ASMAH HAMID^{1*}, MAZLYZAM ABDUL LATIF¹, FARAH SYAHIBAH
MOHD HARIRI¹, NURUL IZZATY MOHD AZIZNUDIN¹, KHAIRANA
HUSAIN²

¹Center for Toxicology & Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia,
50300 Kuala Lumpur, Malaysia

²Centre for Drug and Herbal Development, Faculty of Pharmacy, Universiti Kebangsaan Malaysia, 50300
Kuala Lumpur, Malaysia

*Correspondence: asmah0901@ukm.edu.my

ABSTRACT

Alzheimer's disease (AD) is a progressive, incurable neurodegenerative disorder driven by abnormal beta-amyloid (A β) deposition, tau hyperphosphorylation, and neuronal death. Aluminium chloride (AlCl₃) serves as a significant environmental risk factor that accelerates cognitive decline by inducing severe oxidative stress. With the alarming global rise in neurodegenerative conditions, exploring natural products for neuroprotection is vital. This study highlights the therapeutic potential of *Zingiber zerumbet* (Lempoyang) ethyl acetate extract and its bioactive sesquiterpene, zerumbone (ZER), as multi-targeted botanical agents. Integrating in vitro (H19-7 hippocampus cells) and in vivo (Wistar rats) models, our combined results demonstrate robust, systemic efficacy against AlCl₃-induced neurotoxicity. In vitro, cells pre-treated with ZER (IC₁₀ = 1.26 μ g/mL) and *Z. zerumbet* extract (IC₁₀ = 17.10 μ g/mL) exhibited a statistically significant reduction in lipid peroxidation ($p < 0.05$), protein carbonyls, and cellular DNA damage ($p < 0.05$). Furthermore, both interventions significantly enhanced antioxidant defenses through increased superoxide dismutase and glutathione (GSH) activities while reducing A β accumulation and other neurodegeneration-associated biomarkers ($p < 0.05$). In vivo, AlCl₃ administration (17–25 mg/kg) triggered profound memory impairment and accelerated neuronal death. Treatment with *Z. zerumbet* extract (200 and 400 mg/kg/day) and pure ZER (10, 20 and 30 mg/kg) effectively reversed these deficits. Behaviorally, these biochemical restorations translated to the successful prevention of neuronal cell death and the complete rescue of spatial working memory in the Y-Maze and restored recognition memory in the Novel Object Recognition (NOR) test, without impairing baseline locomotor activity. Notably, proactive protective dosing regimens yielded statistically superior cognitive and biochemical outcomes ($p < 0.01$) compared to delayed therapeutic regimens. In conclusion, these synergistic statistical findings firmly establish *Z. zerumbet* and ZER as highly promising natural products. Their potent ability to prevent heavy-metal-induced cognitive impairment underscores their strong commercial potential for development into scalable, high-value nutraceuticals and targeted botanical drugs for AD management.

Keywords: Alzheimer's disease; neuroprotection; therapeutic; Zerumbone; *Zingiber zerumbet*

i-104

From Play to Practice: Effectiveness of Portion Masters™ in Improving Portion Size Knowledge and Attitude in Overweight and Obese Youths

YEE XING YOU^{1*}, CHING YEE FOO², SEW LI CHEW²

¹*Dietetic Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Osentai Enterprise, L9-19, The Holmes, Bandar Tun Razak, 56000 Kuala Lumpur*

**Correspondence: youyeexing@ukm.edu.my*

ABSTRACT

Obesity among Malaysian youths is on the rise due to unhealthy eating habits and poor understanding of appropriate portion sizes. Interactive tools such as Portion Masters™, a board game designed to teach correct portion sizes, offer a novel approach to address this issue. This quasi-experimental study aimed to evaluate the acceptance and effectiveness of the Portion Masters™ game in improving knowledge, attitude, and practice related to portion sizes among overweight and obese youths in Klang Valley. A total of 40 participants aged 18–30 years with BMI >23 kg/m² were recruited using purposive sampling. Participants completed validated pre- and post-intervention questionnaires assessing knowledge, attitude, and practice. The Portion Masters™ game was conducted in small group sessions. A post-test assessing knowledge was administered immediately after the game, while a follow-up post-test evaluating attitudes and practices was conducted one month later. Data was analysed using SPSS version 29.0. Participants showed significant improvement in knowledge ($24 \pm 15.5\%$ to $84 \pm 8.84\%$, $p < 0.001$) and attitude (33.1 ± 4.4 to 37.7 ± 2.8 , $p < 0.05$) scores. However, the change in practice scores was not statistically significant (2.58 ± 1.5 to 2.85 ± 1.7 , $p = 0.112$). Most participants positively accepted the game and supported its use as an educational tool. Portion Masters™ is an effective and engaging tool for improving knowledge and attitudes about portion sizes among youths. Beyond its educational impact, the game demonstrates strong translational and commercialisation potential. The product has been developed in collaboration with an industry partner, Osentai Enterprise, with ongoing efforts in product refinement and market readiness. Intellectual property considerations, including copyright and trademark protection for Portion Masters™, have been registered to support future licensing opportunities. The board game is low-cost, scalable, and adaptable across schools, community programmes, and clinical settings, with potential for broader industry adoption and integration into national health promotion initiatives. The product aligns with global priorities on obesity prevention and supports sustainable health promotion strategies.

Keywords: Attitude; educational tool; knowledge; nutrition; Portion Masters™

i-105

Beyond Geography: Immersive Virtual Reality Nature for Mental Well-Being in Medical Students

MUHAMMAD HIZRI HATTA¹, FARAH DEENA ABDUL SAMAD^{1,2*},
SIEW KOON CHONG⁴, SURIATI MOHAMED SAINI^{2,3}

¹Department of Psychiatry, Faculty of Medicine, Universiti Sultan Zainal Abidin, 21300 Terengganu, Malaysia

²Department of Psychiatry, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia

³Hospital Canselor Tuanku Muhriz, 56000 Cheras, Kuala Lumpur, Malaysia

⁴Department of Psychiatry, Hospital Kuala Lumpur, 50586 Kuala Lumpur, Malaysia

*Correspondence: drfarahdeena@ukm.edu.my

ABSTRACT

Medical students experience high levels of psychological distress, highlighting the need for accessible and scalable mental health interventions. Immersive 360-degree virtual reality (VR) nature exposure has emerged as a promising strategy; however, the influence of geographical familiarity on its therapeutic effectiveness remains unclear. This study aimed to compare the efficacy of local (Malaysian) versus overseas (Western European) virtual natural environments on mental well-being among medical students. A randomised trial was conducted among 84 senior medical students equally allocated into two groups viewing either local or overseas VR environments. Participants underwent two 15-minute VR sessions over two weeks featuring forest, beach, river, and park scenes. Psychological outcomes were assessed using the Depression, Anxiety and Stress Scale-21 and the World Health Organisation-5 Well-Being Index at baseline, post-intervention one, and post-intervention two. Data were analysed using repeated-measures analysis of variance. The intervention demonstrated significant reductions in anxiety ($p < 0.001$) and stress ($p < 0.001$), alongside improved well-being ($p < 0.001$), with no significant change in depression ($p = 0.122$). Importantly, no significant differences were observed between local and overseas groups across all outcomes ($p > 0.05$). The novelty of this study lies in demonstrating that the psychological benefits of VR nature exposure are independent of geographical familiarity, suggesting a universal mechanism of psychological restoration. These findings support VR nature intervention as a scalable, low-resource strategy for improving mental well-being. The commercialisation potential includes implementation in educational institutions, healthcare settings, and digital mental health platforms without the need for costly localised content production.

Keywords: 360-degree VR video; geographic comparison; medical student; mental wellness; virtual reality

i-106

MyRAKAN Autism: Innovating Social Skills Training through Video Modelling and Virtual Reality

**WAN SALWINA WAN ISMAIL^{1,5,6*}, NORAZLIN KAMAL NOR^{2,6},
NOORLAILI MOHD TOHIT^{3,5}, FARAH DEENA ABDUL SAMAD^{1,5}, MOHD
AKIF FARHAN AHMAD BASRI⁴**

¹*Department of Psychiatry, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia*

²*Department of Paediatrics, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia*

³*Department of Family Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000 Cheras, Kuala Lumpur, Malaysia*

⁴*Department of Psychology, School of Liberal Arts and Sciences, Faculty of Social Sciences and Leisure Management, Taylor's University, 47500 Subang Jaya, Selangor, Malaysia*

⁵*Hospital Canselor Tuanku Muhriz, Universiti Kebangsaan Malaysia 56000 Cheras, Kuala Lumpur, Malaysia*

⁶*Hospital Tunku Ampuan Besar Tuanku Aishah Rohani, Hospital Pakar Kanak-Kanak UKM, 56000 Cheras, Kuala Lumpur, Malaysia*

**Correspondence: wan@hctm.ukm.edu.my*

ABSTRACT

Autism spectrum disorder (ASD) is a neurodevelopmental condition that impacts social interaction and communication. Given that ASD has no cure, psychosocial interventions are essential to improve social skills, particularly for youths with high-functioning ASD (HFASD). In Malaysia, limited access to trained therapists underscores the need for innovative adjunct interventions. This study aimed to develop and evaluate MyRAKAN Autism, a social skills intervention for youths with HFASD, using a mixed-methods approach. Phase 1 explored social skill deficits and cognitive behavioural therapy techniques through qualitative analysis. Phase 2 established content validity via the Fuzzy Delphi method to achieve expert consensus. The intervention integrates psychoeducation, modelling, relaxation, and skills rehearsal, delivered through video modelling (VM) and spherical video-based virtual reality (SVVR). Phase 3 involved formative evaluation with HFASD youths and experts, indicating good feasibility and acceptability. Phase 4 evaluated efficacy through a randomised controlled trial involving 52 participants (aged 18–24), assigned to either a treatment or delayed-treatment control group. Results showed significant improvements in the treatment group in social responsiveness ($p = 0.046$; between-group $p = 0.001$), communication ($p = 0.038$; between-group $p = 0.001$), and imagination ($p = 0.048$; between-group $p = 0.048$). Significant between-group effects were also observed for social awareness, social cognition, and quality of life. Overall, these findings suggest that MyRAKAN Autism is a feasible, acceptable, and effective social skills intervention for youths with HFASD. It has secured intellectual property protection and is currently

positioned for industry collaboration with apoQlar Medical to enhance functionality, support clinical implementation, and advance towards commercialisation as an adjunct intervention in social skills training for youths with HFASD.

Keywords: High-functioning autism spectrum disorder; quality of life; social skills

i-107

Redefining Denture Clasp Design: A Novel Gradient Flexibility Approach Using 3D-Printed Flexible Resin

AIEMEEZA RAJALI^{1*}, ROHANA AHMAD¹, SITI MARIAM AB GHANI¹,
SOLEHUDDIN SHUIB², AMMAR YASSER ABDUL HAKIM³

¹Faculty of Dentistry, Universiti Teknologi MARA, 47000 Sungai Buloh, Selangor, Malaysia

²Faculty of Mechanical Engineering, Universiti Teknologi MARA, 42300 Puncak Alam, Selangor, Malaysia

³Faculty of Dentistry, Universiti Sains Islam Malaysia, 56100 Ampang, Malaysia

*Correspondence: aiemeeza7738@uitm.edu.my

ABSTRACT

Conventional removable partial denture clasps are designed for rigid metal materials, often leading to poor aesthetics and excessive stiffness. With the rise of 3D printing, flexible denture resin offers new possibilities; however, material-specific design principles remain limited. This study aims to develop a material-specific digital design system incorporating tapered geometry for clasps tailored to the mechanical behaviour of 3D-printed flexible denture resin. A computer-aided design (CAD) workflow was established to generate four clasp configurations based on retentive arm position relative to the survey line and controlled undercut engagement. A tapered width profile was introduced, with shoulder width set at 3 mm for premolars and 4 mm for molars, gradually reducing to 1.5 mm at the retentive tip. Clasp thickness was maintained at a constant 1.5 mm. Designs were printed using 3D-printed flexible denture resin. The novelty of this innovation lies in a material-specific clasp design that introduces a tapered width geometry to create a controlled stiffness gradient along the clasp, tailored to the mechanical behaviour of 3D-printed flexible denture resin. The design optimises load distribution through controlled gradient flexibility, reducing stress on abutment teeth while maintaining retention. This enhances patient comfort, prolongs prosthesis longevity, and improves aesthetics with a metal-free solution. Designed for immediate translation, this innovation can be embedded into existing CAD systems as a standardised digital protocol, enabling scalable, high-throughput production of next-generation denture clasps with consistent precision across clinical and laboratory environments.

Keywords: 3D-printed flexible denture resin; clasp design; digital dental design; stiffness gradient; tapered geometry

Digital Smart Health Hub

AIN SOFIYA INSYIROH^{1*}, PUTERI NUR FARZANA¹, NURUL NADIAH FARHANAH², NURUL AIN SYAFIQAH³, NURUL IZZAH INSYIRAH⁴

¹*Optometry Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

²*Pharmacy Programme, Faculty of Pharmacy, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

³*Dietetics Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

⁴*Nutrition Science Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia*

**Correspondence: a212157@siswa.ukm.edu.my*

ABSTRACT

Social media often spreads misleading health advice, highlighting the need for trustworthy sources. This application introduces an integrated Home Medication Review (HMR) model that unites pharmacists and dietitians to deliver comprehensive and patient-centered care. The project follows four qualitative phases, including semi-structured interviews with healthcare professionals and patients, to identify post-discharge challenges. This app development focused on intuitive UI/UX and expert validation for safety and accuracy. A key innovation lies in combining patient autonomy with digital health technology, streamlining the Home Medication Review process in a proactive and user-driven manner. The application features a medical information search engine curated by licensed practitioners to minimise misinformation and unsafe self-medication. Core components include post-discharge monitoring, risk stratification, and nutrition management, as well as a scanning and upload feature to digitise prescribed medication for easy reference, reducing errors and delays. The application demonstrates strong commercial potential by targeting post-discharge patients, elderly individuals with polypharmacy, and caregivers, with revenue generated through subscriptions and partnerships with healthcare providers. Supported by licensed practitioners, this market-ready solution is practical and reliable for multidisciplinary healthcare settings.

Keywords: Digital technology; home medication review; integrated model; professional healthcare